# A study to assess the feasibility and effects of introducing remote ischaemic preconditioning in elective abdominal aorta aneurysm repair

Submission date	Recruitment status No longer recruiting	<ul><li>Prospectively registered</li><li>Protocol</li></ul>	
16/03/2012			
Registration date	Overall study status	Statistical analysis plan	
16/03/2012	Completed	[X] Results	
Last Edited	Condition category	[] Individual participant data	
28/08/2015	Circulatory System		

#### Plain English summary of protocol

Background and study aims

An abdominal aneurysm is an enlarged and weakened section of the abdominal aorta - the main artery that carries blood from the heart. As an aneurysm increases in size it can split open, which carries a very high risk of death. Therefore, when it reaches a certain size it is best to repair it surgically before it becomes dangerous. During this operation the surgeon replaces the weak section in the aorta with a graft. To do this the surgeon will clamp the artery above and below the aneurysm to prevent blood flowing through the area during surgery. This situation, when blood flow to an organ is cut off, is called ischaemia. When the blood supply to an organ returns after a period of ischaemia (a process called reperfusion), damage can occur. The absence of oxygen and nutrients during ischaemia creates a condition in which the restoration of the blood supply results in inflammation and damage to tissues. This can lead to heart and kidney problems and even death. However, we know that if the blood supply is first halted for a short time and then restored quite quickly the tissue cells seem to learn to deal with later, longer periods of loss of blood flow in a process called ischaemic preconditioning. Studies suggest that ischaemic preconditioning may protect the heart and other organs during big operations. In this study we wish to investigate if remote ischaemic preconditioning (done in a simple and safe way by inflating a normal blood pressure cuff on the patient's arm for 5 minutes) protects the patient and reduces heart and kidney problems after abdominal aneurysm operations. The aim is to assess the feasibility and consequences of remote ischaemic preconditioning and the value of conducting a larger study to accurately assess its full usefulness as part of routine care.

#### Who can participate?

Patients aged 30 to 100 undergoing elective abdominal aorta aneurysm repair.

#### What does the study involve?

Before undergoing surgery participants were randomly allocated to receive either remote ischaemic preconditioning or a sham procedure. Remote ischaemic preconditioning was applied by inflating a blood pressure cuff on the arm of the patient for 3 cycles of 5 minutes, allowing 5 minutes of reperfusion in between.

What are the possible benefits and risks of participating? Not provided at time of registration.

Where is the study run from? Southmead Hospital (UK).

When is the study starting and how long is it expected to run for? December 2010 to December 2012.

Who is funding the study? National Institute of Health Research (NIHR) (UK).

Who is the main contact?
Dr Ronelle Mouton
Ronelle.Mouton@nbt.nhs.uk

#### Contact information

#### Type(s)

Scientific

#### Contact name

Dr Ronelle Mouton

#### Contact details

Dept of Anaesthesia Southmead Hospital Southmead Road Westbury-On-Trym Bristol United Kingdom BS10 5NB

Ronelle.Mouton@nbt.nhs.uk

#### Additional identifiers

Protocol serial number 9300

## Study information

#### Scientific Title

A study to assess the feasibility and effects of introducing remote ischaemic preconditioning in elective abdominal aorta aneurysm repair: a randomised controlled pilot study

#### Study objectives

In this study we are piloting the introduction of remote ischaemic preconditioning (RIPC) in patients undergoing elective surgery for repair of abdominal aortic aneurysm (AAA), using a simple, non-invasive, non-surgical technique. We wish to determine the feasibility and

acceptability of the procedure and to inform a proposal to conduct an adequately powered, multi-centre Phase III randomised controlled trial (RCT).

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

South West 5 REC, 29/07/2010, ref: 10/H0107/36

#### Study design

Randomised; Interventional; Design type: Not specified

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Topic: Cardiovascular; Subtopic: Cardiovascular (all Subtopics); Disease: Valve Disease

#### **Interventions**

60 subjects: 30 undergoing elective abdominal aneurysm repair via the traditional open procedure (15 in RIPC arm and 15 controls)

30 undergoing endovascular aorta aneurysm repair (EVAR) with again 15 patients per arm for RIPC and controls

Remote ischaemic preconditioning applied by inflating a blood pressure cuff on the arm of the patient to 40mmHg above normal systolic blood pressure. The intervention is applied for 3 cycles of 5 mins, allowing 5 mins of reperfusion inbetween.

#### Intervention Type

Procedure/Surgery

#### Primary outcome(s)

The main outcomes of this pilot trial relate to the planning requirements for the multi-centre RCT, including key information on recruitment method and consent procedures and rates, patient and clinical staff acceptability and opinion, data on the logistics of the delivering the RIPC, blinding, data collection and capture, management processes, and expected variations in practice across sites.

Primary clinical outcomes recorded in the pilot trial are renal injury and myocardial injury, including arrhythmias and myocardial infarction.

#### Key secondary outcome(s))

- 1. The length of ICU stay and duration of overall postoperative hospital stay
- 2. Incidence of stroke and death
- 3. Variation in patient-reported outcomes

We shall use variation in these outcomes to inform power calculations for the larger RCT, taking information on procedural differences between centres into consideration.

#### Completion date

31/12/2012

# **Eligibility**

#### Key inclusion criteria

- 1. Over a 24-month period all new patients presenting to Southmead Hospital for elective AAA repair will be invited to participate in the trial
- 2. Upper Age Limit 100 years; Lower Age Limit 30 years

#### Participant type(s)

**Patient** 

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Sex

All

#### Key exclusion criteria

- 1. Unable to give informed consent
- 2. If they are taking sulphonylurea oral hypoglycaemic drugs or nicorandil, which have been shown to influence preconditioning

#### Date of first enrolment

01/12/2010

#### Date of final enrolment

31/12/2012

#### Locations

#### Countries of recruitment

United Kingdom

England

# Study participating centre Southmead Hospital

Bristol United Kingdom BS10 5NB

# Sponsor information

#### Organisation

North Bristol NHS Trust (UK)

#### ROR

https://ror.org/036x6gt55

# Funder(s)

#### Funder type

Government

#### **Funder Name**

National Institute of Health Research (NIHR) (UK) - Research for Patient Benefit Programme (RfPB)

### **Results and Publications**

Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Not provided at time of registration

#### **Study outputs**

Output type	<b>Details</b> results	Date created Date added Peer reviewed? Patient-facing?		
Results article		25/08/2015	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	5 No	Yes