

Effect of high-flow nasal cannula on functional exercise capacity in hospitalized adult lung transplant candidates: a modified cross-over study

Submission date 17/01/2025	Recruitment status Recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 20/01/2025	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 20/01/2025	Condition category Respiratory	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

This study aims to find out whether a high-flow nasal cannula can increase the meters walked in the 6-minute walk test (6MWT) compared to conventional oxygen therapy, and also whether it improves exercise tolerance and reduces dyspnea (shortness of breath) and recovery times in patients with chronic respiratory disease.

Who can participate?

Hospitalized adults (18 years of age or older) with pulmonary hypertension or pulmonary fibrosis who are candidates for lung transplantation

What does the study involve?

Participants will perform four 6MWTs: one with supplemental oxygen delivered by reservoir mask and three with high-flow nasal cannula at 40, 50 or 60 LPM. The researchers will evaluate 6MWT performance and physiological response to exercise.

What are the possible benefits and risks of participating?

If a high-flow nasal cannula improves performance and provides an adequate physiological response in the 6MWT, it would support using this device for better results in pulmonary rehabilitation programs in hospitalized patients. However, during the performance of the 6MWT, some adverse events may occur, such as falls, syncope (fainting), or increased oxygen requirements or respiratory support.

Where is the study run from?

Instituto Nacional del Tórax (Chile)

When is the study starting and how long is it expected to run for?

June 2024 to December 2026

Who is funding the study?
Instituto Nacional del Tórax (Chile)

Who is the main contact?
Ruvistay Gutierrez-Arias, rgutierrez@torax.cl

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Prof Ruvistay Gutierrez-Arias

ORCID ID

<https://orcid.org/0000-0003-1881-9316>

Contact details

José Manuel Infante #717

Santiago

Chile

7500000

+56 (0)2 25755000

rgutierrez@torax.cl

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

CECSSMOriente060824-1

Study information

Scientific Title

Effect of high-flow nasal cannula on the 6-minute walk test in hospitalized patients with pulmonary hypertension or pulmonary fibrosis who are candidates for lung transplantation: a modified cross-over study

Acronym

HELCOS

Study objectives

It is hypothesised that using high-flow nasal cannula with flow levels of 40, 50, and 60 LPM increases the metres walked in the 6-minute walk test (6MWT) compared to conventional

oxygen therapy. In addition, it improves exercise tolerance and reduces dyspnoea levels and recovery times in hospitalised patients with pulmonary hypertension or pulmonary fibrosis who are candidates for lung transplantation. Performance on the 6MWT and physiological response to exercise should improve the higher the flows used.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 06/08/2024, Scientific Ethics Committee of the Servicio de Salud Metropolitano Oriente (Santiago, Chile). (Avenida Salvador #364, Santiago, 7500000, Chile; +56 (0)9 9832 8739, +56 (0)9 8156 6931; comite@cec-ssmoriente-adultos.cl), ref: CECSSMOriente060824-1

Study design

Modified randomized cross-over study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Hospitalised adults with pulmonary hypertension or pulmonary fibrosis who are candidates for lung transplantation

Interventions

The study participants will perform four 6MWTs: one with supplemental oxygen delivered via reservoir mask and three with high-flow nasal cannula at 40, 50 or 60 LPM. An inspired fraction of oxygen will be delivered during each test to maintain oxygen saturation above 90-92%. The temperature will be adjusted to 31-37°C in the high-flow nasal cannula according to patient-reported comfort. The order of performance of the four 6MWTs will be randomized for each patient. The tests will be performed at an interval of 3 days. On the day the tests are conducted there will be no physical training sessions before the evaluations. In addition, during rest days, patients will undergo low-intensity physical rehabilitation.

A computer program will generate the randomization sequence for the order in which the 6MWTs are performed (randomized.org). The researcher who recruits the participants and performs the evaluations will not learn about this assignment until the first test is administered.

Intervention Type

Other

Primary outcome(s)

Functional capacity to perform exercise: the distance walked in each 6MWT will be measured. nt. The four tests will be performed at an interval of 3 days.

Key secondary outcome(s)

1. Physiological response to physical exertion:

1.1. Respiratory rate will be monitored with a multiparameter transport monitor every minute during all 6MWT and at 1, 3, 6, and 10 minutes after the end of the tests.

- 1.2. Heart rate will be monitored with a multiparameter transport monitor every minute during all 6MWT and at 1, 3, 6, and 10 minutes after the end of the tests.
- 1.3. Peripheral oxygen saturation will be monitored with a multiparameter transport monitor every minute during all 6MWT and at 1, 3, 6, and 10 minutes after the end of the tests.
- 1.4. Lower extremity fatigue will be assessed with a simple numerical scale (NRS) from 0 to 10 every minute during all 6MWT and at 1, 3, 6, and 10 minutes after the end of the tests.
- 1.5. Dyspnea will be assessed with the modified Borg scale every minute during all 6MWT and at 1, 3, 6, and 10 minutes after the end of the tests.
2. Stops during the 6MWT: stops during each test will be counted, and the patient will be asked for the reasons.
3. Walking time: Patients' total walking time will be determined. This time will be calculated by subtracting the time spent stopped from the total time of the 6MWT (6 minutes).
4. Time to return to calm: when vital signs, lower extremity fatigue and dyspnea return to baseline values will be determined. This will be assessed up to minute 10 of each 6MWT.
5. Adverse events: adverse events directly associated with the performance of the 6MWT will be recorded. Falls, syncope, and increases in oxygen requirements and respiratory support during the same day of the four 6MWTs will be considered. Deaths and the need for invasive mechanical ventilation or extracorporeal circulatory support will also be considered.

Completion date

01/12/2026

Eligibility

Key inclusion criteria

1. Adult persons (18 years of age or older)
2. Persons with a group I or III pulmonary hypertension diagnosis as defined by the World Health Organization (WHO). Diagnosis can be made by echocardiogram or cardiac catheterization
3. Persons with a diagnosis of pulmonary fibrosis. Diagnosis may be made by computed tomography (CT) or lung biopsy
4. Persons under study to enter the lung transplant list or enlisted at the time of study execution
5. Stable medical condition, defined as no change in medication in the last 14 days before assessment
6. Indication from the treating medical team to perform out-of-bed motor rehabilitation
7. Having previously undergone a 6MWT

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

70 years

Sex

All

Key exclusion criteria

1. Persons with exacerbation of pathology due to infectious, inflammatory process or disease progression
2. Persons with a resting heart rate greater than 120 beats per minute
3. Persons with unstable hemodynamics (MAP <60 mmHg or >100 mmHg)
4. Persons connected to a high-flow nasal cannula with flows equal to or greater than 40 LPM at rest
5. Persons in need of continuous or intermittent connection to noninvasive mechanical ventilation
6. Persons with neuromusculoskeletal alterations or need for technical aids that limit independent walking
7. Persons who do not understand the instructions necessary to perform the 6MWT
8. Persons who do not sign the informed consent

Date of first enrolment

15/08/2024

Date of final enrolment

15/08/2026

Locations**Countries of recruitment**

Chile

Study participating centre

Instituto Nacional del Tórax

José Manuel Infante #717

Santiago

Chile

7500000

Sponsor information**Organisation**

Instituto Nacional del Tórax

ROR

<https://ror.org/00t0z3q71>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Instituto Nacional del Tórax

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Ruvistay Gutierrez-Arias (rgutierrez@torax.cl).

The type of data that will be shared: anonymized Excel file.

Dates of availability: 5 years after the study is published.

Whether consent from participants was required and obtained: yes. All study participants will sign a consent form.

Comments on data anonymization: the data will be stored anonymously in an Excel spreadsheet.

Any ethical or legal restrictions: the data may be presented with the authorization of the study's corresponding author.

Any additional comments: the data may be used for meta-analysis of individual patient data with the prior authorization of the study's corresponding author.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes