# Valdoxan ImproVes depRession with anxiEty symptoms (VIVRE)

Submission date	Recruitment status	<ul><li>Prospectively registered</li></ul>
10/05/2012	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
19/06/2012	Completed	[X] Results
Last Edited	Condition category	Individual participant data
11/09/2019	Mental and Behavioural Disorders	

### Plain English summary of protocol

Background and study aims

Depression is currently one of the most frequent psychiatric diseases. Patients with depression suffer very much of depressed mood, loss of interest or pleasure in activity, and they are often impaired emotionally and socially in their daily functioning. Depressed patients also often complain about symptoms of anxiety in combination with their symptoms of depression. The aim of this study is to observe the effects of Valdoxan®, a drug that acts as an antidepressant and specifically resynchronizes circadian rhythms, such as diurnal variation of the mood, disrupted daily rhythms and impaired daytime alertness, as well as disturbed sleep/wake-rhythms. Besides that, the effect on social and emotional functioning is observed, and the effect on anxiety symptoms in combination with depression. We also intend to assess how safe and tolerated this drug is.

### Who can participate?

Ambulatory (who can walk) adult patients (no age and gender limitation) with major depression, who have been prescribed Valdoxan® before entering the study.

### What does the study involve?

All patients observed in this study will be treated with Valdoxan®. The doctor will keep a record of the baseline status of the patient, including other diseases and other medications. The doctor will document the patient and control the symptoms of depression within the further course of treatment. The follow-up visits, according to his clinical routine, will take place after about three weeks, after about six weeks and a last control after about 12 weeks. During these visits a routine practice investigation will be carried out and the doctor will ask about the symptoms of depression and complete a case report form regarding the severity of depression. The patients will be asked to fill out the brief patients' questionnaire for orientation concerning the severity of depressive symptoms (adapted CGI) and social and emotional impairment (Sheehan disability scale) at each visit. According to the antidepressant effect, the dosage of Valdoxan® can be adapted at first follow-up visit (after about week 3) or afterwards in the course of treatment. The duration of the observation of the patients will be three months.

What are the possible benefits and risks of participating?

There are no particular benefits or risks. The treatment that the patient receives corresponds to

the daily routine of the doctor. The doctor prescribes the drug according to the patients' diagnosis and the indication of the drug. Patients are free to withdraw from the study at any time without giving a particular reason and without any consequences on further treatment.

Where is the study run from?

The study will take place all over Germany. It is planned that the study will be carried out by approximately 1000 doctors, mainly psychiatrists (practice-based or in outpatient clinics).

When is the study starting and how long is it expected to run for? From 23 April 2012 until 31 January 2013. The recruitment of participants runs until 31 July 2012.

Who is funding the study? Servier Deutschland, GmbH.

Who is the main contact?
Dr Bettina Barthel
bettina.barthel@de.netgrs.com

# Contact information

Type(s)

Scientific

#### Contact name

Dr Martin Kühn

#### Contact details

Servier Deutschland GmbH Elsenheimer Str. 53 Munich Germany 80687 +49 89 57095 308 martin.kuehn@de.netgrs.com

# Additional identifiers

**EudraCT/CTIS** number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers IC4-20098-128-DEU

# Study information

#### Scientific Title

Valdoxan ImproVes depRession with anxiEty symptoms: a non-interventional study

### Acronym

**VIVRE** 

### Study objectives

Effects of Valdoxan therapy on depressive symptoms, emotional and social functioning and anxiety symptoms within depression in adult patients with episodes of major depression under daily routine in an observational, nin-interventional prospective multicentre trial by psychiatrists.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Freiburger International Ethics Committee, 13/02/2012, ref: 012/1098

### Study design

Observational non-interventional prospective multicentre study

### Primary study design

Observational

### Secondary study design

Cohort study

### Study setting(s)

Hospital

### Study type(s)

**Treatment** 

### Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

### Health condition(s) or problem(s) studied

Episodes of major depression

#### **Interventions**

Observational period: 3 months, where the following will be documented:

- 1. Course of depressive symptoms (CGI, adapted version of CGI for patients)
- 2. Evolution of social and emotional functioning (SDS)
- 3. Evolution of anxiety symptoms within depression (COVI)
- 4. Adverse drug reactions
- 5. Liver function tests, if tested

### Intervention Type

Drug

#### Phase

Not Applicable

### Drug/device/biological/vaccine name(s)

Valdoxan

### Primary outcome measure

Obtain information on Valdoxan therapy under daily routine practice by psychiatrists:

- 1. Changes in depressive symptoms under daily conditions via CGI (Clinical Global impression) and anxiety symptoms within depression via COVI-scale
- 2. Effects of the treatment on depressive symptoms via patients' questionnaire (adapted CGI) and emotional and social functioning via SDS (Sheehan disability scale)
- 3. Get information about how Valdoxan SmPC and patients information are followed via standardized documentation of the dosage of Valdoxan, of comedicatins and concomittant diseases
- 4. Analysis of the general tolerability of Valdoxan under routine conditions via standardized adverse drug reactions' documentation and standardized documentation of therapy discontinuation
- 5. Analysis of unknown adverse drug reactions via standardized documentation
- 6. Get further information on known adverse drug reactions under routine practice in comorbide and comedicated patients via standardized adverse drug reactions' documentation and laboratory parameters (liver function testing)

### Secondary outcome measures

No secondary outcome measures

### Overall study start date

23/04/2012

### Completion date

31/01/2013

# Eligibility

### Key inclusion criteria

Adult patients with episodes of major depression according to the SmPC

### Participant type(s)

Patient

### Age group

Adult

#### Sex

Both

### Target number of participants

3000 patients

### Key exclusion criteria

Does not meet inclusion criteria

### Date of first enrolment

# Date of final enrolment 31/07/2012

# Locations

# Countries of recruitment

Germany

# Study participating centre Servier Deutschland GmbH Munich

Germany 80687

# Sponsor information

### Organisation

Servier Deutschland GmbH (Germany)

# Sponsor details

Elsenheimer Str. 53 Munich Germany 80687 +49 89 57095 01 bettina.barthel@de.netgrs.com

### Sponsor type

Industry

### Website

http://www.servier.de/

### **ROR**

https://ror.org/05wk4ae67

# Funder(s)

# Funder type

Industry

### Funder Name

Servier Deutschland GmbH (Germany)

# **Results and Publications**

# Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

### IPD sharing plan summary

Not provided at time of registration

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results		18/01/2019	Yes	No
Results article	results		18/01/2019	Yes	No