# Assessment of the efficacy of Lanreotide Autogel® in the reversal of sleep apnoea, left ventricular hypertrophy and hypertension when used as primary medical therapy in acromegaly

Submission date	<b>Recruitment status</b> No longer recruiting	<ul><li>Prospectively registered</li></ul>	
11/07/2011		☐ Protocol	
Registration date	Overall study status	Statistical analysis plan	
28/09/2011	Completed	[X] Results	
<b>Last Edited</b> 29/01/2018	Condition category Nutritional, Metabolic, Endocrine	Individual participant data	

# Plain English summary of protocol

Background and study aims

Acromegaly is a relatively rare disorder, affecting 4-6 people per million per year. In the majority of cases, a swelling on the pituitary gland releases an excess of growth hormone (GH), which can cause the hands, feet, nose and jaw to enlarge and the skin to become greasy. The condition itself has often been present for some time before the diagnosis is made, and is frequently complicated by high blood pressure, diabetes, enlargement of the heart and sleep disturbance (typically manifesting as snoring). Left untreated, people with acromegaly have a higher risk of developing diseases such as angina, heart attacks and stroke, as well as suffering from problems directly related to the pituitary swelling (e.g. headaches and visual disturbance). Surgery by an experienced neurosurgeon remains the mainstay of treatment, although complete removal of the pituitary swelling is often not possible because of the risk of damage to important surrounding structures, including the nerves to the eyes. Accordingly, further treatment (either in the form of radiotherapy, drugs or, on occasion, both) is often required to control the condition. In recent years it has become standard clinical practice to consider medical treatment prior to surgery with a class of drug commonly referred to as somatostatin receptor ligands (SRLs). These drugs can lower GH levels and improve symptoms and in some instances this is associated with shrinkage of the pituitary swelling, which may improve the surgical outcome. Many centres now routinely offer 'pre-treatment' to newly diagnosed cases of acromegaly. However, it remains unclear as to how effective these drugs are in terms of correcting the sleep disturbance, high blood pressure, metabolic upset, blood vessel changes and heart enlargement of acromegaly. This study aims to examine the effectiveness of a drug called 'Lanreotide Autogel' in reversing blood pressure, cardiac and breathing pattern changes seen in many acromegaly patients before they proceed to surgery.

# Who can participate?

Newly diagnosed patients with acromegaly, aged over 18, who have not undergone any previous medical or surgical treatment or radiotherapy for their acromegaly.

What does the study involve?

Participants are asked to score various symptoms of acromegaly (e.g. headache, sweating, fatique, arthralgia, carpal tunnel syndrome) using a 5-point scale. A full physical examination is performed. Clinical assessments include height, weight, waist and hip circumference, blood pressure measurement, ring size and neck size. At the first visit, blood tests are carried out to confirm the diagnosis of acromegaly. Participants in whom the diagnosis is confirmed are then invited for a second visit during which a more detailed assessment is undertaken with hourly blood sampling to estimate average GH levels. Liver function, fasting glucose and lipid profiles are also checked. Detailed studies of the heart and blood vessels are performed using ultrasound and a magnetic resonance imaging (MRI) scan is performed to determine the pituitary tumour volume. Ultrasound scans of the liver and gallbladder are performed to look for evidence of gall stones and gallbladder polyps. The former can sometimes develop in patients receiving treatment with SRLs and so it is helpful for us to know if these are already present before treatment begins. Participants also undergo overnight sleep studies (polysomnography) to determine the severity of their sleep disturbance. All participants will receive an injection of short-acting SRL octreotide to gauge their likely response to a longer course of SRL treatment. Participants are then treated with the longer acting Lanreotide Autogel, with the dose determined based on their response to octreotide. At each visit, patients are asked to re-score their acromegaly symptoms. At the end of the study, participants are assessed again, as outlined above.

What are the possible benefits and risks of participating? Not provided at time of registration.

Where is the study run from? University of Cambridge (UK).

When is study starting and how long is it expected to run for? March 2004 to March 2014.

Who is funding the study? IPSEN Fund, National Institute for Health Research, and Cambridge Biomedical Research Centre.

Who is the main contact? Dr Mark Gurnell mg299@medschl.cam.ac.uk

# **Contact information**

Type(s)

Scientific

#### Contact name

Dr Mark Gurnell

# **Contact details**

Institute of Metabolic Science and Department of Medicine University of Cambridge Box 289 Addenbrooke's Hospital Hills Road Cambridge United Kingdom CB2 0QQ +44 (0)1223 245 151 mg299@medschl.cam.ac.uk

# Additional identifiers

#### Protocol serial number

v.3 December 2009

# Study information

#### Scientific Title

A comprehensive study of clinical, biochemical, radiological, cardiovascular and sleep parameters in an unselected cohort of patients with newly diagnosed ACROmegaly undergoing Pre-surgical somatostatin Analogue Therapy with Lanreotide Autogel® (ACROPAT)

#### Acronym

**ACROPAT** 

# Study objectives

Medical therapy with Lanreotide Autogel® may improve biochemical parameters like growth hormone (GH) and Insulin-like growth factor 1 (IGF-1) levels which may benefit a comprehensive improvement in radiological, sleep and cardiovascular sequelae of acromegaly in a newly diagnosed acromegaly cohort

# Ethics approval required

Old ethics approval format

# Ethics approval(s)

Cambridgeshire 2 Research Ethics Committee, 09/12/2003, ref: 03/354

# Study design

Prospective open-label cohort study

# Primary study design

Interventional

# Study type(s)

Treatment

# Health condition(s) or problem(s) studied

Acromegaly and medical pre treatment with Lanreotide Autogel®

#### **Interventions**

Phase 1: Baseline assessments

1. Clinical

Subjects will be asked to score various symptoms of acromegaly (e.g. headache, sweating, fatigue, arthralgia, carpal tunnel syndrome) using a 5-point scale (ranging from 0 = not present

to 4 = severe/incapacitating). A full physical examination will be performed. Clinical assessments include height, weight, waist and hip circumference, blood pressure measurement, ring size and neck size estimations.

#### 2. Biochemical

At the first visit, the diagnosis of acromegaly will be confirmed with blood tests, including a glucose challenge test. Those subjects in whom the diagnosis is confirmed will then be invited for a second visit during which a more detailed characterisation of residual pituitary status will be undertaken together with a growth hormone (GH) day profile (1 hourly blood sampling from 09:00 hrs - 17:00 hrs) to permit an estimate of average GH levels pre-treatment. Liver function, fasting glucose and lipid profiles will also be checked.

#### 3. Cardiovascular

Detailed studies of the heart and blood vessels using ultrasound will be performed in conjunction with the Clinical Pharmacology Unit at Addenbrooke's Hospital.

# 4. Radiological

During the second visit, magnetic resonance imaging (MRI) of the pituitary fossa will be performed to permit accurate determination of pituitary tumour volume.

Ultrasound of the liver and gallbladder will also be performed to look for evidence of gall stones and gallbladder polyps. The former can sometimes develop in patients receiving treatment with SRLs and so it is helpful for us to know if these are already present before treatment is commenced.

# 5. Sleep disturbance

All subjects will be invited to the internationally recognised Respiratory Support and Sleep Centre (RSSC), Papworth Hospital for evidence of sleep disturbance. Patients will undergo formal overnight sleep studies (so-called polysomnography) to determine the degree/severity of sleep-disordered breathing.

## Phase 2: Somatostatin analogue administration and interim assessments

At completion of the baseline assessments, all subjects will be started on an injection of subcutaneous (sc) octreotide 100mcg to gauge their likely response to a longer course of SRL therapy - octreotide is a short-acting SRL. Following a repeat GH series and IGF-1 measurement patients will be commenced on the longer acting depot preparation Lanreotide Autogel®, with the dose (60, 90 or 120mg by deep subcutaneous injection every 4 weeks) determined on the basis of mean GH level and observed response to the octreotide test dose.

Repeat GH series and IGF-1 measurements will be performed at 12 weeks to determine whether further dose adjustments are necessary.

At each visit, patients will be asked to re-score their symptoms of acromegaly.

#### Phase 3: Post-treatment assessments

At the completion of the study (6 months post-transition to Lanreotide Autogel®), subjects will be invited to attend for clinical, biochemical, cardiovascular, radiological and sleep-apnoea status re-assessment, as outlined above

# Intervention Type

Drug

#### Phase

Not Applicable

# Drug/device/biological/vaccine name(s)

Lanreotide Autogel®

# Primary outcome(s)

To determine the extent to which pre-surgical medical treatment with the SRL Lanreotide Autogel® ameliorates/reverses the following in newly diagnosed previously untreated acromegaly subjects:

- 1. Sleep disordered breathing
- 2. Cardiac hypertrophy and dysfunction
- 3. Hypertension and arterial dysfunction

# Key secondary outcome(s))

To determine the extent to which pre-surgical medical treatment with the SRL Lanreotide Autogel®:

- 1. Reduces GH and IGF1 levels
- 2. Reduces pituitary tumour volume
- 3. Corrects 'metabolic dysfunction'
- 4. Induces biliary 'sludge'/gallstone formation

## Completion date

01/03/2014

# Eligibility

## Key inclusion criteria

- 1. Treatment naive newly diagnosed subjects with biochemically confirmed acromegaly
- 2. Males and females
- 3. More than 18 years of age

# Participant type(s)

**Patient** 

# Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

All

# Key exclusion criteria

Subjects who had prior medical therapy, surgery or radiotherapy or sight-threatening tumours requiring urgent surgical decompression

#### Date of first enrolment

01/03/2004

#### Date of final enrolment

01/03/2014

# Locations

# Countries of recruitment

**United Kingdom** 

England

Study participating centre University of Cambridge Cambridge United Kingdom CB2 0QQ

# Sponsor information

# Organisation

Addenbrooke's NHS Trust R&D

#### **ROR**

https://ror.org/055vbxf86

# Funder(s)

# Funder type

Industry

# **Funder Name**

IPSEN Fund (UK)

# Alternative Name(s)

# **Funding Body Type**

Private sector organisation

# **Funding Body Subtype**

Other non-profit organizations

# Location

**United Kingdom** 

#### Funder Name

National Institute for Health Research (UK)

# Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

# **Funding Body Type**

Government organisation

# **Funding Body Subtype**

National government

## Location

**United Kingdom** 

#### Funder Name

Cambridge Biomedical Research Centre (UK)

# **Results and Publications**

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Not provided at time of registration

# **Study outputs**

Output type	Details	Date created Date add	ed Peer reviewed	? Patient-facing?
Results article	results	01/03/2013	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/202	25 No	Yes