Effect of local anesthesia on pain sensation during colposcopy

Submission date	Recruitment status No longer recruiting	Prospectively registered		
05/11/2012		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
05/03/2013		[X] Results		
Last Edited	Condition category	[] Individual participant data		
24/01/2019	Surgery			

Plain English summary of protocol

Background and study aims

Traditionally colposcopy [is a procedure where the surface of the cervix is closely examined using a magnifying instrument called a colposcope] with punch biopsies [a biopsy that is performed by using a punch, an instrument for cutting and removing a disk of tissue] is performed without any local anesthesia in Finland. We have observed that using local anesthesia reduces pain and discomfort during this treatment.

Who can participate?

Any woman who have been referred to diagnostic colposcopy in Helsinki University Hospital. Only if they have allergy to anesthetic agent, if they are pregnant, or if there is a suspicion of cancer they are not asked to participate in this study.

What does the study involve?

Half of the participants will be randomly allocated to receive local anesthesia before punch biopsies. Other half will be treated without local anesthesia. Pain will be measured by visual analogue scale (VAS). Participants from both groups will be asked to fill in a questionnaire, Beck's anxiety inventory before the treatment.

What are the possible benefits and risks of participating?

The possible benefit is probably less pain during the punch biopsies.

The only additional risk for participating in the study is possible allergy to anaesthetic agent that the subject is not aware of. All the subjects would be treated the same way despite of participating in the study or not.

Where is the study run from?

The study takes place in Helsinki University Hospital, Department of Obsterics and Gynaecology.

When is the study starting and how long is it expected to run for? The study started in November, 2012 and will last until May 2013.

Who is funding the study?

This study has received funding from Helsinki University Research Foundation.

Who is the main contact? Dr Mari Kiviharju mari.kiviharju@hus.fi

Contact information

Type(s)

Scientific

Contact name

Dr Pekka Nieminen

Contact details

Helsinki University Hospital Bo-BOX 140 Helsinki Finland 00029

Additional identifiers

Clinical Trials Information System (CTIS)

2012-005106-21

Protocol serial number

2012-005106-21

Study information

Scientific Title

Effect of local anesthesia on pain sensation during colposcopy: a randomized trial

Study objectives

The main objective is to measure pain by using visual analogy scale (VAS) during colposcopy and biopsies. We hypothesize that local anesthesia will reduce pain and discomfort.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Helsinki University Institutional Review Board, 2.10.2012, ref: (292/13/03/03/2012)

Study design

Randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Colposcopy: pain during punch biopsies

Interventions

Woman referred to colposcopy will be randomized into two groups

- 1. Colposcopy with local anesthesia
- 2. Colposcopy without local anesthesia

Pain will be measured by VAS scale after biopsies. We will also ask Beck's anxiety inventory for both groups before the treatment.

Intervention Type

Procedure/Surgery

Phase

Not Applicable

Primary outcome(s)

Pain, measured using the Visual Analogue Scale (VAS) score (0 = no pain, 10 = unbearable pain) after punch biopsies.

Key secondary outcome(s))

- 1. Abnormal bleeding after colposcopy
- 2. Beck's anxiety inventory score

Completion date

30/05/2013

Eligibility

Key inclusion criteria

All eligible women who have been referred to diagnostic colposcopy for any cytological abnormality

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

- 1. Allergy to local anesthesia (prilocain + felypressin)
- 2. Pregnancy
- 3. Suspicion of cancer

Date of first enrolment 12/11/2012

Date of final enrolment 30/05/2013

Locations

Countries of recruitment

Finland

Study participating centre Helsinki University Hospital Helsinki Finland 00029

Sponsor information

Organisation

Helsinki University Hospital (Finland)

ROR

https://ror.org/02e8hzf44

Funder(s)

Funder type

University/education

Funder Name

Helsinki University Research Foundation (Finland)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summaryNot provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/04/2017	24/01/2019	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes