

Positioning after idiopathic macular hole surgery

Submission date 02/01/2008	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 30/01/2008	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 18/07/2008	Condition category Eye Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Catherine Creuzot-Garcher

Contact details
Service d'Ophtalmologie
University Hospital Dijon (CHU) General Hospital
Dijon
France
21000

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title
Comparison of face-down and seated position after idiopathic macular hole surgery

Study objectives
To test whether a face-down position is required in every patient operated on idiopathic macular hole, whatever the size of the macular hole.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee for Protection of Human Subjects in Biomedical Research, Bourgogne (Comité Consultatif de protection des personnes dans la recherche biomédicale de Bourgogne). Date of approval: 6 May 2004 (ref: 2004/26)

Study design

Prospective, comparative, randomized controlled trial.

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Idiopathic macular hole

Interventions

All patients are subjected to the same surgical procedure: an extensive three-port pars plana vitrectomy using 20-gauge instrumentation. A peristaltic or venturi pump is used with maximum vacuum set between 200 and 500 mmHg according to the surgeon's preference, to obtain posterior vitreous detachment. The posterior hyaloid is removed. Then the Internal Limiting Membrane (ILM) is systematically removed using microforceps without indocyanine green or any other dye. Vitrectomy is completed, especially at the vitreous base. Finally, total FluidAir Exchange (FAE) was performed and a nonexpanding mixture of air and SF₆ (20%) is used for pneumatic tamponade in Idiopathic Macular Holes (IMHs) less than 500 µm, air and C₂F₆ (17%) in IMHs larger than 500 µm, and air and C₃F₈ (14%) in IMHs larger than 800 µm.

After there procedures, the patients were allocated to two groups. The P0 group is asked to keep a seated position and P1 patients a strict face-down position 8 h a day for 5 days.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Anatomical closure of the hole after one surgical procedure confirmed by OCT. Timepoints of assessment: before, 3 and 6 months after the surgery.

Key secondary outcome(s))

Best corrected visual acuity change between the preoperative and the 6-month visit (expressed as LogMAR)

Completion date

01/02/2006

Eligibility

Key inclusion criteria

Patients with stage 2, 3 and 4 idiopathic macular holes according to a scale developed by Gass and confirmed by Optical Coherence Tomography (OCT)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

1. Axial length longer than 27 mm
2. Previous macular surgical procedure
3. Stage I macular holes
4. Posttraumatic or other secondary macular holes
5. Inability to assume a correct face-down position

Date of first enrolment

01/07/2004

Date of final enrolment

01/02/2006

Locations

Countries of recruitment

France

Study participating centre

Service d'Ophtalmologie

Dijon

France

21000

Sponsor information

Organisation

Burgundy Association for Research in Ophthalmology (ABPRO) (France)

Funder(s)

Funder type

Research organisation

Funder Name

Burgundy Association for Research in Ophthalmology (ABPRO) (France)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results	01/07/2008		Yes	No