

Clinical trial on the efficacy of an intervention package to improve adherence to treatment for systolic blood pressure (BP) reduction in primary care patients with poorly controlled or uncontrolled BP

Submission date 17/06/2010	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 27/07/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 13/01/2015	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Miss Lucia Moreno

Contact details
Gerencia de Atención Primaria
C/ Reina Esclaramunda nº 9
Palma de Mallorca
Spain
07003
-
lmoreno@ibsalut.caib.es

Additional identifiers

Protocol serial number
PS09/01456

Study information

Scientific Title

A randomised controlled trial on the efficacy of a multifactorial intervention to improve adherence to treatment for systolic blood pressure (BP) reduction in primary care patients with poorly controlled or uncontrolled BP

Study objectives

Background:

Lowering blood pressure (BP) with antihypertensive drugs has an important risk reduction effect on cardiovascular events, mainly stroke and total mortality. However lack of adherence to antihypertensive medication reduces its effectiveness and increases the risk for adverse effects to the medication. Improvement of patient adherence to the treatment could be similar in relative risk reduction than the development of a new drug.

Hypotheses:

1. In primary care patients with poorly- or un-controlled BP and low adherence to treatment, a 9-month multifactorial intervention with the aim of improving communication between medical staff and patient and thus facilitating the taking of medication will result in decreased systolic blood pressure (SBP) at 12 months compared with patients in the control group
2. The estimated direct cost at 12 months resulting from the use of health services associated with hypertension in the intervention group will be at least 10% lower than the control group
3. The number needed to treat (NNT) will be 10 - 15

Please note that as of 16/01/2013, the following changes were made to the record:

1. The public title was previously "Clinical trial on the efficacy of an intervention package to improve adherence to treatment for systolic blood pressure (BP) reduction in primary care patients with poorly controlled or uncontrolled BP and low adherence to treatment"
2. The scientific title was previously "A randomised controlled trial on the efficacy of a multifactorial intervention to improve adherence to treatment for systolic blood pressure (BP) reduction in primary care patients with poorly controlled or uncontrolled BP and low adherence to treatment"

Ethics approval required

Old ethics approval format

Ethics approval(s)

The Ethical Committee on Human Research of Balearic Islands approved on the 26th May 2010

Study design

Cluster randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Non-optimally controlled hypertension

Interventions

Patients in the intervention group will attend of 3 sessions consisting of a 30 minute semi-structured motivational interview and a 15-minute health education programme combined with simplifying dosing regimens, family and/or social support, blood pressure self-monitoring and use of medication reminders.

Patients in the control group will receive treatment as usual.

The duration of the intervention will be 6 months. The total duration of follow-up will be 12 months.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Change in systolic BP at 12 months

Key secondary outcome(s)

1. Change in diastolic BP
2. Proportion of participants with adequate BP control at 12 months
3. Total direct cost

Completion date

15/12/2012

Eligibility

Key inclusion criteria

Current inclusion criteria as of 16/01/2013:

1. Patients between 18 and 80 years of age
2. Non controlled BP according to European Societies of Hypertension and Cardiology (ESH/ESC) guidelines (i.e. BP less than 140/90 mmHg or less than 130/80 mmHg in diabetic or renal failure patients)

Previous inclusion criteria until 16/01/2013:

1. Patients between 18 and 80 years of age
2. Non controlled BP according to European Societies of Hypertension and Cardiology (ESH/ESC) guidelines (i.e. BP less than 140/90 mmHg or less than 130/80 mmHg in diabetic or renal failure patients)
3. Adherence below 80% to AHT by Hayness-Sacket Test

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Non-essential hypertensive patients
2. Haemodialysis patients
3. Institutionalised patients

Date of first enrolment

15/10/2010

Date of final enrolment

15/12/2012

Locations**Countries of recruitment**

Spain

Study participating centre

Gerencia de Atención Primaria

Palma de Mallorca

Spain

07003

Sponsor information**Organisation**

Health Service of the Balearic Islands (Servei de Salut de les Illes Balears [IB-salut]) (Spain)

ROR

<https://ror.org/00d9y8h06>

Funder(s)**Funder type**

Government

Funder Name

Health Research Fund (Fondo de Investigaciones Sanitarias [FIS]) (Spain)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	27/09/2010		Yes	No
Other publications	resultst	05/12/2014		Yes	No