

Intermediate care in nursing home

Submission date 19/07/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 26/07/2013	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 12/08/2015	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Intermediate care services are thought to provide adequate care closer to home, preventing hospital admissions and facilitating early discharge. The goal of this study is to assess the effectiveness of intermediate care services. We will compare patients transferred to intermediate care service with patients staying in hospital. The study's findings will help us choose the correct treatment and care for elderly persons with acute illness.

Who can participate?

The study recruited men and women 70 years and older, living at home who were admitted to hospital for acute illness in Bergen, Norway. Patients could participate if they had a medical or orthopedic condition (disorders related to muscles and joints) which did not require surgery or intensive care.

What does the study involve?

Patients were randomly allocated to be either treated in hospital as usual or transferred to intermediate care in a nursing home unit for a stay of maximum three weeks. For the next 12 months, patients' mortality, subsequent admissions to hospital or nursing home and use of home care services were recorded.

What are the possible benefits and risks of participating?

A potential benefit in transfer to intermediate care could be longer stay before returning home. A potential disadvantage could be less access to diagnostic procedures like x-ray. The two treatment options were considered equally safe and appropriate for these patients.

Where is the study run from?

The study was initiated by Bergen municipality, Haraldsplass Deaconess Hospital and Haukeland University Hospital and set up in cooperation with Kavli Research Centre for Ageing and Dementia and Institute of Medicine, University of Bergen, Norway. Centers taking part in the study are Storetveit Nursing Home, Haraldsplass Deaconess Hospital and Haukeland University Hospital, all located in Bergen, Norway.

When is the study starting and how long is it expected to run for?

Recruitment started in August 2007 and ended in June 2008. The study period lasted for a further 12 months after enrolment of the last patient.

Who is funding the study?

Funding has been provided by The Western Norway regional Health Authority and Kavli Research Centre for Ageing and Dementia.

Who is the main contact?

Dr Jo Kåre Herfjord

fmhojkh@fylkesmannen.no

Contact information

Type(s)

Scientific

Contact name

Dr Jo Kåre Herfjord

Contact details

Fylkesmannen i Hordaland

Kaigaten 9

Bergen

Norway

5020

-

fmhojkh@fylkesmannen.no

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

Intermediate care in nursing home after hospital admission: a randomized controlled trial with one year follow-up

Study objectives

The objective of the study is to evaluate effect and safety of early transfer to community-based intermediate care, compared to traditional hospital treatment.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The Regional Committee for Medical Research Ethics in Norway approved the study on 21/05 /2007, ref: S-07096a. Extension of the observation period and new outcome measures were approved on 24/09/2009, ref: 2009/821.

Study design

Parallel-group randomized intervention study. Blinding was not possible. The setting of the study and data collection took place at a nursing home and two hospitals.

Primary study design

Interventional

Secondary study design

Randomised parallel trial

Study setting(s)

Other

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Acute illness or trauma in frail elderly patients with co-morbidity, resulting in hospital admission. Choice of post-acute management and care. Collaboration between hospital and community health services.

Interventions

1. Patients in the intervention group went to intermediate care unit within the next working day after randomization. A multi-disciplinary team of physicians, nurse and physiotherapists examined each patient after a model of comprehensive geriatric assessment. As well as continuing treatment for the acute illness or injury, therapy focused on mobilization, physical exercise, optimal nutrition and critical evaluation of prescribed medication. The aim was maximum independence, enabling patients to resume living at home. Information about home situation and caregivers were obtained and if necessary increased home care services were requested. Length of stay in the unit was limited to maximum 3 weeks
 2. Patients in the control group received usual care in hospital.
- For both treatment arms there was a follow-up period of 12 months.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Our pre-specified primary outcome was the number of days patients would be alive and living at home throughout the first year after randomization. Data were obtained from patient registers at the hospitals and the community health care services for a period of 365 days for each patient after randomization.

Secondary outcome measures

Secondary outcome measures were mortality, days spent in hospital, days in nursing homes and use of home care services.

Overall study start date

15/08/2007

Completion date

02/06/2009

Eligibility

Key inclusion criteria

1. Patients aged 70 years or older
2. Living at home
3. Resident in Bergen municipality
4. Acutely admitted to medical or orthopaedic departments
5. Respiratory and circulatory stable
6. Deemed able to return home within three weeks
7. Intermediate care considered suitable by attending physician

Participant type(s)

Patient

Age group

Senior

Sex

Both

Target number of participants

400

Key exclusion criteria

1. Severe dementia
2. Delirium
3. Need for surgery or intensive care treatment

Date of first enrolment

15/08/2007

Date of final enrolment

01/06/2008

Locations

Countries of recruitment

Norway

Study participating centre

Fylkesmannen i Hordaland, Kaigaten 9

Bergen

Norway

5020

Sponsor information

Organisation

Kavli Research Centre for Ageing and Dementia (Norway)

Sponsor details

Haraldsplass Deaconess Hospital

Ulriksdal 8

Bergen

Norway

5009

-

ida.kristine.sangnes@haraldsplass.no

Sponsor type

Research organisation

Website

<http://kavlisenter.no/index.aspx?pageId=1>

ROR

<https://ror.org/03t3p6f87>

Funder(s)

Funder type

Government

Funder Name

The Western Norway Regional Health Authority (Norway)

Funder Name

Kavli Research Centre for Ageing and Dementia (Norway)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	09/12/2014		Yes	No