

# Double blind, randomised controlled cross over trial comparing the antihypertensive effects of amiloride and spironolactone in black hypertensives with and without the T594M mutation

<b>Submission date</b> 08/03/2002	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 08/03/2002	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 14/03/2017	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Dr PA Swift

### Contact details

Blood Pressure Unit  
Department of Physiological Medicine  
St George's Hospital Medical School  
Cranmer Terrace  
London  
United Kingdom  
SW17 0RE  
+44 20 8725 3176  
pswift@sghms.ac.uk

## Additional identifiers

### Protocol serial number

PG/2001084



# Study information

## Scientific Title

Double blind, randomised controlled cross over trial comparing the antihypertensive effects of amiloride and spironolactone in black hypertensives with and without the T594M mutation

## Study objectives

Not provided at time of registration

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Study design

Randomised controlled trial

## Primary study design

Interventional

## Study type(s)

Not Specified

## Health condition(s) or problem(s) studied

Hypertension

## Interventions

1. Amiloride 5 mg bd for 2 weeks, increased to 10 mg bd for further 6 weeks. Then cross over to spironolactone as below, then to a combination of amiloride 5 mg bd plus spironolactone 25 mg bd
2. Spironolactone 25 mg bd for 2 weeks then increased to 50 mg bd for further 6 weeks. Then cross over to amiloride as above, then to a combination of amiloride 5 mg bd plus spironolactone 25 mg bd

## Intervention Type

Drug

## Phase

Not Specified

## Drug/device/biological/vaccine name(s)

amiloride and spironolactone

## Primary outcome(s)

Not provided at time of registration

## Key secondary outcome(s)

Not provided at time of registration



**Completion date**

31/12/2002

## Eligibility

**Key inclusion criteria**

1. Hypertensives of African origin
2. Subjects with the T594M polymorphism and controls with the wild-type sodium channel

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Sex**

All

**Key exclusion criteria**

Not provided at time of registration

**Date of first enrolment**

01/01/2002

**Date of final enrolment**

31/12/2002

## Locations

**Countries of recruitment**

United Kingdom

England

**Study participating centre**

**Blood Pressure Unit**

London

United Kingdom

SW17 0RE

## Sponsor information



**Organisation**

British Heart Foundation (UK)

**ROR**

<https://ror.org/02wdwnk04>

**Funder(s)****Funder type**

Charity

**Funder Name**

British Heart Foundation (UK)

**Alternative Name(s)**

the\_bhf, The British Heart Foundation, BHF

**Funding Body Type**

Private sector organisation

**Funding Body Subtype**

Trusts, charities, foundations (both public and private)

**Location**

United Kingdom

**Results and Publications****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration