

PREPARE – A clinical trial on a group treatment for distressed caregivers of cancer patients

Submission date 31/03/2016	Recruitment status Stopped	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 07/04/2016	Overall study status Stopped	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 13/02/2020	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Being a caregiver for someone with cancer can be very difficult emotionally. For many the patient is a family member or friend, and so this can cause the caregiver a great deal of psychological distress. There are currently very few studies looking into the effectiveness of psychological treatments for distressed caregivers of cancer patients. This study is looking at tailored group therapy program for carers of cancer givers. The aim of this study is to find out whether this program can help to lower levels of distress in cancer caregivers and help them to better cope with caring for someone with the illness.

Who can participate?

Adult carers of patients who have been diagnosed with cancer in the last six months.

What does the study involve?

Participants are randomly allocated to one of two groups. Those in the first group take part in the tailored group therapy program, over a course of five sessions spread over eight weeks. The sessions are structured group talks and discussions which revolve around pre-specified themes: coping and acceptance, distress, handling cancer-related stressful situations, self-care and a final session for stabilisation. Those in the second group receive a leaflet containing brief information about what individual counselling services are available. One week after the eight week treatment period and again six months later, participants in both groups complete a number of questionnaires in order to find out if their distress levels are lower as well as how well they are coping.

What are the possible benefits and risks of participating?

It is possible that caregivers who take part in the tailored group therapy program will benefit from lower levels of distress and improved coping. There are no notable risks involved with taking part in this study.

Where is the study run from?

Heidelberg University Hospital (Germany)

When is the study starting and how long is it expected to run for?
February 2016 to August 2017

Who is funding the study?
Heidelberg University (Germany)

Who is the main contact?
Dr Markus W. Haun
markus.haun@med.uni-heidelberg.de

Contact information

Type(s)
Scientific

Contact name
Dr Markus W. Haun

ORCID ID
<http://orcid.org/0000-0003-1851-3747>

Contact details
Department of General Internal Medicine and Psychosomatics
Heidelberg University Hospital
Heidelberg
Germany
D-69120
+49 6221 56 38 39 6
markus.haun@med.uni-heidelberg.de

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number
NCT02739243

Secondary identifying numbers
S-655/2015

Study information

Scientific Title
Addressing the hidden patient - A randomised-controlled trial of a tailored group intervention for caregivers of cancer patients

Acronym
PREPARE

Study objectives

A caregiver-specific group intervention is significantly more effective in reducing disease-related distress than treatment as usual that includes brief information and individual referral to the local outpatient counselling centre.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Institutional Review Board of the Medical Faculty Heidelberg, 08/02/2016, ref: S-129/2015

Study design

Randomised controlled pilot trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Community

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet (in German)

Health condition(s) or problem(s) studied

Disease-specific distress in caregivers of cancer patients

Interventions

Gender-stratified 1:1 randomisation with allocation concealment (computed-assisted randomisation in RANDI2 by a blinded data manager). Total treatment duration is eight weeks for both groups. Follow-up for both study arms is at 1-week and 6-months post treatment.

Experimental condition: Participants take part in the tailored group intervention which consists of five 90-minute group sessions over eight weeks. The sessions are in the form of structured group talk/discussions following specific themes:

Session 1 - Coping and acceptance: identification of common themes from participants' feedback, provision of information about support services, mindfulness practices

Session 2 - Distress: introduction of the distress model, identification of resources, breathing practices

Session 3 - Handling of stressful situations: individual identification in tandems, discussion of common pitfalls in the group

Session 4 - Self-care: inspection of individual daily routines, group collects positive experiences

with „windows“ for self-care

Session 5 - Feedback and outlook: stabilisation, retrospective reflection on the group experience and its benefits and, if applicable, potential adverse events

Control condition: Participants are provided with brief information in a leaflet, about the possibilities for individual counselling including referral to the local outpatient cancer counselling centre providing client-centred counselling and financial social work.

Participants in both groups are followed up 1 week and 6 months post-treatment.

Intervention Type

Behavioural

Primary outcome measure

Distress is measured using the Questionnaire on Stress in Caregivers of Cancer Patients Revised version (QSC-R10C) at baseline, 1 week, and 6 months post treatment.

Secondary outcome measures

1. Depression is measured using the Patient Health Questionnaire-9 (PHQ-9) at baseline, 1 week, and 6 months post treatment
2. Anxiety is measured using the Generalized Anxiety Disorder-7 (GAD-7) at baseline, 1 week, and 6 months post treatment
3. General distress is measured using the Distress Thermometer (NCCN DT) at baseline, 1 week, and 6 months post treatment
4. Unmet needs are measured using the Supportive Care Needs Survey – Partners & Caregivers (SCNS-P&C-G) at baseline, 1 week, and 6 months post treatment
5. Self-efficacy in coping with the cancer disease is measured using the adapted General Perceived Self-Efficacy Scale, Positive Bonding Scale of the Group Questionnaire (GQ-D) at baseline, 1 week, and 6 months post treatment
6. Utilisation of primary healthcare is measured via the number of visits with the individual primary care physician at baseline, 1-week, and 6-months post treatment

Overall study start date

01/02/2016

Completion date

30/08/2017

Reason abandoned (if study stopped)

Participant recruitment issue

Eligibility

Key inclusion criteria

1. Time since initial diagnosis (of patient) not longer than six months previously
2. Significant distress against the background of the cancer disease (QSC-R10C exceeding 16 points)
3. Provision of informed consent
4. Aged 18 or over, either gender

Participant type(s)

Carer

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

24

Key exclusion criteria

1. Inability to give informed consent
2. Insufficient German language knowledge
3. Cognitive impairment impeding handling of questionnaires
4. Severe psychiatric disease (acute psychosis or acute suicidality)

Date of first enrolment

15/04/2016

Date of final enrolment

14/05/2016

Locations

Countries of recruitment

Germany

Study participating centre

Heidelberg University Hospital

Department of General Internal Medicine and Psychosomatics

Im Neuenheimer Feld 410

Heidelberg

Germany

D-69120

Sponsor information

Organisation

Department of General Internal Medicine and Psychosomatics

Sponsor details

Heidelberg University Hospital
Im Neuenheimer Feld 410
Heidelberg
Germany
D-69120

Sponsor type

University/education

Website

<https://www.klinikum.uni-heidelberg.de/innere2>

ROR

<https://ror.org/013czdx64>

Funder(s)

Funder type

University/education

Funder Name

Heidelberg University

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Stored in repository