The biological variation of insulin resistance and cardiovascular risk factors in patients with non-alcoholic fatty liver disease compared to type 2 diabetes

| Submission date | Recruitment status | Prospectively registered |
|-------------------|-----------------------------------|---|
| 27/04/2009 | Stopped | ∐ Protocol |
| Registration date | Overall study status | Statistical analysis plan |
| 13/05/2009 | Stopped | ☐ Results |
| Last Edited | Condition category | Individual participant data |
| 18/06/2013 | Nutritional, Metabolic, Endocrine | Record updated in last year |

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Stephen Atkin

Contact details

HS Brocklehurst Building Hull Royal Infirmary Anlaby Road Hull United Kingdom HU3 2RW

Additional identifiers

Protocol serial number R0783

Study information

Scientific Title

The biological variation of insulin resistance and cardiovascular risk factors in patients with nonalcoholic fatty liver disease compared to type 2 diabetes: an observational prospective study

Acronym

BV NASH-T2DM

Study objectives

Patients with Non-alcoholic Fatty Liver Disease (NAFLD) have at least as great intra-individual variance of insulin resistance measured using the homeostasis model assessment (HOMA-IR) as those with the paradigm of insulin resistance; Type 2 Diabetes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Hull and East Riding Research Ethics Committee, approved on 13/03/2009 (ref: LREC 09/H1304/2).

Study design

Observational prospective cross-sectional study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Non-alcoholic fatty liver disease and type 2 diabetes

Interventions

18/06/2013: Please note that this trial was stopped in April 2011.

Twenty type 2 diabetes patients and 20 NAFLD/NASH patients will be recruited for this study (total n = 40).

Over 5 weeks fasting blood samples taken to determine the biological variation of insulin resistance and clotting structure and function and on one occasion endothelial function will be measured using Endo-PAT2000® in the 2 groups - NASH/NAFLD and type 2 diabetes.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

To determine the biological variation of insulin resistance measured by HOMA in patients with NAFLD compared to type 2 diabetes.

Key secondary outcome(s))

- 1. To show statistically significant greater intra-individual variance of insulin resistance measured by HOMA in patients with NASH
- 2. To determine the level of endothelial dysfunction in subjects with NASH compared to NAFLD compared to type 2 diabetes
- 3. To determine the fibrin clot structure and function in subjects with NAFLD/NASH compared to type 2 diabetes

Completion date

30/09/2010

Reason abandoned (if study stopped)

Participant recruitment issue

Eligibility

Key inclusion criteria

For both groups:

Both males and females, over age of 16 years

For NAFLD/non-alcoholic steatohepatitis (NASH) group:

- 1. Patients with confirmed NAFLD/NASH
- 2. Able to give informed consent
- 3. Agreeing to consent to inform GP's regarding participation in the study
- 4. No change in medication for 3 months prior to starting the study

For type 2 diabetes group:

- 1. Diabetes diagnosed according to World Health Organization (WHO) criteria
- 2. No change in medication for 3 months prior to starting study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

For NAFLD/NASH group:

- 1. Patients with concurrent illnesses
- 2. Patients not wishing to allow disclosure to their GP's
- 3. Alcohol intake more than 14 units/week for women, 21 units a week for men
- 4. Diabetic patients
- 5. Pregnancy or breastfeeding women
- 6. Smokers

For type 2 diabetes group:

- 1. Abnormal results from liver function tests (LFTs)
- 2. Patients with concurrent illnesses
- 3. Patients not wishing to allow disclosure to their GP's
- 4. Alcohol intake more than 14 units/week for women, 21 units a week for men
- 5. Pregnancy or breastfeeding women
- 6. Smokers

Date of first enrolment

14/04/2009

Date of final enrolment

30/09/2010

Locations

Countries of recruitment

United Kingdom

England

Study participating centre HS Brocklehurst Building

Hull United Kingdom HU3 2RW

Sponsor information

Organisation

Hull and East Yorkshire Hospitals NHS Trust (UK)

ROR

https://ror.org/01b11x021

Funder(s)

Funder type

University/education

Funder Name

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet
Participant information sheet
11/11/2025 No Yes