

Rapid reduction versus abrupt quitting for smokers who want to stop soon

Submission date 14/10/2008	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
		<input checked="" type="checkbox"/> Protocol
Registration date 22/10/2008	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 16/03/2016	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Paul Aveyard

Contact details
Primary Care Clinical Sciences
University of Birmingham
Edgbaston
Birmingham
United Kingdom
B15 2TT

Additional identifiers

Clinical Trials Information System (CTIS)
2008-006433-28

Protocol serial number
RG_08_082

Study information

Scientific Title

Rapid reduction versus abrupt quitting for smokers who want to stop soon: a randomised controlled non-inferiority trial

Study objectives

The trial will investigate the abstinence success rates of participants reducing cigarette smoking by 50% for two weeks prior to smoking quit date whilst using nicotine replacement therapy (NRT), when compared with abrupt cessation with no cutting down, whilst also utilising NRT for two weeks prior to quitting.

Ethics approval required

Old ethics approval format

Ethics approval(s)

1. Nottingham Research Ethics Committee 2, 20/01/2009, ref: 08/H0408/213
2. Medicines and Healthcare products Regulatory Agency (MHRA), 12/12/2008, ref: 21761/0222/001-0001

Study design

Randomised controlled unblinded non-inferiority trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Smoking cessation in response to nicotine addiction

Interventions

Participants will be seen at an assessment session, similar to that used by the stop smoking services. Here they will be randomised 1:1 to reduction or abrupt cessation arm of the trial.

In the rapid reduction arm participants will decide on a quit date in two weeks time, and in the two weeks leading up to this will reduce smoking consumption by 50%, whilst using nicotine patches (Niquitin, Nicotinell, Nicorette), usually 21 mg, and replacing missing cigarettes with an acute form of nicotine replacement therapy (for example nicotine gum). Participants will be asked to complete a daily diary (explained at the assessment visit). The target number of cigarettes will be completed for each day, and at the end of each day participants will be asked to put aside the next day's cigarettes into a separate pack to encourage adherence to target.

In the abrupt cessation arm participants will also decide on a quit date in two weeks time and be given homework to identify critical cigarettes, which will be the basis for the pre-quit discussion the next week. During the two-week period leading up to the quit date participants will also use patches (usually 21 mg), but will not utilise acute NRT.

Both arms will then go on to have five weekly sessions on quit week and weekly thereafter, following the typical seven-session UK withdrawal-orientated therapy programme. Follow-up

will also be carried out eight weeks and six months after quit day. All NRT used will have the brand name Niquitin, Nicotinell or Nicorette, and participants will be able to have some input in the type(s) of NRT and dosage that they use.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Nicotine (Niquitin, Nicotinell, Nicorette)

Primary outcome(s)

Abstinence at four weeks in all treatment arms, measured according to the Russell standard (allows a two-week grace period from quit day for slips).

Key secondary outcome(s)

Point prevalence at each follow-up and prolonged abstinence at eight weeks and six months (according to the Russell standard). Throughout the trial and at follow-up, the following will be monitored:

1. Urges and withdrawal (using the Mood and Physical Symptoms Scale [MPSS])
2. Self-efficacy
3. Exhaled carbon monoxide (using a CO monitor)
4. Salivary cotinine levels
5. Daily smoking and NRT use (using written diaries)
6. Satisfaction from smoking particular cigarettes (using the Cigarette Evaluation Scale)

Completion date

17/09/2012

Eligibility

Key inclusion criteria

Participants will be recruited by writing to patients on GP practice lists recorded as smokers, writing to people on the stop smoking service's database who have tried and failed to stop, and by offering the treatment to those booking with the stop smoking service. Any participants will be eligible if they are:

1. Prepared to stop either abruptly in two weeks or cut down and stop over two weeks
2. Aged 18 or over, either sex
3. Willing and able to complete data collection

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

Pregnancy will be the only exclusion for insurance purposes

Date of first enrolment

26/06/2009

Date of final enrolment

13/12/2011

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

University of Birmingham

Birmingham

United Kingdom

B15 2TT

Sponsor information**Organisation**

University of Birmingham (UK)

ROR

<https://ror.org/03angcq70>

Funder(s)**Funder type**

Charity

Funder Name

British Heart Foundation (BHF) (UK) (ref: PG/08/047/25082)

Alternative Name(s)

The British Heart Foundation, the_bhf, BHF

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	03/05/2016		Yes	No
Protocol article	protocol	14/08/2009		Yes	No
HRA research summary			28/06/2023	No	No