Can a structured health dialogue with a primary care nurse reduce risk factors for heart disease, offered to a population 55 years old

Submission date	Recruitment status No longer recruiting	Prospectively registered		
25/11/2015		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
10/12/2015	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
02/12/2020	Circulatory System			

Plain English summary of protocol

Background and study aims

There are many risk factors associated with cardiovascular heart disease and stroke. These include having high blood pressure, being overweight, not taking enough exercise, high cholesterol and a unhealthy diet. This study is investigating whether a one hour session with a district nurse on how to reduce these risk factors can improve cardiovascular (heart and circulation) health among people over the age of 55.

Who can participate?

Adults aged 55 and registered at one of the participating primary care (GP) centers.

What does the study involve?

Participants are randomly allocated into one of three groups. Those in group 1 (intervention) fill in a health questionnaire, visit a laboratory for blood tests and then meet a district nurse for a structured health dialogue. During this health dialogue the participant is given information from the district nurse about results from their laboratory tests and questionnaire, shown as a health curve. He or she also, with the support of the district nurse, has the opportunity to plan changes to their lifestyle. Those in group 2 (control group) fill in the same questionnaire and also have blood tests but do not participate in the structured health dialogue with a district nurse. Those in group 3 (comparison group) only fill in the same questionnaire and have the same blood tests a year later, along with the participants in the other two groups.

What are the possible benefits and risks of participating?

Participants who have the health dialogue have the opportunity to get information and also motivation for change and improvement to their lifestyle. All participants get their laboratory test results. Participants visit a doctor if laboratory tests indicates a need for that.

Where is the study run from?

Västmanland County Council (Department of Public Health/Centre for Clinical Research), Sweden.

When is study starting and how long is it expected to run for? May 2010 to December 2012

Who is funding the study?

- 1. County Council of Västmanland, Sweden
- 2. Centre for Clinical Research, Västmanland, Sweden.

Who is the main contact? Mr Mats Hellstrand mats.hellstrand@ltv.se

Contact information

Type(s)

Public

Contact name

Mr Mats Hellstrand

Contact details

Department of Public Health County Council of Västmanland Västerås Sweden 72189 +46 21 17 64 33 mats.hellstrand@ltv.se

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

A health dialogue intervention reduces cardiovascular risk factor levels: a population based randomized controlled trial in primary care setting with one year follow up

Study objectives

Population at the age of 55 who participated in a primary care nurse led health dialogue have at one year follow up significant improved CVD risk factor levels concerning life style, stress and biological markers compared to control.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Regional Ethical Review Board of Uppsala, Sweden, 15/12/2010, ref: 2010/427

Study design

A single-centre interventional study.

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Cardiovascular disease prevention

Interventions

A computerized randomization was made by the Center of Clinical Research, Västerås Hospital, from a list of 1320 inhabitants born 1956-1957 belonging to five primary care centers. Randomization was conducted within each primary care center. The study population of men and women was assigned to three groups, intervention (n=440), control (n=440) and comparison group (n=440).

Intervention group filled in a health questionnaire, visited a laboratory for blood tests and thereafter met a district nurse for a structured health dialogue.

Control group filled in the same questionnaire and visited the same laboratory examination, but received no health dialogue.

The comparison group was invited only at one year follow up, answering the same questionnaire and doing the same laboratory examination.

Intervention Type

Behavioural

Primary outcome(s)

- 1. Weight, height, waist, hip, systolic and diastolic blood pressure, measured by a district nurse.
- 1.1.Weight was measured with light clothing and without shoes
- 1.2. Blood pressure was measured in the right arm in sitting position after five minutes rest.
- 1.3. BMI (kg/m2) was calculated as weight/ (height x height).
- 1.4. Waist circumference was measured in the standing position at the level midway between the 1.5. lower rib margin and the iliac crest.
- 1.6. Hip circumference was measured at the widest point between hip and buttock
- 1.7. Waist–hip ratio (WHR) was calculated as waist circumference divided by hip circumference (cm/cm)
- 1.8. Height was measured without shoes on a fixed wall measure to the nearest centimetre.
- 2. Cholesterol, HbA1c and fasting plasma glucose measured by laboratory staff.
- 3. Self reported variables were physical activity, alcohol consumption last week, diet, smoking habits and stress.

All outcomes are measured one year (11-13 months) after baseline measurement and on all participants in the intervention and control group. Baseline measurement was carried out April-December 2011.

Key secondary outcome(s))

Sub group analysis on high risk groups based on the same measurements as for primary outcomes.

Definition of high risk factor levels: BMI \geq 27 for men and BMI \geq 29 for women, WHR \geq 0.95 for men and \geq 0.83 for women, systolic blood pressure \geq 140, diastolic blood pressure \geq 90, inactive or light physical activity following four degree scale, diet points \geq 6 following eleven degree scale, alcohol \geq 25 cl 40% liquor equivalents/week for men and \geq 17 for women and cholesterol \geq 5.0 mmol/l.

Completion date

31/12/2012

Eligibility

Key inclusion criteria

Inhabitants at the age of 54 or 55 years old at baseline year 2011, registered at one of the five participating primary care centres.

Participant type(s)

Other

Healthy volunteers allowed

No

Age group

Adult

Sex

Αll

Total final enrolment

880

Key exclusion criteria

No exclusion criteria were specified.

Date of first enrolment

01/04/2011

Date of final enrolment

31/12/2012

Locations

Countries of recruitment

Sweden

Study participating centre

Västmanland County Council (Department of Public Health/Centre for Clinical Research), Sweden Västerås Sweden 72189

Sponsor information

Organisation

County Council of Västmanland

ROR

https://ror.org/04vz7gz02

Funder(s)

Funder type

Government

Funder Name

Landstinget Västmanland (County Council of Västmanland)

Funder Name

Center of Clinical Research, County Council of Västmanland

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	22/08/2017	02/12/2020	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes