

(Cost-)effectiveness of a physical, a cognitive-behavioural and an integrated treatment in chronic low back pain disability

Submission date 30/04/2005	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 10/06/2005	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 04/10/2017	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Not provided at time of registration

Study website

<http://www.lobadis.nl/lobadis/study3/default.htm>

Contact information

Type(s)

Scientific

Contact name

Dr Rob Smeets

Contact details

Rehabilitation Centre Blixembosch
P.O. Box 1355
Eindhoven
Netherlands
5602 BJ
+31 (0)40 2642878
rsmeets@iae.nl

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

014-32-007

Study information

Scientific Title

(Cost-)effectiveness of a physical, a cognitive-behavioural and an integrated treatment in chronic low back pain disability: a randomised controlled trial

Acronym

Lobadis 3

Study objectives

The treatment of non-specific chronic low back pain (CLBP) is often based on different models regarding the maintenance of pain and disability:

1. Loss of muscle strength and endurance including aerobic capacity (physical deconditioning model)
2. Avoidance of activity due to learning, cognitive processes and environmental influences (cognitive behavioral model)
3. The combination of both models (biopsychosocial model)

Four differing treatments were developed:

1. Active Physical Treatment (APT)
2. Cognitive-Behavioural Treatment (CBT)
3. Combination Treatment consisting of APT plus CBT (CT)
4. Waiting List control group (WL)

It is hypothesised that:

1. All three active treatments are more effective in reducing low back pain disability than a waiting list control treatment (short term analysis)
2. CT is more effective and cost-effective than APT and CBT

Additional research questions are:

1. Does deconditioning exist in CLBP-patients?
2. What factors mediate the decrease in outcome (especially the effect of pain catastrophising and internal control of pain will be investigated)
3. Can prognostic factors for each of the treatments be identified?
4. Do physiological and psychological factors that are often described as very important, influence the level of performance in CLBP-patients?
5. Psychometric research on the Roland Disability Questionnaire and Quebec Back Pain Disability Scale (stability and responsiveness)

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Treatment

Participant information sheet**Health condition(s) or problem(s) studied**

Non-specific chronic low back pain

Interventions

Active Physical Treatment (APT):

Cardiovascular and leg/back extending muscle strengthening (three sessions/week for ten weeks)

Cognitive-Behavioural Treatment (CBT):

Combination of graded activity of patient relevant activities and problem solving training (19 and ten sessions in ten weeks respectively)

Combination Treatment (CT):

Combination of APT and CBT in the same frequency

Waiting List (WL):

After ten weeks of no treatment regular rehabilitation treatment is offered

Post-treatment WL will be compared to APT, CBT and CT. Also, CT will be compared to APT and CT. Six and 12 months post-treatment CT will be compared to APT and CBT.

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Roland Disability Questionnaire (RDQ)

Secondary outcome measures

1. Three patient specific main complaints by using the patient specific approach method (Tugwell et. al.)
2. Current pain by using the Visual Analogue Scale (VAS) for pain at this moment
3. Pain Rating Index (total score) of the McGill Pain Questionnaire
4. Beck Depression Inventory

5. Patients global assessment of overall result measured by a transitional seven-point ordinal scale (1 = vastly worsened, 7 = completely recovered)
6. Treatment satisfaction by using VAS for the overall treatment provided to the patient
7. Six performance tests:
 - a. five minutes walking (meters)
 - b. fifty foot walking (seconds)
 - c. five times sit to stand, performed twice and average time needed to perform a series of five times is calculated (seconds)
 - d. loaded forward reaching by holding a stick with a weight of 2.25 or 4.5 kg in front of the body at shoulder height and extend as far as possible (centimeters)
 - e. one minute stair climbing (number of stairs)
- f. Progressive Isoinertial Lifting Evaluation (PILE)-test weight lifting from floor to waist

Overall study start date

01/04/2002

Completion date

31/12/2004

Eligibility

Key inclusion criteria

1. Patients for the first time referred to three outdoor rehabilitation centers in the Netherlands
2. Age between 18 and 65 years
3. Non specific CLBP with or without irradiation to leg for more than three months resulting in disability (Roland Morris Disability score more than three)
4. Ability to walk at least 100 m without interruption

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

223

Key exclusion criteria

1. Vertebral fracture
2. Spinal inflammatory disease
3. Spinal infections or malignancy
4. Current nerve root pathology
5. Spondylolysis or -listhesis
6. Lumbar spondylodesis

7. Medical co-morbidity making intensive exercising impossible (e.g. cardiovascular or metabolic disease)
8. Ongoing diagnostic procedures or treatment for their CLBP at the time of referral
9. Not proficient in Dutch
10. Pregnancy
11. Substance abuse that could intervene with the rehabilitation treatment
12. Patients have to stop other therapies except pain medication for their back complaints
13. The Symptom Checklist (SCL-90) and the Dutch Personality Questionnaire (NPV) are used to check for psychopathology that would hamper individual or group processes

Date of first enrolment

01/04/2002

Date of final enrolment

31/12/2004

Locations

Countries of recruitment

Netherlands

Study participating centre

Rehabilitation Centre Blixembosch

Eindhoven

Netherlands

5602 BJ

Sponsor information

Organisation

Netherlands Organisation for Health Research and Development (ZonMw)

Sponsor details

Laan van Nieuw Oost Indië 334

P.O. Box 93245

Den Haag

Netherlands

2509 AE

+31 (0)70 349 5111

info@zonmw.nl

Sponsor type

Research organisation

ROR

Funder(s)

Funder type

Research organisation

Funder Name

,950 Euro funded by Netherlands Organisation for Health Research and Development (ZonMw)
(014-32-007)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	20/01/2006		Yes	No