Study to see if GSK2110183 with chemotherapy can treat ovarian cancer

Submission date	Recruitment status No longer recruiting	Prospectively registered		
16/01/2014		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
16/01/2014		[X] Results		
Last Edited	Condition category	[] Individual participant data		
14/04/2021	Cancer			

Plain English summary of protocol

http://www.cancerresearchuk.org/cancer-help/trials/a-study-looking-cell-changes-ovariancancer-aktres

Contact information

Type(s)

Scientific

Contact name

Ms Maria Martinez

Contact details

Hammersmith Hospital Du Cane Road London United Kingdom W12 0HS

Maria.martinez@imperial.nhs.uk

Additional identifiers

Clinical Trials Information System (CTIS)

2012-002483-27

ClinicalTrials.gov (NCT)

NCT01653912

Protocol serial number

13614, PKB116611

Study information

Scientific Title

AKTRES study: A Biologic Study of the early effects and determinants of AKT inhibition using GSK2110183 alongside chemotherapy in patients with platinum RESistant adenocarcinoma of the ovary

Acronym

AKTRES

Study objectives

Activation of the AKT signalling pathway in ovarian cancer cells causes chemotherapy resistance. An AKT inhibitor, given alongside chemotherapy, could potentially reverse this resistance and enable chemotherapy to be effective once again. This study is designed for women whose ovarian cancer has relapsed within 6 months of receiving carboplatin or cisplatin-containing chemotherapy and who are receiving an AKT inhibitor given alongside 3 weekly carboplatin and paclitaxel in the PKB116611 trial. Participants in the AKTRES study undergo a maximum of two biopsies from their tumour and some additional blood tests. The aims of the study are to better understand the how chemotherapy-resistant cancer cells are affected by AKT inhibitors.

Ethics approval required

Old ethics approval format

Ethics approval(s)

London Hampstead, 07/11/2012, ref. 12/LO/1174

Study design

Non-randomised; Interventional; Design type: Process of Care, Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: National Cancer Research Network; Subtopic: Gynaecological Cancer; Disease: Ovary /Fallopian tube

Interventions

Blood sampling, [3 x 10 ml whole blood collected in greentopped (heparinised) tube] will be collected at baseline (within 14 days prior to first dose), on day 1 (immediately preceding chemotherapy), day 8 and day 15 of cycle 1 only. Blood sampling will subsequently be performed at day 1 only of subsequent chemotherapy treatment cycles and thereafter 3 weekly for those on maintenance GSK2110183 alone. A final 3 x 10 ml blood sample will be taken at end of study visit.

Coagulation profile, will be collected immediately prior to the first biopsy during the screening period and postcycle 1 biopsy on day 22. This is in addition to the screening coagulation blood test required for PKB116611 study.

Collection of ascites: If patients are undergoing ascitic or pleural drainage for symptom relief

prior to, or during, this study a sample of ascites (or pleural effusion) can be taken and stored. The preferred times for collection are during screening, day 22 and at study completion, but additional/alternative collections can be obtained at the investigators discretion. Tumour Biopsy: A tumour biopsy performed under image guidance within 14 days prior to first dose. Three tumour cores will be taken, two will be fresh frozen and one will be formalin fixed. Following one complete treatment cycle (day 22), =3 days prior to starting cycle 2 day 1, a second tumour biopsy will be performed under image guidance. Again, three tumour cores will be taken, two will be fresh frozen and one will be formalin fixed. Patients can decline the second biopsy if they wish to.

Intervention Type

Drug

Phase

Phase I/II

Drug/device/biological/vaccine name(s)

GSK2110183 (afurasertib)

Primary outcome(s)

Overall Response, CA125 response and Progression free survival measured by ELISA detected decrease in phosphorylated PRAS40 (as ratio of total PRAS40) and increase in pAKT S473

Key secondary outcome(s))

Not provided at time of registration

Completion date

04/12/2014

Eligibility

Key inclusion criteria

- 1. Female at least 18 years of age at the time of signing the informed consent form
- 2. Capable of giving written informed consent
- 3. Having measurable tumour tissue and Platinum-resistant ovarian cancer as defined as radiological evidence of disease progression within 6 months of completion of platinum-containing chemotherapy
- 4. Tumour tissue measuring >2cm long axis amenable to direct biopsy or biopsy via imagequidance

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Total final enrolment

59

Key exclusion criteria

Concurrent medication with warfarin or low molecular weight heparin (heparin use is acceptable if it has been discontinued for 2 days preceding biopsy)

Date of first enrolment

10/12/2012

Date of final enrolment

04/12/2014

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Hammersmith Hospital

London United Kingdom W12 0HS

Sponsor information

Organisation

Imperial College of Science, Technology and Medicine (UK)

ROR

https://ror.org/041kmwe10

Funder(s)

Funder type

Charity

Funder Name

Ovarian Cancer Action (UK)

Alternative Name(s)

Funding Body Type

Private sector organisation

Funding Body Subtype

Other non-profit organizations

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/03/2019	14/04/2021	Yes	No
Basic results				No	No
Basic results				No	No
Participant information she	Participant information sheet	11/11/2025	11/11/2025	No	Yes