Optimal keratoplasty for the correction of presbyopia and hypermetropia

Submission date 04/03/2017	Recruitment status No longer recruiting	 Prospectively registered Protocol
Registration date 14/03/2017	Overall study status Completed	
Last Edited 15/02/2022	Condition category Eye Diseases	[] Individual participant data

Plain English summary of protocol

Background and study aims

Presbyopia is a condition associated with aging of the eye that leads to s progressively worsening ability to focus clearly on close objects. In a younger person, the lens is soft and flexible so it can easily reshape the lens to focus on close and distant objects. With age, the lens becomes harder so it is unable to reshape as easily. Hyperopia, also known as farsightedness, is another forma of presbyopia which makes it difficult for people to focus on things that are close. Optimal keratoplasty is a new form of laser eye treatment to treat presbyopia. It works by remodeling the shape of the cornea (transparent layer forming the front of the eye) to help sharpen distance vision and near vision. The aim of this study is to investigate the safety and effectiveness of optimal keratoplasty in correcting hyperopia (long-sightedness) and presbyopia.

Who can participate?

Adults aged 40 years and over who have hyperopia and presbyopia

What does the study involve?

All participants receive optimal keratoplasty, which involves having a special laser applied to their eyes while they are lying down. The procedure takes around 10-15 minutes altogether. Before the procedure and then one hour, one day, one week and one, three and six months after the procedure, participants have an eye test in order to find out if there has been any improvement to their vision.

What are the possible benefits and risks of participating?

Participants may benefit from an improvement to their eyesight. There is a small risk that some participants may experience minor vision problems for a short time after the procedure, such as glare (light entering the eye and interfering with vision) and halos (bright circles that surround a light source).

Where is the study run from? Istituto Europeo di Microchirurgia Oculare (Italy)

When is the study starting and how long is it expected to run for? June 2014 to June 2016 Who is funding the study? Istituto Europeo di Microchirurgia Oculare (Italy)

Who is the main contact? Professor Paolo Lanzetta

Contact information

Type(s) Public

Contact name Prof Paolo Lanzetta

ORCID ID http://orcid.org/0000-0003-3746-141X

Contact details Istituto Europeo di Microchirurgia Oculare via Fiducio 8 Udine Italy 33100

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers I-2014A27

Study information

Scientific Title

In patients with presbyopia and hypermetropia, do optimal keratoplasty improve near and distance visual acuity?

Study objectives Optimal keratoplasty is safe and effective for the correction of hyperopia and presbyopia.

Ethics approval required Old ethics approval format

Ethics approval(s)

Istituto Europeo di Microchirurgia Oculare (IEMO) Institutional Review Board, 22/03/2014, ref: I-2014A27

Study design

Single-centre single-arm non-randomised study

Primary study design Interventional

Secondary study design Non randomised study

Study setting(s) Hospital

Study type(s) Quality of life

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Presbyopia and hypermetropia

Interventions

All patients undergo laser irradiation of the corneal tissue using the NTK Optimal Keratoplasty (Opti-K) System.

While the patient is lying in a supine position, an output beam is directed onto the cornea in a symmetrical square ring pattern (6 and 7.2 mm diameter), so that the corneal epithelium is protected from thermal damage with a sapphire application window/suction ring. Thereafter, laser is applied at 1.93 µm wavelength. The laser is typically operated with a total delivered power of 0.80-1.28 W for a period of 150 ms. The duration of the entire procedure is approximately 10-15 minutes.

Patients are evaluated at baseline, 1 hour, 1 day, 1 week, 1 month, 3, and 6 months after treatment. The duration of a visit is about an hour. Baseline and follow-up examinations include measurement of uncorrected distance visual acuity (UDVA), uncorrected near visual acuity (UNVA), CDVA, manifest refraction, corrected near visual acuity (CNVA), presbyopic add, corneal optical coherence tomography (OCT), corneal pachymetry, slit lamp examination of the anterior segment, and dilated fundus examination.

Intervention Type

Procedure/Surgery

Primary outcome measure

Mean uncorrected near visual acuity is measured by ETDRS charts and standardized procedures at baseline and 6 months.

Secondary outcome measures

Uncorrected distance visual acuity in hypemetropic/presbyopic eyes is measured by ETDRS charts and standardized procedures at baseline and 6 months.

Overall study start date

01/06/2014

Completion date

01/06/2016

Eligibility

Key inclusion criteria

1. Aged 40 years and over

2. Low to moderate hypermetropia (manifest refraction: sphere between +1 to +2.5 D, absolute cylinder ≤1 D) or presbyopia (with presbyopic adds between +1D to +2.75 D)

3. Documented stable refraction

4. Corrected distance visual acuity (CDVA) of 33 Early Treatment Diabetic Retinopathy Study (ETDRS) letters or better

Participant type(s)

Patient

Age group Adult

Sex

Both

Target number of participants

35

Key exclusion criteria

- 1. Nystagmus
- 2. Corneal diameter ≤9 mm
- 3. Central corneal thickness ≤500 µm
- 4. Dry eye disease
- 5. Severe blepharitis
- 6. Residual, recurrent or active corneal disease or abnormality

Date of first enrolment

01/07/2014

Date of final enrolment 01/05/2016

Locations

Countries of recruitment Italy Study participating centre Istituto Europeo di Microchirurgia Oculare via Fiducio 8 Udine Italy 33100

Sponsor information

Organisation Istituto Europeo di Microchirurgia Oculare

Sponsor details via Fiducio 8 Udine Italy 33100

Sponsor type Other

ROR https://ror.org/02t9kcf24

Funder(s)

Funder type Other

Funder Name Istituto Europeo di Microchirurgia Oculare

Results and Publications

Publication and dissemination plan Planned publication in a Medline indexed peer reviewed journal.

Intention to publish date 31/08/2017

Individual participant data (IPD) sharing plan

Participant level data are stored as medical charts at the center. Charts are anonymised with a numeric code. Patients signed an informed consent allowing data storage.

IPD sharing plan summary

Stored in repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		18/04/2017	15/02/2022	Yes	No