

The optimal warming strategy to reduce low body temperature during surgery

Submission date 08/03/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 11/03/2024	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 16/06/2025	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Patients having surgery with anesthesia often lose body heat and become cold during the surgery. This can be associated with discomfort and, if significant, with medical side effects such as abnormal heart rhythm. A variety of methods and devices exist to warm patients up and/or prevent loss of body heat. While these are used routinely and are approved by regulatory agencies around the world, and are safe and (mostly) external to the body, the optimal combination and timing of their use is not entirely clear. The purpose of this study is to find out which of four combinations of warming strategies is optimal for these patients.

Who can participate?

Men and women at least 18 years of age, who are having elective abdominal, gynecologic, breast surgery, plastic/reconstructive or urological surgery under general anesthesia, where the operation is projected to last longer than 1.5 hours but less than 4 hours.

What does the study involve?

The patients will be randomly assigned to four groups that represent combinations of warming strategies. Group 1 will have an electrical warming blanket (conductive warming, or CW) used before their surgery and during their surgery. Group 2 will have a forced air blanket (FAW) used before and during their surgery. Group 3 will have no active warming before surgery and CW during surgery, but warm cotton blankets before the surgery will be allowed if patient wants to use them. And 4) will have no active warming before surgery and FAW during the surgery. Again in group 4 warm cotton blankets before the surgery will be allowed. All of these warming methods (CW and FAW) are well-established in practice and are achieved with common equipment. No method in and of itself is novel or lacking regulatory approval. All these methods are non-invasive.

What are the possible benefits and risks of participating?

The possible benefits are that if one of these groups turns out to be a superior strategy for keeping the patients warm, then those who have been assigned to that group will be more comfortable and/or will lose less body heat during their surgery. The risks are that the patient may have been assigned in a group where they will lose more body heat during the surgery.

Where is the study run from?
Cooper University HealthCare, New Jersey, USA.

When is the study starting and how long is it expected to run for?
February 2018 to August 2022

Who is funding the study?
Augustine Medical Inc., Eden Prairie, MN, USA

Who is the main contact?
Dr. Ronak Desai, D.O., Desai-ronak@cooperhealth.edu

Contact information

Type(s)
Public, Scientific, Principal investigator

Contact name
Dr Ronak Desai

ORCID ID
<https://orcid.org/0000-0001-7831-6492>

Contact details
One Cooper Plaza, Department of Anesthesiology
Camden
United States of America
08103
+1 856-342-2425
Desai-ronak@cooperhealth.edu

Additional identifiers

Clinical Trials Information System (CTIS)
Nil known

ClinicalTrials.gov (NCT)
Nil known

Protocol serial number
Nil known

Study information

Scientific Title
The optimal warming strategy to reduce perioperative hypothermia: a prospective randomized non-blinded clinical trial

Study objectives

1. Active warming with a conductive warming system (CW) will be non-inferior in preventing hypothermia when compared to active warming with a forced air warming system (FAW).
2. Active preoperative warming with CW or FAW will lead to a reduction in intraoperative hypothermia when compared with only intraoperative warming with CW or FAW.
3. Active preoperative warming combined with intraoperative warming using CW is a superior strategy compared to only intra-operative warming with the CW, intraoperative warming with FAW only, or pre-operative and intra-operative warming with FAW.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 30/09/2019, Cooper University HealthCare IRB (One Cooper Plaza, Camden, NJ, 08103, United States of America; +1 (856) 757-7832; IRBOnlineHelp@cooperhealth.edu), ref: 18-148

Study design

Prospective randomized non-blinded investigator-initiated clinical trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Patients having surgery with general anesthesia

Interventions

Participants are randomized to four groups, each representing a different preoperative warming approach:

1. Conductive prewarming (CW) and conductive intraoperative warming i.e. CW+CW;
2. Forced air (FAW) prewarming with forced air intraoperative warming i.e. FAW+FAW;
3. No active prewarming (NAPW) with conductive intraoperative warming i.e. NAPW+CW; and
4. No active prewarming (NAPW) with forced air intraoperative warming i.e. NAPW+FAW.

Of note, all of these warming methods are well-established in practice and are achieved with commonly available methods/equipment. No method in and of itself is novel or lacking regulatory approval. All these methods are non-invasive.

Duration of intervention: from signing of informed consent form and/or arrival to pre-operative surgical area (if form was signed ahead of time), to end of surgery.

Duration of follow-up: 30 days post-surgery.

Randomisation: we used a computer generated randomization list (Excel) to allocate subjects to one of four warming strategies in a 1:1:1:1 ratio. We used block randomization for scheduled surgical strata (1.5 h – 2.5 h and 2.5 – 4 h) and type of surgery (abdominal, gynecologic, breast surgery, plastic/reconstructive, urologic surgery) to ensure equal distribution of likely confounding variables.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Intraoperative hypothermia magnitude will be the primary outcome. Hypothermia will be defined as a core temperature $<36^{\circ}\text{C}$ and quantified as the intraoperative area under the curve (AUC) $< 36^{\circ}\text{C}$ (units $^{\circ}\text{C}\cdot\text{hr}$) detected with an esophageal temperature probe.

Key secondary outcome(s)

1. Lowest temperature measured intraoperatively: degrees C, measured via esophageal temperature probe. Recorded in the electronic anesthesia record.
2. The percentage of the time of the case spent hypothermic: unit is minutes. Measured as the time in minutes under 36 degrees C divided by the duration of the operation (surgery) in minutes x 100. Times obtained from the electronic anesthesia record.
3. The absolute incidence of hypothermia: this is a binary score. I.e. if body temperature dropped below 36 degrees C at any point during the surgery, it was a yes. If not, then no.
4. The thermal comfort score: ordinal 5-point patient-reported Likert scale. Obtained by study coordinator at the end of the patient's stay in the pre-operative area, i.e. just before transfer to the operating room. See attached PDF for the Likert scale questions.

Completion date

08/08/2022

Eligibility

Key inclusion criteria

1. Subjects undergoing elective abdominal, gynecologic, breast surgery, plastic/reconstructive or urological surgery under general anesthesia
2. Surgery projected to last longer than 1.5 hours, but no longer than 4 hours

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

100 years

Sex

All

Total final enrolment

184

Key exclusion criteria

1. Cardiac surgery
2. Vascular surgery
3. Pregnancy
4. Age < 18 years
5. Imprisonment
6. Inability to provide written informed consent
7. Inability to speak and/or read English

Date of first enrolment

04/12/2019

Date of final enrolment

08/08/2022

Locations**Countries of recruitment**

United States of America

Study participating centre**Cooper University healthcare**

One Cooper Plaza

Camden, NJ

United States of America

08103

Sponsor information**Organisation**

Cooper University HealthCare, Department of Anesthesiology

Funder(s)**Funder type**

Not defined

Funder Name

Augustine Medical Inc., Eden Prairie, MN, USA

Results and Publications

Individual participant data (IPD) sharing plan

The full dataset will be uploaded into <https://datadryad.org/stash> before submission to a journal for peer review.

IPD sharing plan summary

Stored in publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		12/06/2025	16/06/2025	Yes	No
Participant information sheet			11/03/2024	No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file		22/05/2019	11/03/2024	No	No