

# Pilot study designed to assess functionality of a low-cost negative pressure wound therapy system

<b>Submission date</b>	<b>Recruitment status</b>	<input type="checkbox"/> Prospectively registered
21/05/2018	No longer recruiting	<input type="checkbox"/> Protocol
<b>Registration date</b>	<b>Overall study status</b>	<input type="checkbox"/> Statistical analysis plan
30/05/2018	Completed	<input type="checkbox"/> Results
<b>Last Edited</b>	<b>Condition category</b>	<input type="checkbox"/> Individual participant data
30/05/2018	Other	<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Healyx Labs is developing a Negative Pressure Wound Therapy device (NPWT) designed for low-resource medical settings. The Healyx Labs NPWT device is designed to provide a controlled amount of suction (negative pressure) to a wound site through an airtight dressing applied to a patient's wound. Suction removes fluids from the wound and provides wound contraction and tissue changes that can promote wound healing. The aim of this study is to evaluate the feasibility of using the NPWT device in low-resource medical settings.

### Who can participate?

Patients over 18 years of age who have a wound that can be treated by the device. These wounds include chronic, acute, traumatic, subacute and dehisced wounds, partial-thickness burns, ulcers (such as diabetic, pressure, or venous insufficiency), flaps, grafts and surgical incisions

### What does the study involve?

After wound cleansing and preparation, treatment with the NPWT device begins. The dressing is cut to size and applied. The suction device is attached to the dressing and turned on. The vacuum is left on for up to six weeks, or until the investigator determines that the wound is healed or ready for a further intervention. During this time, caregivers are instructed to check on the treatment daily. Wound dressing changes occur at least twice a week. After completion of the intervention, the patient is contacted four to six weeks later to assess satisfaction with treatment and quality of life.

### What are the possible benefits and risks of participating?

Healyx Labs is not claiming to directly benefit participants. Healyx does expect the NPWT to provide comparable benefits to those of existing negative-pressure wound therapy vacuums, including faster healing. Healyx Labs is not introducing new risks to those listed with standard NPWT devices. Additionally, standard NPWT risk for adverse events is lower than that associated

with makeshift wall and surgical suction devices. All dressings used in the study are certified as sterile and the Healyx device itself provides a pressure of 125 mmHg, considered safe for use in negative pressure wound therapy.

Where is the study run from?  
Kirtipur Hospital (Nepal)

When is the study starting and how long is it expected to run for?  
November 2017 to May 2018

Who is funding the study?  
Healyx Labs (USA)

Who is the main contact?  
Cam Hutton

## Contact information

### Type(s)

Public

### Contact name

Mr Cam Hutton

### Contact details

2500 Grant Rd G01  
Mountain View  
United States of America  
94040

## Additional identifiers

### Protocol serial number

CLN100

## Study information

### Scientific Title

Pilot study designed to assess functionality of a low-cost negative pressure wound therapy system in patients with chronic, acute, traumatic, subacute and dehisced wounds, partial-thickness burns, ulcers (such as diabetic, pressure, or venous insufficiency), flaps, grafts and surgical incisions

### Acronym

Healyx NPWTD Pilot Study

### Study objectives

Primary Objectives: To determine NPWT device ease of use in a low-resource setting  
Secondary Objectives: To assess safety of device design; to assess healing progression and quality of life for patients

## **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Public Health Concern Trust, Nepal (phect-NEPAL) Institutional Review Committee (IRC) , 14/02 /2018, ref: 1824

### **Study design**

Prospective single-arm interventional pilot study in a single site

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Chronic, acute, traumatic, subacute and dehisced wounds, partial-thickness burns, ulcers (such as diabetic, pressure, or venous insufficiency), flaps, grafts and surgical incisions

### **Interventions**

The Healyx Labs NPWT suction unit provides suction at the gold standard level for Negative Pressure Wound Therapy (-125mmHg +/- 10%) from a built-in diaphragm pump powered by a DC motor. The pressure setting at -125 mmHg is the standard pressure used in predicate devices and has application in treatment for majority of wound types. The suction unit has a casing that holds a disposable suction container that collects fluids leaving the patient's sealed wound bed.

After wound cleansing and preparation, treatment with the NPWT device begins. The dressing is cut to size and applied. The suction device is attached to the dressing and turned on. The vacuum is left on for up to six weeks, or until the investigator determines that the wound is healed or ready for a further intervention. During this time, caregivers are instructed to check on the treatment daily. Wound dressing changes occur at least twice a week. After completion of the intervention, the patient is contacted four to six weeks later to assess satisfaction with treatment and quality of life indicators.

### **Intervention Type**

Device

### **Primary outcome(s)**

Ease of use measured by user reported rankings of device procedures at each dressing change:

1. Users (caregivers) are confident in setting up the device and creating successful suction.

Device user completes a device assessment at each dressing change (twice per week) and at device removal measuring:

1.1. Overall satisfaction with a Likert scale

1.2. Usability compared to current wound therapies

2. Users are able to successfully remove the device during dressing changes. Device user completes a device assessment at each dressing change and at device removal measuring:

2.1. Overall satisfaction with a Likert scale

2.2. Usability compared to current wound therapies

## **Key secondary outcome(s)**

1. Safety of device design evaluated by the incidence and intensity of device-related adverse events (AEs). Patient is interviewed and observed visually for device-related adverse events at treatment initiation and each dressing change
2. Healing progression at six weeks or conclusion of therapy, if earlier
3. Pain measured using a validated pain scale VAS recorded at 1x dressing change per week
4. Quality of life measured using questionnaire modified using the EuroQol Derived Single Index (EQ-DSI) recorded at 1x dressing change per week

## **Completion date**

26/05/2018

## **Eligibility**

### **Key inclusion criteria**

1. Patient is 18 years of age or older
2. Patient presents with an open wound
3. Patient is able to understand study requirements, is willing to undergo subject procedures, provide any required subject feedback, and provide informed consent to participate in the study

### **Participant type(s)**

Patient

### **Healthy volunteers allowed**

No

### **Age group**

Adult

### **Lower age limit**

18 years

### **Sex**

All

### **Key exclusion criteria**

1. Patient who is pregnant or nursing
2. Patient with a non-survivable wound, life expectancy of 7 days or less
3. Patient's wound is of malignant or vasculitis origin, or shows presence of necrotic tissue
4. Patient's wound has active bleeding, exposed blood vessels or organs
5. Patient with wound-related comorbidities, including diabetic ketoacidosis, Hyperglycaemic Hyperosmolar Nonketotic Coma, vascular disease, or arthritis
6. Patient currently using medications that cause immune suppression
7. Patient with wound considered too difficult to seal based on location
8. Patient unable to understand study or provide informed consent to participate in the study

### **Date of first enrolment**

26/02/2018

### **Date of final enrolment**

18/04/2018

## Locations

### Countries of recruitment

Nepal

### Study participating centre

Kirtipur Hospital

Thapathali-11

Kathmandu

Nepal

44618

## Sponsor information

### Organisation

Healyx Labs

## Funder(s)

### Funder type

Industry

### Funder Name

Healyx Labs

## Results and Publications

### Individual participant data (IPD) sharing plan

The company is not making the participant level data available as the device does not yet have regulatory approval and the data are confidential and proprietary. The data are held at the clinical trial site in Nepal.

### IPD sharing plan summary

Not expected to be made available

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>	Participant information sheet	11/11/2025	11/11/2025	No	Yes

