

# Dialectical Behavior Therapy for men with borderline personality disorder and antisocial behavior

<b>Submission date</b> 02/06/2014	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 11/09/2014	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
<b>Last Edited</b> 25/06/2020	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Dialectical behavior therapy (DBT) is a psychological treatment that is known to work well for women with borderline personality disorder (BPD). It has been developed from cognitive behavior therapy (CBT) but modified to meet the needs of people with BPD. CBT focuses on helping people change unhelpful ways of thinking and behaving. DBT does this too, but it also encourages people to accept who they are and includes elements of mindfulness, a mind-body approach developed to help people change how they think about experiences and how to reduce feelings of stress and anxiety. However, to date, there has been a lack of DBT-treatment studies looking at how it may help men with BPD being treated in outpatient clinics. Here, we will look at how well DBT performs when treating men with BPD with a history of anti-social behavior. We want to know to what extent 12 months of DBT might reduce suicidal, impulsive and antisocial behaviors, how much it improves psychological well-being and whether it is considered useful by the participants involved. We also want to investigate variables that might influence the effect of the treatment.

### Who can participate?

Men with a diagnosis of borderline personality disorder and antisocial behaviors within Stockholm Psychiatry

### What does the study involve?

After the diagnosis of BPD has been made, participants are asked to self-record their antisocial and impulsive behaviors for 4 weeks before the treatment starts. They also attend an interview where they recall their antisocial behavior. They then start a 12 month course of DBT therapy, where they learn how to manage their emotional reactions and destructive behaviors. Participants are asked to self-record their antisocial/impulsive behaviors and attend further interviews 6 months after the treatment starts and at the end of treatment.

What are the possible benefits and risks of participating?

All participants receive intensive DBT treatment, which is known to be helpful for women with BPD. Based on earlier studies, we expect the treatment to help most participants reduce their self-harming, impulsive and anti-social behaviors.

Where is the study run from?

The treatment will be conducted at two psychiatric outpatient clinics in Stockholm.

When is the study starting and how long is it expected to run?

December 2013 to December 2016

Who is funding the study?

Swedish Prison and Probation Service (Sweden)

Who is the main contact?

Pia Enebrink, Ph.D.

[pia.enebrink@ki.se](mailto:pia.enebrink@ki.se)

## Contact information

### Type(s)

Scientific

### Contact name

Dr Pia Enebrink

### Contact details

Division of Psychology

Karolinska Institutet

Nobels väg 9-11

Solna

Sweden

171 65

## Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

## Study information

### Scientific Title

Effectiveness trial of Dialectical Behavior Therapy for men with borderline personality disorder and antisocial behavior

**Study objectives**

1. What effect does Dialectical Behavior Therapy (DBT) have on self-injurious behavior, depression, anxiety, as well as antisocial, criminal behaviors for men with borderline personality disorder?
2. How do participants experience the treatment, and are there any components of the treatment they find particularly helpful?
3. What is the potential cost-effectiveness of the intervention?
4. Which variables predict (moderators), or explain (mediators) the outcome of DBT?

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

The regional ethical committee in Stockholm, 06/03/2012, ref. 2012/91-31/4

**Study design**

Intervention study with within-subjects design

**Primary study design**

Interventional

**Secondary study design**

Other

**Study setting(s)**

Hospital

**Study type(s)**

Treatment

**Participant information sheet****Health condition(s) or problem(s) studied**

Borderline personality disorder

**Interventions**

Dialectical Behavior Therapy. DBT is based on cognitive behavior therapy (CBT) with the addition of acceptance and mindfulness practices. In treatment, participants learn skills to manage their intensive emotional reactions and destructive behaviors. The treatment is characterized by a combination of individual sessions and skills treatment in group and telephone support for the client, and supervision for the therapists. The treatment lasts for 12 months.

**Intervention Type**

Other

**Phase**

Not Applicable

**Primary outcome measure**

1. Structured interviews collecting data on earlier antisocial behaviors using the Timeline Follow Back Method
2. Daily registration of antisocial behaviors: participants will register antisocial and impulsive behaviors during a 4-week baseline period before treatment starts, as well as throughout the treatment period
3. Swedish National Centre for Crime Prevention, reported offenses

### **Secondary outcome measures**

Self-ratings:

1. Beck Depression Inventory II (BDI)
2. Beck Anxiety Inventory (BAI)
3. Alcohol use disorders identification test (AUDIT)
4. Drug use disorders identification test (DUDIT)
5. Karolinska affective and borderline symptom scale- self rating (KABOSS-S)
6. ASEBA-ASR (Adult Behavior Checklist)
7. DBT-skills rating scale
8. Health and occupation survey

Questionnaire for relatives:

1. ASEBA-ABCL (Adult Behavior Checklist)

Therapist ratings:

1. Attendance of therapy sessions
2. Homework completion

### **Overall study start date**

09/12/2013

### **Completion date**

31/12/2016

## **Eligibility**

### **Key inclusion criteria**

Men with borderline personality disorder and antisocial behavior

### **Participant type(s)**

Patient

### **Age group**

Adult

### **Sex**

Male

### **Target number of participants**

32

### **Total final enrolment**

**Key exclusion criteria**

1. Psychotic disorder
2. Severe substance abuse
3. Autistic disorders

**Date of first enrolment**

09/12/2013

**Date of final enrolment**

31/12/2016

**Locations****Countries of recruitment**

Sweden

**Study participating centre****Division of Psychology**

Solna

Sweden

171 65

**Sponsor information****Organisation**

Swedish Prison and Probation Service (Sweden)

**Sponsor details**

Kriminalvården

601 80 Norrköping

Norrköping

Sweden

601 80

**Sponsor type**

Government

**ROR**

<https://ror.org/04030gz13>

**Funder(s)**

**Funder type**  
Government

**Funder Name**  
Swedish Prison and Probation Service (Sweden)

## Results and Publications

**Publication and dissemination plan**  
Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**  
Not provided at time of registration

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/02/2020	25/06/2020	Yes	No