Post-op oesophageal Doppler trial: effect on clinical outcome of extension of oesophageal Doppler goal-directed fluid optimisation into the post-operative recovery period

Submission date	Recruitment status No longer recruiting	Prospectively registered		
12/07/2012		[X] Protocol		
Registration date	Overall study status Completed Condition category Digestive System	Statistical analysis plan		
14/09/2012		Results		
Last Edited		Individual participant data		
18/05/2018		Record updated in last year		

Plain English summary of protocol

Background and study aims

Cardiac output is the volume of blood pumped by the heart per minute. It can be measured by placing a probe into the oesophagus (food pipe) via the mouth or nose, a procedure called oesophageal Doppler monitoring. Guiding treatment with cardiac output monitoring has been shown to benefit patients undergoing major colorectal surgery, improving outcomes and reducing the length of hospital stay. Traditionally this monitoring has only been used during surgery. With the introduction of flexible probes, this monitoring can now be used in awake patients after surgery. The group most likely to benefit are high-risk patients. We aim to find out about the effect of continuing the use of this monitoring to guide treatment after surgery in high-risk colorectal surgery patients.

Who can participate?

High-risk patients aged 18 and over undergoing colorectal surgery.

What does the study involve?

Participants are randomly allocated to one of two groups. Participants in one group undergo oesophageal Doppler cardiac output monitoring for 16 hours (overnight) after their operation. Participants in the other group receive standard care with the probe in place but not used for monitoring. We then compare the outcomes of the two groups during their hospital stay and at follow-up.

What are the possible benefits and risks of participating?

Oesophageal Doppler monitoring has already been tested in cardiac and trauma patients. Patients can request early removal of the probe before the end of the 16-hour monitoring period if they wish. Patients who may be at risk of nasal or oesophageal injury from the probe are excluded from the study.

Where is the study run from? St Thomas' Hospital (UK).

When is the study starting and how long is it expected to run for? May 2011 to October 2012

Who is funding the study? Guy's and St. Thomas' Charity (UK).

Who is the main contact? Mr Jonathan van Dellen jvandellen@doctors.org.uk

Contact information

Type(s)

Scientific

Contact name

Mr Andrew Williams

Contact details

Department of Colorectal Surgery St Thomas' Hospital Westminster Bridge Road London United Kingdom SE1 7EH

Type(s)

Scientific

Contact name

Dr Jonathan van Dellen

ORCID ID

https://orcid.org/0000-0002-1343-6140

Contact details

N/A London United Kingdom N/A N/A jvandellen@doctors.org.uk

Additional identifiers

Protocol serial number

RJ1 11/N160

Study information

Scientific Title

Effect on clinical outcome of extension of oesophageal Doppler goal-directed fluid optimisation into the post-operative recovery period: a randomised control trial.

Study objectives

Extension of OD-guided goal-directed fluid optimisation into the immediate 16-hour post-operative period, in high risk colorectal ERP patients, achieves an oxygen delivery index ≥600ml /min/ m2 for the majority of the duration of this period, and subsequently improves patient outcomes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

NRES Committee London - Westminster, 06/04/2011, ref: 11/H0802/9

Study design

Single-centre prospective single-blinded (assessor blinded to outcomes) randomised control study within an academic tertiary referral hospital setting

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Improving perioperative care for elective colorectal surgery within an Enhanced Recovery Programme

Interventions

Intervention group will have 16 hours post-operative oesophageal Doppler cardiac output monitoring with nurse-led stroke volume optimisation in an overnight intensive recovery (OIR) unit.

Control group will have an equivalent OIR admission without cardiac output monitoring (nasal oesophageal Doppler probe still in-situ for 16h measurements).

Follow-up of outcomes during hospital admission and standard telephone and outpatient clinic follow-up.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Post-operative length of stay (in days)

Key secondary outcome(s))

- 1. Oxygen delivery index at end of 16h post-operative recovery period
- 2. Mean oxygen delivery index over 16h post-operative recovery period
- 3. Morbidity
- 4. Functional recovery (mobility and return of gastrointestinal function)
- 5. In-hospital and 30-day mortality

Completion date

01/10/2012

Eligibility

Key inclusion criteria

- 1. Consecutive elective or semi-elective patients due to undergo colorectal surgery as part of Enhanced Recovery Protocol involving resection or anastamosis of bowel
- 2. High-risk participants will be defined as having (1) pre-operative ASA grade III+ OR anaerobic threshold <11 ml/min/kg on cardiopulmonary exercise testing OR undergoing major complex surgery (as per OPCS 4.5 classification Office of Population, Censuses and Surveys Classification of Surgical Operations and Procedures)
- 3. Ages 18 and over (no upper limit)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. Emergency surgery
- 2. Lack of capacity to consent
- 3. Pregnancy
- 4. Contraindications to oesophageal Doppler probe:
- 4.1. Nasal injuries or polyps
- 4.2. Severe oesophageal/laryngeal/pharyngeal disease
- 4.3. Recent oesophageal/laryngeal/pharyngeal surgery
- 4.4. Thoracic aortic aneurysm
- 4.5. Severe bleeding diathesis
- 4.6. Long-term systemic steroid therapy
- 4.7. Portal hypertension

All excluded patients from patient groups to be reported in results

Date of first enrolment

24/05/2011

Date of final enrolment

20/09/2012

Locations

Countries of recruitment

United Kingdom

England

Study participating centre St Thomas' Hospital

London United Kingdom SE1 7EH

Sponsor information

Organisation

Guy's & St Thomas' NHS Foundation Trust (UK)

ROR

https://ror.org/00j161312

Funder(s)

Funder type

Charity

Funder Name

Guy's and St. Thomas' Charity (UK) ref: G100706

Alternative Name(s)

Guy's and St Thomas' Charity, Guy's and St Thomas' Foundation, GSTTFoundation

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date. Current plans are for submission and storage with King's College London Data Management system.

IPD sharing plan summary

Stored in repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet				No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	version v1	14/10/2011		No	No