

# Benefit of machine learning to diagnose deep vein thrombosis compared to the gold standard ultrasound

<b>Submission date</b> 11/05/2023	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 15/05/2023	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 27/05/2025	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Deep vein thrombosis (DVT) is a term that describes blood clots (thrombi) that can form in the deep veins. The deep leg veins are commonly affected (such as the proximal veins: the femoral vein or the popliteal vein) or the deep veins of the pelvis. The standard approach to making a diagnosis involves an algorithm combining pre-test probability, a blood test called the D-dimer test, and the patient undergoing an ultrasound of the leg veins. Ultrasound is currently completed by a trained expert (e.g. sonographer or radiologist). However, handheld ultrasound probes have recently become available and they have enabled 'app-based' ultrasonography to be performed. ThinkSono has developed software (AutoDVT software) allowing non-specialists to perform DVT ultrasound, hoping it has the same accuracy for diagnosing DVT as the standard ultrasound. If this study has a positive outcome, it would mean that DVT could be diagnosed at the point of care by non-experts such as nurses, junior doctors, GPs and other healthcare staff. By diagnosing DVT early in the clinical pathway (for example, at GP practices), the technology could reduce emergency department admissions and free up specialists to focus on other clinical tasks. These improvements could also potentially reduce the financial burden of the DVT diagnostic service on healthcare systems.

### Who can participate?

Patients aged 18 years and over, coming for a check to see if they have a DVT and have symptoms suggesting that they need an ultrasound scan

### What does the study involve?

Participants undergo two compression ultrasound scans. One is carried out by a non-radiology staff member (e.g. a nurse) using AI software to guide them and another ultrasound scan will be carried out as already scheduled by a sonographer or radiologist.

### What are the possible benefits and risks of participating?

This study will not benefit participants directly in the short term but it may benefit patients having an ultrasound for a DVT in the future. The results from this study will improve knowledge of how software may be able to help diagnose blood clots accurately and quickly. Ultrasound is a

very safe method of confirming a DVT or not and is used already as standard care in hospitals. There are no risks of taking part. The scan does involve some pressing on the leg but if it is painful or participants want to stop they can let the researchers know.

Where is the study run from?

Helios Klinikum Emil von Behring, Klinik für Angiologie (Germany)

When is the study starting and how long is it expected to run for?

January 2022 to September 2022

Who is funding the study?

1. Imperial College London (UK)

2. Helios Klinikum Emil von Behring, Klinik für Angiologie (Germany)

Who is the main contact?

Sven Mischkewitz (Sponsor contact), [hello@thinksono.com](mailto:hello@thinksono.com)

## Contact information

### Type(s)

Scientific

### Contact name

Mrs Kerstin Nothnagel

### ORCID ID

<https://orcid.org/0009-0004-5767-8346>

### Contact details

University of Bristol

Bristol Medical School

Room 1.10 (Desk 3)

Canynges Hall,

Clifton

Bristol

United Kingdom

BS8 2PN

+4915141406431

[kerstin.nothnagel@bristol.ac.uk](mailto:kerstin.nothnagel@bristol.ac.uk)

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil Known

### ClinicalTrials.gov (NCT)

Nil known

### Protocol serial number

V1.0

# Study information

## Scientific Title

Benefit of machine learning to diagnose deep vein thrombosis compared to the gold standard ultrasound

## Study objectives

Measure the sensitivity and specificity of clinical decision-making based on images recorded by a non-specialist using AutoDVT AI-based software versus the ground truth produced by the local imaging specialist.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved on 15/03/2022, Ethics Committee of the Berlin Medical Association (Ethik-Kommission der Ärztekammer Berlin), (Working Committee Research II [Arbeitsausschuss Forschung II] Ärztekammer Berlin KdöR, Friedrichstr 16, 10969 Berlin, Germany; +49 30 408 06 26 01; ek@aekb.de), ref: Eth-07/21

## Study design

Non-randomized prospective study

## Primary study design

Observational

## Study type(s)

Diagnostic

## Health condition(s) or problem(s) studied

Proximal deep vein thrombosis

## Interventions

Patient recruitment

Patients will consent when a DVT ultrasound exam is requested in the angiology department.

An AI-assisted scan with the AutoDVT software is performed by a non-specialist (nurse). A follow-up gold-standard scan is performed by a local specialist (compression ultrasound), i.e. angiologist. That same-day follow-up scan represents the standard of care.

The images collected by the non-specialist are presented to a remote, qualified clinician who evaluated image quality according to the quality scale of the American College of Emergency Physicians (ACEP) and consequently, if the image quality is sufficient, assesses whether the veins of the patient are compressible, incompressible or indeterminate.

## Study arms

Single-arm study. Every patient will receive an AI-guided scan which was followed up by a reference scan which represents the standard of care.

#### Intervention provider

The nursing staff carried out the AI-guided ultrasound scan. They had no prior ultrasound experience at all. The remote, qualified clinician assessing the images collected by the AI-guided scan are qualified to diagnose DVT, i.e., radiologists.

#### Modes of delivery

The AI-guided scan is performed face-to-face. The images that are presented to the remote, qualified clinician will be evaluated retrospectively via an internet platform. This remote, qualified clinician will not see the patient. The AI-guided scan was carried out in the rooms of the angiology department.

#### Intervention Type

Device

#### Phase

Not Applicable

#### Drug/device/biological/vaccine name(s)

AutoDVT

#### Primary outcome(s)

Sensitivity and specificity measured using the AI-guided ultrasound and a local imaging specialist performing the gold-standard ultrasound exam at one timepoint. The gold-standard exam is performed on the same day.

#### Key secondary outcome(s)

Image quality of the AI-guided ultrasound assessed by remote qualified clinicians, according to the American College of Emergency Physicians (ACEP) scoring scale from 1 to 5, at one timepoint

#### Completion date

01/09/2022

## Eligibility

#### Key inclusion criteria

1. Aged 18 years old and over
2. Subject can consent and consent has been signed
3. Subject has symptoms of DVT and ultrasound is indicated

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

**Sex**

All

**Total final enrolment**

91

**Key exclusion criteria**

1. Below the age of 18 years old
2. No Wells score was calculated prior to the ultrasound
3. Distal DVT
4. Did not consent

**Date of first enrolment**

18/04/2022

**Date of final enrolment**

09/08/2022

**Locations****Countries of recruitment**

Germany

**Study participating centre**

Helios Emil von Behring Klinikum Zehlendorf

Walterhöferstrasse 11

14165 Berlin Germany

Berlin

Germany

14165

**Sponsor information****Organisation**

ThinkSono GmbH

**Funder(s)****Funder type**

Hospital/treatment centre

**Funder Name**

# Results and Publications

## Individual participant data (IPD) sharing plan

The datasets generated and analysed during the current study will be available upon request from Sven Mischkewitz (hello@thinksono.com) if a request is made by relevant authorities and a statement about the use of the data has been made. Anonymised ultrasound data may be shared. No patient data will be shared due to patient confidentiality.

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		02/01/2025	27/05/2025	Yes	No
<a href="#">Participant information sheet</a>	Participant information sheet	11/11/2025	11/11/2025	No	Yes