A trial of tongue-tie division and its immediate effect on breastfeeding

Submission date	Recruitment status No longer recruiting	Prospectively registered		
22/06/2011		Protocol		
Registration date 15/07/2011	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 24/09/2015	Condition category	[] Individual participant data		
24/09/2015	Other			

Plain English summary of protocol

Background and study aims

Ankyloglossia, more commonly known as "tongue-tie", is a common congenital (present from birth) condition which affects the way the tongue moves in the mouth. About 4% of children are born with this problem, in which the cord-like skin (frenulum) that joins the bottom of the tongue to the floor of the mouth is too short, limiting the tongue's range of movement. In some cases, babies may have difficulties breastfeeding, as they are unable to open their mouths wide enough or to suck properly. This can lead to painful nipples for the mother, and can mean that the baby does not gain weight as it should. If feeding is affected, then a tongue-tie division procedure is recommended. This simple operation involves cutting the frenulum to allow the tongue to move. Tongue-tie division is often carried out when the child is a few months old, and afterwards the baby is able to feed normally straight away. It has been suggested that the fact that mothers report an immediate improvement in breastfeeding could be down to the placebo effect (that it works only because it is expected to). The aim of this study is to investigate the possibility that the reported improvements in breastfeeding are due to the placebo effect.

Who can participate?

Babies under four months of age with tongue-tie, who have difficulties with breastfeeding.

What does the study involve?

Participants are randomly allocated into two groups. Those in the first group receive the tonguetie division procedure, and those in the second group did not receive the procedure. All infants are returned to their mothers with a square of gauze under their tongues so that the mothers do not know whether their child has had the procedure or not. Before and after the procedure, mothers are asked to feed their babies and to give their opinions on how well their child is feeding, and their level of pain.

What are the possible benefits and risks of participating? Not provided at time of registration.

Where is the study run from?
Wessex Regional Centre for Paediatric Surgery (UK)

When is the study starting and how long is it expected to run for? October 2003 to July 2004

Who is funding the study? Southampton University Hospitals NHS Trust (UK)

Who is the main contact? Mr Mervyn Griffiths

Contact information

Type(s)

Scientific

Contact name

Mr Mervyn Griffiths

Contact details

Wessex Regional Centre for Paediatric Surgery Southampton General Hopital Tremona Road Southampton United Kingdom SO16 6YD

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers 120/01/t

Study information

Scientific Title

A double-blind, randomised, controlled trial of tongue-tie division and its immediate effect on breastfeeding

Study objectives

That division of tongue-tie causes an immediate change in breastfeeding which the mother can detect and is not due to placebo effect

Ethics approval required

Old ethics approval format

Ethics approval(s)

Southampton and South West Hampshire Local Research Ethics Committees, 6 November 2003 ref: 120/01/t

Study design

Single centre double-blind randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Quality of life

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Tongue-tie

Interventions

- 1. Both participants & the observer were blinded to the intervention
- 2. Tongue-tie division and non-division
- 3. Unblinded follow up at 1 day & 3 months

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

- 1. Subjective and objective change in breastfeeding
- 2. Maternal pain score

Secondary outcome measures

- 1. Subjective changes in breastfeeding
- 2. Breastfeeding rates
- 3. Complications and acceptability of procedure to parents

Overall study start date

30/10/2003

Completion date

31/07/2004

Eligibility

Key inclusion criteria

- 1. Less than 4 months of age
- 2. Tongue-tie present
- 3. Symptomatic of breastfeeding problem

Participant type(s)

Patient

Age group

Child

Upper age limit

4 Months

Sex

Both

Target number of participants

60

Key exclusion criteria

Does not meet inclusion criteria

Date of first enrolment

30/10/2003

Date of final enrolment

31/07/2004

Locations

Countries of recruitment

England

United Kingdom

Study participating centre Wessex Regional Centre for Paediatric SurgerySouthampton

Southampton United Kingdom SO16 6YD

Sponsor information

Organisation

Southampton University Hospitals NHS Trust (UK)

Sponsor details

Southampton General Hospital Tremona Road Southampton England United Kingdom SO16 6YD

Sponsor type

Hospital/treatment centre

Website

http://www.suht.nhs.uk/home.aspx

ROR

https://ror.org/0485axj58

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Southampton University Hospitals NHS Trust (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/06/2012		Yes	No