

# INTERVAL study: To determine whether the interval between blood donations in England can be safely and acceptably decreased

<b>Submission date</b> 29/11/2011	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 25/01/2012	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 29/08/2019	<b>Condition category</b> Other	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

In order to ensure that there is enough supply of blood for hospitals and patients who need it, many people are asked to donate their blood. By giving blood, donors can help save lives. Blood donation is a very simple process, where a needle is inserted into a vein in the arm and around a pint of blood is collected. The entire process takes around five to ten minutes. However, there are often shortages in blood supply, especially for rare blood types. These shortages can be life threatening. One way to increase the blood supply is to ask previous blood donors to give more frequently, however donors must maintain enough time between donations in order to ensure their body replaces their blood iron levels. No one knows the ideal amount of time that should be waited between donations and the intervals between donations differ across countries. In England, the NHS Blood and Transplant team (NHSBT) invites men to donate every 12 weeks and women to donate every 16 weeks but in Europe some countries ask donors to donate every eight weeks. This study aims to recruit 50,000 blood donors to compare different intervals between blood donations. The goal is to find the optimum interval for which it is safe for different donors to give blood and to look at whether intervals should be tailored by age, gender, genetic profile, and other characteristics. The study's findings should help to improve the well-being of future blood donors in England and enhance the country's blood supplies.

### Who can participate?

Adults over the age of 17 who attend an NHS blood donation clinic in England.

### What does the study involve?

Participants are asked to join this study while they are at a blood donation clinic. Participants must pass the screening haemoglobin (blood iron) finger-prick test in order to give blood. Participants are randomly allocated to one of two groups. Those in the first group are asked to give blood at the usual donation intervals (based on their gender). Those in the second group are asked to donate blood more frequently. Men are asked to donate every eight, ten or 12 weeks and women every 12, 14 or 16 weeks. The study lasts two years in total. Participants are

asked to give an additional blood samples at the beginning and at the end of the study to test for the blood iron levels and for DNA. Participants also complete online questionnaires and assessments every six-months during the study period.

What are the possible benefits and risks of participating?

There will be no immediate direct benefit to those taking part. But there should be benefits to future blood donors and to the country's future blood supply because the results of the study are likely to influence how the NHSBT collects blood donations. The main risk of giving blood more frequently is iron deficiency and the related anaemia because of a low haemoglobin level. Therefore, NHSBT will continue to follow its routine safety procedures to monitor haemoglobin level before donation. Participants will receive the usual finger-prick screening test for haemoglobin levels and will need to be within the safe range to be eligible to donate.

Where is the study run from?

The INTERVAL study is being run by the Universities of Cambridge and Oxford and takes place in NHSBT blood donation clinics across England.

When is the study starting and how long is it expected to run for?

November 2011 to November 2021

Who is funding the study?

NHS Blood and Transplant team (NHSBT) (UK)

Who is the main contact?

1. Prof. John Danesh

john.danesh@phpc.cam.ac.uk

2. Prof. David Roberts

david.roberts@ndcls.ox.ac.uk

**Study website**

<http://www.intervalstudy.org.uk>

## Contact information

**Type(s)**

Scientific

**Contact name**

Prof John Danesh

**Contact details**

Department of Public Health and Primary Care

University of Cambridge

Wort's Causeway

Cambridge

United Kingdom

CB1 8RN

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

NCT01610635

**Secondary identifying numbers**

11-01-GEN

## **Study information**

### **Scientific Title**

A randomised trial to determine whether the interval between blood donations in England can be safely and acceptably decreased

### **Acronym**

INTERVAL

### **Study objectives**

The number of donations made by English blood donors will be greater with reduced versus standard inter-donation intervals. The null hypothesis is that there will be no difference in donations between treatment groups; this may arise if reduced inter-donation intervals result in a greater number of donation deferrals (due to low haemoglobin) and/or an unacceptable burden to donors.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

NRES Committee East of England - Cambridge East, 16/02/2012, ref: 11/EE/0538

### **Study design**

Two-year open randomised parallel-group multi-site trial

### **Primary study design**

Interventional

### **Secondary study design**

Randomised controlled trial

### **Study setting(s)**

Other

### **Study type(s)**

Other

### **Participant information sheet**

The patient information sheet is available on the trial website <http://www.intervalstudy.org.uk>

## **Health condition(s) or problem(s) studied**

Optimum intervals for blood donation

## **Interventions**

The study will involve 25,000 male and 25,000 female whole blood donors recruited at donation visits to the 25 permanent NHSBT clinics across England. Participation will be over a two-year period.

Male participants will be assigned to standard 12-week vs. 10-week vs. 8-week inter-donation intervals and female participants to 16-week vs. 14-week vs. 12-week intervals.

Added 14/03/2017:

INTERVAL participants were recruited between June 2012 and June 2014, and were studied over a two-year period. Following ethical approval of a substantial amendment to the protocol, some donors (who had completed their 2-year participation) were invited to continue on their randomised donation frequency for an additional period of 6 months – 2 years, and were randomised to receive either routine or more intensive reminders to attend for blood donation. This study extension will enable additional questions to be answered on the acceptability and sustainability of shorter donation intervals, as well as on the effectiveness of different appointment-reminder approaches.

## **Intervention Type**

Other

## **Phase**

Not Applicable

## **Primary outcome measure**

Total blood collected after two years, expressed in units (470ml) per person per year

## **Secondary outcome measures**

Assessed two years after recruitment and will include:

1. Donor quality of life using the SF-36 health survey (this is the key secondary outcome)
2. Number of donation 'deferrals' (i.e. temporary rejections) of donors due to low haemoglobin and other factors
3. Markers of iron status (serum ferritin and reticulocyte haemoglobin)
4. Cognitive ability (reasoning, attention and memory)
5. Levels of physical activity
6. Cost effectiveness
7. Donor attitudes, beliefs and values

## **Overall study start date**

01/11/2011

## **Completion date**

01/11/2021

## **Eligibility**

### **Key inclusion criteria**

1. Age >17 years and fulfilling all normal criteria for blood donation
2. Willing to be assigned to any of the study intervention groups
3. Registered at one of the permanent donation clinics at the time of enrolment

**Participant type(s)**

Healthy volunteer

**Age group**

Adult

**Sex**

Both

**Target number of participants**

50,000

**Key exclusion criteria**

Current exclusion criteria as of 22/05/2012:

1. As the aim of the study is to be almost paper-less, it will involve remote web-based data collection. Hence, participants who do not have internet access and/or are not willing to provide an email address for study correspondence will be excluded

Previous exclusion criteria:

1. Donors who are providing extra samples to NHSBT at their initial visit (e.g. for the British bone marrow registry) as there will be insufficient blood left in the sample pouch for the collection of study samples
2. As the aim of the study is to be almost paper-less, it will involve remote web-based data collection. Hence, participants who do not have internet access and/or are not willing to provide an email address for study correspondence will be excluded
3. Donors who have been identified for special panels by NHSBT

**Date of first enrolment**

11/06/2012

**Date of final enrolment**

15/06/2014

**Locations****Countries of recruitment**

England

United Kingdom

**Study participating centre**

**University of Cambridge**

Department of Public Health and Primary Care

Strangeways Research Laboratory

Worts Causeway

Cambridge  
United Kingdom  
CB1 8RN

## Sponsor information

### Organisation

NHS Blood and Transplant (NHSBT) (UK)

### Sponsor details

c/o Professor Marion Scott  
National R&D Manager  
500 North Bristol Park  
Northway  
Filton  
Bristol  
United Kingdom  
BS34 7QH

### Sponsor type

Government

### ROR

<https://ror.org/0227qpa16>

## Funder(s)

### Funder type

Government

### Funder Name

NHS Blood and Transplant (ref: 11-01-GEN)

### Alternative Name(s)

National Health Service Blood and Transplant, UK National Health Service Blood and Transplant, NHSBT

### Funding Body Type

Government organisation

### Funding Body Subtype

Local government

### Location

## Results and Publications

### Publication and dissemination plan

Planned publication in a high-impact peer reviewed journal

### Intention to publish date

01/11/2022

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request and approval from the INTERVAL Data Access Committee (helpdesk@intervalstudy.org.uk)

### IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol article</a>	protocol	17/09/2014		Yes	No
<a href="#">Results article</a>	results	20/09/2016		Yes	No
<a href="#">Results article</a>	results	25/11/2017		Yes	No
<a href="#">Results article</a>	extension study results	01/10/2019	07/08/2019	Yes	No