

A phase II randomised feasibility study of chemoresection or surgical management in low risk non muscle invasive bladder cancer

Submission date 14/01/2015	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 15/01/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 27/11/2025	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

<http://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-trial-looking-at-chemotherapy-and-surgery-for-early-bladder-cancer-caliber>

Contact information

Type(s)

Scientific

Contact name

Mr Steven Penegar

Contact details

Clinical Trials & Statistics Unit at the Institute of Cancer Research (ICR-CTSU)
The Institute of Cancer Research
London
United Kingdom
SM2 5NG

Additional identifiers

Clinical Trials Information System (CTIS)

2013-005095-18

ClinicalTrials.gov (NCT)

NCT02070120

Protocol serial number

17640

Study information

Scientific Title

A phase II randomised feasibility study of chemoresection or surgical management in low risk non muscle invasive bladder cancer

Acronym

CALIBER

Study objectives

Current hypothesis as of 10/04/2017:

To demonstrate that chemoresection has sufficient activity against NMIBC to warrant further investigation of its role as a potential alternative to surgical intervention for low risk NMIBC recurrence.

Previous hypothesis:

CALIBER aims to demonstrate that chemoresection will enable 60% of participants to avoid surgical intervention for low risk NMIBC recurrence, as assessed by response rate at 3 months.

Ethics approval required

Old ethics approval format

Ethics approval(s)

NRES Committee South Central - Hampshire B, 29/08/2014, ref: 14/SC/1223

Study design

Randomized; Interventional

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Cancer; Subtopic: Bladder Cancer; Disease: Bladder (superficial)

Interventions

Chemoresection group: 4 once weekly instillations of 40 mg MMC as outpatients

Surgical management group: Surgery according to local practice

Added 27/11/2025:

Additional Data Linkage Information:

Participants from this trial will also be included in the INTERACT project which will link to their data held by NHS England. For more information, please see the INTERACT website:

<https://www.icr.ac.uk/interact>.

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

Mitomycin C

Primary outcome(s)

Complete response to chemoresection 3 months post-treatment

Key secondary outcome(s)

Added 12/10/2016:

In the chemoresection group:

1. Treatment compliance

In both groups:

2. Time to recurrence in patients disease free at 3 months
3. Transurethral resection and biopsy rates
4. Progression-free survival
5. Toxicity
6. Quality of life
7. Health service utilisation

Completion date

30/09/2019

Eligibility

Key inclusion criteria

Current inclusion criteria as of 29/03/2017:

1. Written informed consent
2. NMIBC recurrence following original diagnosis of low risk NMIBC (defined as Ta G1 or Ta G2 (Ta low grade) with a risk of recurrence score of ≤ 6 using EORTC risk tables).
3. Histologically confirmed TCC at original diagnosis
4. Aged 16 or over
5. Satisfactory pre-treatment haematology values haemoglobin > 100 g/L and serum creatinine $< 1.5 \times \text{ULN}$
6. Negative pregnancy test for women of child-bearing potential

Previous inclusion criteria from 12/10/2016 to 29/03/2017:

1. Written informed consent
2. NMIBC recurrence following original diagnosis of low risk NMIBC (defined as Ta G1 or Ta G2 (Ta low grade) with a risk of recurrence score of ≤ 5 using EORTC risk tables).
3. Histologically confirmed TCC at original diagnosis
4. Aged 16 or over
5. Satisfactory pre-treatment haematology values haemoglobin > 100 g/L and serum creatinine $< 1.5 \times \text{ULN}$
6. Negative pregnancy test for women of child-bearing potential

Original inclusion criteria:

1. Written informed consent

2. NMIBC recurrence following original diagnosis of low risk NMIBC (defined as Ta G1 or Ta G2 NMIBC with a risk of recurrence score of ≤ 5 using EORTC risk tables)
3. Histologically confirmed TCC at original and any subsequent diagnoses
4. Aged 16 or over
5. Satisfactory pre-treatment haematology values Hb > 100 g/L and serum creatinine < 1.5xULN
6. Negative pregnancy test for women of child-bearing potential

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

16 years

Upper age limit

100 years

Sex

All

Total final enrolment

82

Key exclusion criteria

Current exclusion criteria as of 29/03/2017:

1. Any history of: grade 3/high grade or $\geq T1$ transitional cell carcinoma, concomitant carcinoma in situ, more than 7 tumours at one diagnosis or more than 1 recurrence per year since initial diagnosis or in the past five years, whichever is shorter
2. Any history of histologically confirmed non-TCC bladder cancer
3. Trial entry recurrence identified within 11.5 months of the date of the original diagnosis
4. Any prior treatment of the trial entry recurrence (including biopsy)
5. Previous MMC chemotherapy other than a single instillation at diagnostic surgery
6. Known allergy to MMC
7. Carcinoma involving the prostatic urethra or upper urinary tract (participants should have had imaging of the upper urinary tract within 2 years prior to randomisation)
8. Known or suspected reduced bladder capacity (<100ml)
9. Significant bleeding disorder
10. Female patients who are breast-feeding or are of childbearing potential and unwilling or unable to use adequate non-hormonal contraception. Male patients should also use contraception if sexually active.
11. Active or intractable urinary tract infection
12. Urethral stricture or anything impeding the insertion of a catheter
13. Large narrow neck diverticula
14. Significant urinary incontinence

15. Any other conditions that in the Principal Investigator's opinion would contraindicate protocol treatment

16. Unable or unwilling to comply with study procedures or follow up schedule

Previous exclusion criteria from 12/10/2016 to 29/03/2017:

1. Any history of: grade 3/high grade or T1 transitional cell carcinoma, concomitant carcinoma in situ, more than 7 tumours at one diagnosis or more than 1 recurrence per year since initial diagnosis or in the past five years, whichever is shorter

2. Any history of histologically confirmed non-TCC bladder cancer

3. Trial entry recurrence identified within 11.5 months of the date of the original diagnosis

4. Previous MMC chemotherapy other than a single instillation at diagnostic surgery

5. Known allergy to MMC

6. Carcinoma involving the prostatic urethra or upper urinary tract (participants should have had imaging of the upper urinary tract within 2 years prior to randomisation)

7. Known or suspected reduced bladder capacity (<100ml)

8. Significant bleeding disorder

9. Female patients who are breast-feeding or are of childbearing potential and unwilling or unable to use adequate non-hormonal contraception. Male patients should also use contraception if sexually active

10. Active or intractable urinary tract infection

11. Urethral stricture or anything impeding the insertion of a catheter

12. Large narrow neck diverticula

13. Significant urinary incontinence

14. Any other conditions that in the Principal Investigator's opinion would contraindicate protocol treatment

15. Unable or unwilling to comply with study procedures or follow up schedule

Original exclusion criteria:

1. Risk of recurrence score >5 at original or any subsequent diagnoses (including any history of: grade 3 or T1 transitional cell carcinoma, concomitant carcinoma in situ, more than 7 tumours at one diagnosis or more than 1 recurrence per year)

2. Previous MMC chemotherapy other than a single instillation at diagnostic surgery

3. Known allergy to MMC

4. Carcinoma involving the prostatic urethra or upper urinary tract (participants should have received an ultrasound of the upper urinary tract within 2 years prior to randomisation)

5. Known or suspected reduced bladder capacity (<100ml)

6. Significant bleeding disorder

7. Female patients who are breast-feeding or are of childbearing potential and unwilling or unable to use adequate non-hormonal contraception. Male patients should also use contraception if sexually active

8. Any other malignancy in the past 2 years (except: non-melanomatous skin cancer cured by excision, adequately treated carcinoma in situ of the cervix, DCIS/LCIS of the breast or prostate cancer in patients who have a life expectancy of >5 years upon trial entry)

9. Active or intractable urinary tract infection

10. Urethral stricture or anything impeding the insertion of a catheter

11. Large narrow neck diverticula

12. Significant urinary incontinence

13. Any other conditions that in the Principal Investigator's opinion would contraindicate protocol treatment

14. Unable or unwilling to comply with study procedures or follow up schedule

Date of first enrolment

28/01/2015

Date of final enrolment

04/09/2017

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Medway Maritime Hospital

Windmill Road

Gillingham

England

ME7 5NY

Study participating centre

Royal Bournemouth Hospital

Castle Lane East

Bournemouth

England

BH7 7DW

Study participating centre

Cumberland Infirmary

Newtown Road

Carlisle

England

CA2 7HY

Study participating centre

West Cumberland Hospital

Homewood Rd

Whitehaven

England

CA28 8JG

Study participating centre

Wythenshawe Hospital

Southmoor Road
Wythenshawe
England
M23 9LT

Study participating centre

Withington Community Hospital

Nell Lane
Manchester
England
M20 2LR

Study participating centre

Royal Devon and Exeter Hospital

Barrack Road
Exeter
England
EX2 5DW

Study participating centre

Royal Surrey County Hospital

Egerton Road
Guildford
England
GU2 7XX

Study participating centre

Basingstoke & North Hampshire Hospital

Aldermaston Road
Basingstoke
England
RG24 9NA

Study participating centre

St James's University Hospital

Beckett Street
Leeds
England
LS9 7TF

Study participating centre
Croydon University Hospital
London Road
Thornton Heath
England
CR7 7YE

Study participating centre
Freeman Hospital
Freeman Road
Newcastle upon Tyne
England
NE7 7DN

Study participating centre
Royal Oldham Hospital
Rochdale Road
Manchester
England
OL1 2JH

Study participating centre
Basingstoke & North Hampshire Hospital
Aldermaston Road
Basingstoke
England
RG24 9NA

Study participating centre
Ipswich Hospital
Heath Road
Ipswich
England
IP4 5PD

Study participating centre
Leicester General Hospital
Gwendolen Road
Leicester

England
LE5 4PW

Study participating centre
New Cross Hospital
Wednesfield Road
Wolverhampton
England
WV10 0QP

Study participating centre
Darent Valley Hospital
Darenth Wood Road
Dartford
England
DA2 8DA

Study participating centre
Broomfield Hospital
Court Road
Chelmsford
England
CM1 7ET

Study participating centre
James Cook University Hospital
Marton Road
Middlesbrough
England
TS4 3BW

Study participating centre
Hereford County Hospital
Union Walk
Hereford
England
HR1 2ER

Study participating centre

University College Hospital
250 Euston Road
London
England
NW1 2PG

Study participating centre
Pinderfields Hospital
Aberford Road
Wakefield
England
WF1 4DG

Study participating centre
Macclesfield District General Hospital
Victoria Road
Macclesfield
England
SK10 3BL

Study participating centre
Dorset County Hospital
Williams Avenue
Dorchester
England
DT1 2JY

Study participating centre
St Richard's Hospital
Spitalfield Lane
Chichester
England
PO19 6SE

Study participating centre
Worthing Hospital
Lyndhurst Road
Worthing
England
BN11 2DH

Study participating centre
Worcestershire Royal Hospital
Charles Hastings Way
Worcester
England
WR5 1DD

Study participating centre
Royal Hallamshire Hospital
Glossop Road
Sheffield
England
S10 2JF

Study participating centre
Royal Preston Hospital
Sharoe Green Lane North
Preston
England
PR2 9HT

Study participating centre
Cheltenham General Hospital
Sandford Road
Cheltenham
England
GL53 7AN

Study participating centre
Southampton General Hospital
Tremona Road
Southampton
England
SO16 6YD

Study participating centre
Derriford Hospital
Derriford Road
Plymouth

England
PL6 8DH

Study participating centre
Northwick Park Hospital
Watford Road
Harrow
England
HA1 3UJ

Study participating centre
Princess Alexandra Hospital
Hamstel Road
Harlow
England
CM20 1QX

Sponsor information

Organisation

The Institute of Cancer Research

ROR

<https://ror.org/043jzw605>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from caliber-icrctsu@icr.ac.uk.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2018		Yes	No
Results article	results	01/04/2018		Yes	No
Results article	12-month results	01/03/2019		Yes	No
Results article	results	01/06/2020		Yes	No
HRA research summary			28/06/2023	No	No
Other publications	recruitment aids	01/11/2016		Yes	No
Other publications	recruitment aids	01/05/2017		Yes	No
Protocol file	version 6	20/06/2017	18/08/2022	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes