# Patient controlled analgesia (PCA) versus routine care in the Emergency Department

Submission date 04/07/2011	Recruitment status  No longer recruiting	[X] Prospectively registered		
		[X] Protocol		
Registration date 04/07/2011	Overall study status Completed	Statistical analysis plan		
		[X] Results		
<b>Last Edited</b> 15/08/2016	Condition category Injury, Occupational Diseases, Poisoning	Individual participant data		

## Plain English summary of protocol

Background and study aims

Pain is very common in patients attending Emergency (A&E) Departments. Pain is sometimes difficult to treat, and individual patient satisfaction and levels of pain control also vary. In recent national surveys more than half of patients felt more could be done to treat their pain during their attendance. Normally when a patient is in severe pain in the Emergency Department they receive morphine, a strong painkiller, through a drip. Nurses give the injection, and then return after a while to see if any further treatment is needed. This might involve further doses of morphine or other types of painkiller if more appropriate. When they are admitted to a ward, it is more common for patients to be prescribed painkillers taken by mouth. A patient-controlled analgesia (PCA) device is a syringe that can be connected to a drip in the patients arm, which allows the patient to deliver their own painkiller into a vein by pressing a button attached to the syringe holder. It has a safety device to prevent too much drug being delivered. PCAs are commonly used in a variety of different settings in the hospital (typically after an operation), but they are not usually used for emergency patients. This study aims to see if giving patients a PCA machine in the Emergency Department, and during the first few hours of their stay in hospital, improves pain relief and satisfaction.

# Who can participate?

Patients aged between 18 and 75 with traumatic injuries or non-traumatic abdominal pain.

# What does the study involve?

Patients who agree to take part will be randomly allocated to receive either PCA or standard treatment involving nurses giving pain relief drugs then returning after a period of time to see whether any further medication is required. In addition, all patients in the study will be offered other forms of painkillers as necessary. By using the two methods of administration we will be able to assess whether there is benefit in terms of pain control and satisfaction with treatment between the two methods of managing pain.

What are the possible benefits and risks of participating? We feel that by giving patients control of their own pain relief, they may feel their pain is better managed.

Where is the study run from? Derriford Hospital, Plymouth (UK)

When is the study starting and how long is it expected to run for? July 2011 to January 2013

Who is funding the study? National Institute for Health Research (NIHR) (UK)

Who is the main contact? Dr Jason Smith jasonesmith@nhs.net

# Contact information

# Type(s)

Scientific

#### Contact name

Dr Jason Smith

#### Contact details

Derriford Hospital Derriford Road Plymouth United Kingdom PL6 8DH

•

jasonesmith@nhs.net

# Additional identifiers

Protocol serial number 10515

# Study information

### Scientific Title

An open randomised trial of patient controlled analgesia (PCA) versus routine care in the Emergency Department

#### Acronym

PAin SoluTions In the Emergency Setting (PASTIES)

### Study objectives

Pain is an extremely common presentation to Emergency Departments (EDs), but is often difficult to treat effectively (almost half of patients recently surveyed thought more could be done to treat their pain in the ED). Routine treatment of severe pain involves intermittent doses of intravenous morphine, administered by nurses.

Patient controlled analgesia (PCA) is effective in many clinical settings, but there is very little evidence relating to its use in the ED. No previous study has investigated the issue of continuing a PCA from the emergency department and during the first few hours of a patient's admission to hospital, to optimise their pain relief.

The aim of this study is to compare PCA with routine care (nurse-titrated analgesia) in adult emergency patients who present to the Emergency Department in severe pain from traumatic injuries or non-traumatic abdominal pain, and are then admitted to an inpatient ward. We plan to undertake a non-blinded randomised trial of PCA versus routine care in these patients. The primary outcome is a selfadministered visual analogue scale pain score, completed hourly for 12 hours. Secondary outcomes include total opioid dose and other analgesic use, adverse effects, and patient satisfaction. An economic evaluation will compare PCA with standard care in a cost-effectiveness analysis.

Added 31/03/2014: The study is conceived as two distinct trials within one study protocol, having been powered separately for the two different populations of patients; the results will accordingly be analysed and presented separately.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

South Central REC (Southampton), May 2011, ref: 11/SC/0151

### Study design

Randomised; Interventional; Design type: Not specified, Treatment

# Primary study design

Interventional

# Study type(s)

Treatment

# Health condition(s) or problem(s) studied

Topic: Injuries and Emergencies; Subtopic: Injuries and Emergencies (all Subtopics); Disease: Injuries and Emergencies

### **Interventions**

Number of participants in each group - 400 total, i.e. 200 with non-traumatic abdominal pain and 200 with musculoskeletal injury, half of whom will receive PCA and half will receive standard care.

Patients will be randomised to either standard care involving bolus intravenous (IV) morphine and multimodal analgesia, or PCA. PCA is maximum 1mg morphine per 5 minutes with lock out on PCA device.

### Intervention Type

Drug

#### Phase

Not Applicable

## Drug/device/biological/vaccine name(s)

Morphine

### Primary outcome(s)

Visual Analogue Scale (VAS) Pain Score; Timepoint(s): hourly for 12 hours

### Key secondary outcome(s))

- 1. Total opioid dose and use of other analgesics
- 2. Opioid side-effects
- 3. Patient satisfaction with pain management
- 4. Proportion of study period with VAS >44mm
- 5. Proportion of study period spent sleeping
- 6. Length of hospital stay
- 7. Incremental cost effectiveness ratio (ICER)

### Completion date

03/01/2013

# Eligibility

### Key inclusion criteria

- 1. Adult patients aged between 18 and 75 years of age inclusive
- 2. Traumatic injuries or non-traumatic abdominal pain
- 3. In severe pain on arrival in the Emergency Department
- 4. Admission to hospital is intended at the time of enrolment
- 5. Provision of informed consent to participate.; Target Gender: Male & Female; Upper Age Limit 75 years; Lower Age Limit 18 years

# Participant type(s)

Patient

### Healthy volunteers allowed

No

### Age group

Adult

### Lower age limit

18 years

### Sex

All

### Key exclusion criteria

- 1. Patients over 75 years
- 2. Patients with a reduced conscious level
- 3. Inability to operate a PCA device
- 4. Patients who cannot understand the study information (e.g. due to pre-existing dementia, learning difficulties, or intoxication)
- 5. Patients with chronic pain

- 6. Patients who are opioid tolerant
- 7. Patients with active opioid addiction
- 8. Patients with a history of renal failure
- 9. Allergy or other contraindication to morphine
- 10. Hypotension (systolic blood pressure <90mmHg)
- 11. Patients in police custody, or prisoners
- 12. Inability to gain intarvenous (IV) access
- 13. Patients who are likely to be definitively treated in the ED and discharged
- 14. Patients who are pregnant or breastfeeding
- 15. Open fractures (excluded because this group of patients will undergo surgery within six hours of injury according to national standards)
- 16. Patients on other predetermined analgesia pathway (e.g. regional anaesthesia)
- 17. Previous participation in this study
- 18. Current participation in another Clinical Trial of an Investigational Medicinal Product (CTIMP)

#### Date of first enrolment

04/07/2011

### Date of final enrolment

03/01/2013

# Locations

### Countries of recruitment

United Kingdom

**England** 

# Study participating centre Derriford Hospital

Plymouth United Kingdom PL6 8DH

# Sponsor information

### Organisation

Plymouth Hospitals NHS Trust (UK)

#### **ROR**

https://ror.org/05x3jck08

# Funder(s)

# Funder type

Government

# Funder Name

National Institute for Health Research (NIHR) (UK) - Research for Patient Benefit (RfPB) programme

# **Results and Publications**

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Not provided at time of registration

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	21/06/2015		Yes	No
Results article	results	21/06/2015		Yes	No
Protocol article	protocol	14/02/2013		Yes	No
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes