# Topical use of adrenaline in different concentrations for endoscopic sinus surgery

Submission date	Recruitment status	<ul><li>Prospectively registered</li></ul>
18/12/2007	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
07/01/2008	Completed	Results
Last Edited	Condition category	Individual participant data
10/03/2008	Ear, Nose and Throat	Record updated in last year

#### Plain English summary of protocol

Not provided at time of registration

## Contact information

#### Type(s)

Scientific

#### Contact name

Dr Krishnamurti Sarmento Junior

#### Contact details

Praia de Botafogo, 422 / 1106 Botafogo Rio de Janeiro Brazil +55 22250 040 krishnamurti.sarmento@gmail.com

#### Additional identifiers

**Protocol serial number** N/A

# Study information

Scientific Title

**Study objectives** 

Vasoconstrictors such as cocaine or epinephrine (adrenaline) are routinely applied during functional endoscopic sinus surgery. The goal is to control bleeding because, when surgery is endoscopic, even minor bleeding can be troublesome if it blocks the small end of the endoscope. Nevertheless vasoconstrictors might cause an increase in blood pressure and heart arrhythmias. The ideal adrenaline concentration should be the one to provide optimal operative field with no cardiovascular side effects. Despite its routine use, that concentration is yet to be determined. Concentrations of adrenaline varying from 1:1000 to 1:200,000 have been used both topically and sub-mucosally. Many studies have used both adrenaline and cocaine, which makes it difficult to establish the individual role of each substance on the outcome. We have decided to work with adrenaline solutions applied only topically.

We study the effects of topical use of adrenaline solution on the nasal mucosa in three different concentrations on systemic absorption of the drug, blood pressure, heart rhythm, operative bleeding and operative time during endoscopic sinus surgery for nasal polyposis.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Approved by the ethic committee of Federal University of Rio de Janeiro (UFRJ) on 24 November 2004 (ref: 207/04 CEP)

#### Study design

Randomised double-blind single-centre controlled trial.

#### Primary study design

Interventional

#### Study type(s)

**Not Specified** 

#### Health condition(s) or problem(s) studied

Nasal polyposis

#### Interventions

Patients were submitted to endoscopic sinus surgery under general anesthesia for treatment of nasal polyposis using different concentrations of adrenaline solution applied topically to the nasal mucosa.

#### Concentrations:

- 1. Adrenaline 1:2000 with lidocaine 1% (4 ml)
- 2. Adrenaline 1:10,000 with lidocaine 1% (4 ml)
- 3. Adrenaline 1:50,000 with lidocaine 1% (4 ml)

The control group consisted of patients submitted to tonsillectomies using the same anesthetic protocol as the study groups but without the use of adrenaline solution during surgery.

#### Intervention Type

Drug

#### **Phase**

#### Drug/device/biological/vaccine name(s)

adrenaline

#### Primary outcome(s)

The following data were collected during surgery (No assessment was done post-operatively):

- 1. Variation of blood pressure during surgery (measured by non-invasive blood pressure measures every 3 minutes)
- 2. Variation of heart frequency during surgery (measured continuously by 12 lead ECG)
- 3. Development of heart arrythmias during surgery (measured continuously by 12 lead ECG)
- 4. Operative bleeding (measured both objectively by the amount of blood aspirated during the procedure and subjectively by visual-analogue scale from 01 = no bleeding to 10 = very high bleeding)
- 5. Variation of plasma levels of adrenaline measured by three blood samples obtained during each surgery
- 6. Operative time

#### Key secondary outcome(s))

No secondary outcome measures

#### Completion date

28/02/2007

# Eligibility

#### Key inclusion criteria

- 1. Be over 18 years old
- 2. Need elective endoscopic sinus surgery under general anesthesia for the treatment of nasal polyposis
- 3. Understand and give written consent to participate in the trial
- 4. To be classified as grade I or II according to the American Society of Anesthesiology classification of pre-operative assessment

#### Participant type(s)

**Patient** 

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

Αll

#### Key exclusion criteria

- 1. Patients legally incompetent or unable, for any reason, to read and sign a written consent form
- 2. Patients who withdraw their consent in any time during the course of the trial, even if they have signed the consent form
- 3. Patients with systemic hypertension, coronary diseases, heart arrythmia, coagulation disorders, collagen disorders, renal or liver insufficiency in any degree, previously diagnosed or detected during pre-operative assessment
- 4. Pregnant women
- 5. Patients who do not follow the trial protocol for any reason, unless it is related to the use of the adrenaline solution
- 6. Patients in use of a number of medications that might interfere with blood pressure or coagulaion, such as anti-inflammatory drugs, oral hypoglicemics, beta agonists, etc

# Date of first enrolment 01/03/2005

Date of final enrolment 28/02/2007

## Locations

# Countries of recruitment Brazil

Study participating centre
Praia de Botafogo, 422 / 1106
Rio de Janeiro
Brazil
+55 22250 040

## Sponsor information

#### Organisation

Federal University of Rio de Janeiro (UFRJ) (Brazil)

#### **ROR**

https://ror.org/03490as77

# Funder(s)

#### Funder type

University/education

#### Funder Name

Federal University of Rio de Janeiro (UFRJ) (Brazil)

# **Results and Publications**

Individual participant data (IPD) sharing plan

**IPD sharing plan summary**Not provided at time of registration