# Effect of 2 versus 3 pneumococcal conjugate vaccinations Prevnar on nasopharyngeal carriage, transmission and herd immunity; a randomized, controlled study.

Submission date	<b>Recruitment status</b> No longer recruiting	Prospectively registered		
20/12/2005		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
20/12/2005		[X] Results		
Last Edited	Condition category	Individual participant data		
31/05/2019	Infections and Infestations			

## Plain English summary of protocol

Not provided at time of registration

## Contact information

## Type(s)

Scientific

#### Contact name

Prof E.A.M. Sanders

#### Contact details

University Medical Center Utrecht
Department of Pediatric Immunology
HP 06.063.0
P.O. Box 85090
Lundlaan 6
Utrecht
Netherlands
3508 AB
+31 (0)30 2504000
l.sanders@umcutrecht.nl

## Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

## ClinicalTrials.gov (NCT)

NCT00189020

#### Protocol serial number

MINOES 01, STEG R05 008

## Study information

#### Scientific Title

Effect of 2 versus 3 pneumococcal conjugate vaccinations Prevnar on nasopharyngeal carriage, transmission and herd immunity; a randomized, controlled study.

#### **Acronym**

**MINOES** 

## **Study objectives**

Study hypotheses amended as of 22/05/2007:

- 1. Two vaccinations with 7-valent conjugate vaccine before six months of age will protect against invasive pneumococcal disease in the first 18 to 24 months of life
- 2. Two vaccinations at 2 and 4 months of age without a booster vaccination at 11 months will have less effect on vaccine-type pneumococcal carriage reduction and therefore also herdimmunity and respiratory tract infections but also less pneumococcal replacement by non-vaccine types and other potentially pathogenic bacteria
- 3. Anti-pneumococcal antibody development after two vacinations with 7-valent conjugate vacine before six months of age and revaccination at 24 months of age will be equal or better compared with antibody development after two vaccination before six months of age followed by booster vaccination at 11 and 24 months of age. Later vaccination will improve long term protection

## Previous study hypothesis:

- 1. Two vaccinations before the age of 6 months with the pneumococcal conjugate vaccine Prevnar in infants will protect the children against invasive pneumococcal disease.
- 2. Two vaccinations at 2 and 4 months of age without a booster vaccination at 11 months will have less effect on vaccine-type pneumococcal carriage reduction and therefore also herdimmunity and respiratory tract infections but also less pneumococcal replacement by non-vaccine types and other potentially pathogenic bacteria.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approval received from the local medical ethics board (Stichting Therapeutische Evaluatie Geneesmiddelen [STEG], acknowledged by the Centrale Commissie Mensgebonden Onderzoek [CCMO], The Netherlands) on the 21st June 2005 (ref: STEG R05.008, version 6).

First and second amendment (collection of saliva for immunological analysis and extension microbiology research of collected nasopharyngeal swabs in subgroups of children) approved on the 11th September 2006.

Third amendment (addition of booster vaccination at 24 months and blood sampling) approved on the 2nd March 2007 (ref: STEG R05-008, MINOES version 8 dd february 2007).

### Study design

Randomised controlled trial

## Primary study design

Interventional

## Study type(s)

**Not Specified** 

## Health condition(s) or problem(s) studied

Invasive pneumococcal disease, Respiratory tract infection.

### Added as of 22/05/2007:

Change in: nasopharyngeal carriage after reduced doses 7-valent conjugate vaccine, antipneumococcal antibody development, family transmission of pneumococci

#### **Interventions**

Three groups of each 333 newborns:

Group I: Prevnar at age 2 and 4 months

Group II: Prevnar at age 2, 4 and 11 months

Group III: Prevnar at age 24 months

#### Amendment made as of 22/05/2007:

Groups I and II will receive booster vaccination 7-valent conjugate vaccine at 24 months of age.

## Intervention Type

Drug

#### Phase

**Not Specified** 

## Drug/device/biological/vaccine name(s)

Pneumococcal conjugate vaccine Prevnar

## Primary outcome(s)

Pneumococcal nasopharyngeal carriage before 2 months of age, and at 6, 12, 18 and 24 months of age in vaccinated children (groups I and II) and controls (group III), and NP pneumococcal carriage of siblings and parents/caregivers when the baby is 12 and 24 months of age (herd immunity).

## Key secondary outcome(s))

- 1. Determination of pneumococcal (vaccine and non-vaccine serotypes) colonization of infants in the Netherlands before introduction of Prevnar in the National Vaccination Program (group III)
- 2. Evaluation of replacing pneumococcal serotypes after Prevnar vaccinations
- 3. Influence of Prevnar vaccinations on other colonizing bacterial species like Staphylococcus, H. influenzae, M. catarrhalis and Streptococcus
- 4. Anti pneumococcal antibody levels at 12 and 24 months after 2 doses as compared to 2+1 Prevnar vaccinations
- 5. Relation between invasive pneumococcal disease and pneumococcal colonization in the Netherlands

6. Evaluation of potential reduction of physician-diagnosed acute otitis media and lower respiratory tract infections after pneumococcal conjugate vaccinations

## Added as of 22/05/2007:

- 7. Comparison of anti-pneumococcal antibody levels after vaccinations at 2, 4 and 24 months, at
- 2, 4, 11 and 24 months and after primo-vaccination at 24 months of age in subgroups of 80 children of groups I, II and III
- 8. Comparison of B-memory cells after vaccination at 24 months of age after vaccination schemes of groups I, II and III

## Completion date

01/03/2008

# **Eligibility**

## Key inclusion criteria

1000 healthy newborns (and family members) who will receive childhood vaccinations according to the national vaccination program, starting at 2 months of age.

## Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Neonate

#### Sex

All

#### Total final enrolment

1003

## Key exclusion criteria

Exclusion from the national vaccination program because of the presence of a medical condition requiring treatment that can interfere with the results of vaccinations, known of suspected allergy to components of the vaccine, known or suspected immunodeficiency disease other than IgA or IgG-subclass deficiency, previous treatment with plasma or immunoglobulins, previous vaccinations other than hepatitis B vaccinations, coagulations disorders

## Date of first enrolment

20/06/2005

#### Date of final enrolment

01/03/2008

## Locations

#### Countries of recruitment

Study participating centre
University Medical Center Utrecht
Utrecht
Netherlands
3508 AB

# Sponsor information

## Organisation

University Medical Centre Utrecht (UMCU) (Netherlands)

#### **ROR**

https://ror.org/04pp8hn57

# Funder(s)

## Funder type

Government

#### **Funder Name**

The Netherlands Ministry of Health, Welfare and Sport (The Netherlands)

## **Results and Publications**

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2011	31/05/2019	Yes	No
Results article	results	01/12/2014	31/05/2019	Yes	No
Results article	results	01/08/2014	31/05/2019	Yes	No
Results article	results	02/12/2013	31/05/2019	Yes	No

Results article	results	01/12/2011	31/05/2019	Yes	No
Results article	results	01/02/2014	31/05/2019	Yes	No
Results article	results	08/07/2009	31/05/2019	Yes	No
Results article	results	08/09/2010	31/05/2019	Yes	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes