

# Nebulised procaterol versus nebulised salbutamol for the treatment of moderate acute asthma

<b>Submission date</b> 23/07/2009	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 05/08/2009	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 05/08/2009	<b>Condition category</b> Respiratory	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Hadiarto Mangunegoro

**Contact details**  
Department of Pulmonology  
Faculty of Medicine  
University of Indonesia  
Persahabatan Hospital  
Jl. Persahabatan Raya No. 1  
Jakarta  
Indonesia  
13230

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

## Study information

### Scientific Title

The efficacy of nebulised procaterol versus nebulised salbutamol for the treatment of moderate acute asthma: a randomised, double-blind, parallel group study

### Study objectives

The efficacy of nebule procaterol in improving peak expiratory flow rate (PEFR) in moderate acute asthma patients is superior to nebule salbutamol.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

1. Ethical clearance and registration in local (Indonesian) regulatory authority approved 7th March 2006
2. Ethics Committee of the Medical Research Ethics of the Faculty of Medicine, University of Indonesia approved on 9th October 2006

### Study design

Randomised double-blind two-arm parallel group phase III clinical trial

### Primary study design

Interventional

### Secondary study design

Randomised controlled trial

### Study setting(s)

Hospital

### Study type(s)

Treatment

### Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

### Health condition(s) or problem(s) studied

Moderate acute asthma

### Interventions

Treatment allocation was according to block randomisation using random permuted blocks of size 4. Patients received either one ampule of procaterol, diluted with 2 ml of normal saline, or one ampule of salbutamol three times every 20 minutes (at 0, 20 and 40 minutes). Both drugs were administered via jet-type nebuliser (Pulmoaid™). The PEFR was measured 20, 40, 60 and 120 minutes. At the same time, vital signs, asthma score and adverse events were evaluated. At 120 minutes, the blood gas analysis and the electrocardiogram (ECG) were repeated. Then, the

patient was observed for adverse event(s) that might occur until 280 minutes. Afterwards, the patient was discharged from the study and received 6 tablets of bronchodilator (salbutamol) and 6 tablets of corticosteroid (methylprednisolone).

**Intervention Type**

Drug

**Phase**

Phase III

**Drug/device/biological/vaccine name(s)**

Procaterol, salbutamol, methylprednisolone

**Primary outcome measure**

Difference from baseline in peak expiratory flow rate (PEFR). Measurements performed at 0, 20, 40, 60 and 120 minutes.

**Secondary outcome measures**

Difference from baseline in asthma score. Measurements performed at 0, 20, 40, 60 and 120 minutes.

**Overall study start date**

12/06/2007

**Completion date**

24/04/2008

**Eligibility****Key inclusion criteria**

1. Patients with moderate acute asthma according to Global Initiative for Asthma (GINA) 1998 (patients with asthma score 5 to 11; PEFR less than or same 80% predicted)
2. Patients of both genders, aged 15 to 60 years
3. Patients still have the ability to undergo examinations and give written informed consent

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

140

**Key exclusion criteria**

1. Pregnant and lactating women
2. Smokers

3. Patients with heart disease, hyperthyroidism, diabetes mellitus, chronic obstructive pulmonary disease (COPD) or other chronic diseases

4. Patients with signs of severe infections

**Date of first enrolment**

12/06/2007

**Date of final enrolment**

24/04/2008

## **Locations**

**Countries of recruitment**

Indonesia

**Study participating centre**

**Department of Pulmonology**

Jakarta

Indonesia

13230

## **Sponsor information**

**Organisation**

Otsuka Indonesia

**Sponsor details**

Perkantoran Hijau Arkadia

Tower A, 3rd Floor

Jl. Letjen T.B. Simatupang Kav. 88

Jakarta

Indonesia

12520

**Sponsor type**

Industry

**Website**

<http://www.otsuka.co.id/>

**ROR**

<https://ror.org/01ambd593>

# **Funder(s)**

**Funder type**

Industry

**Funder Name**

Otsuka Indonesia

## **Results and Publications**

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Not provided at time of registration