

Esketamine for reducing perioperative opioid use in laparoscopic colorectal cancer surgery: impact on postoperative nausea and vomiting at 48 hours

Submission date 08/09/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 26/09/2024	Overall study status Completed	<input checked="" type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 14/11/2025	Condition category Digestive System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Colorectal cancer (CRC) is the third most common malignant tumor in the world, and surgical resection is the most effective treatment for CRC. Postoperative nausea and vomiting (PONV) is a common and painful complication after surgery. Opioids, the most commonly used intravenous analgesics for general anesthesia, are considered to be an independent risk factor for PONV, and perioperative reduction of opioid use can reduce the occurrence of PONV. Esketamine has sedative, analgesic and hypnotic effects, and is widely used in clinical anesthesia, which can completely or partially replace opioids. In this study, esketamine, an intravenous anesthetic, will be used for anesthesia and analgesia during the perioperative period of laparoscopic radical resection of colorectal cancer, partially replacing the role of opioids. By observing the occurrence of PONV in patients, the influence of opioid reduction anesthesia based on esketamine on PONV in patients undergoing laparoscopic radical resection of colorectal cancer will be evaluated, to guide clinical practice and optimize anesthesia medication plan.

Who can participate?

Patients aged between 18 and 80 years old undergoing laparoscopic radical resection of colorectal cancer

What does the study involve?

Before entering the hospital, patients were forbidden to drink for 2 h and fast for 6 h, and electrocardiogram, hematuria routine, electrolyte and related tests will be undertaken. After entering the room, the venous channel is opened routinely, and the heart rate, blood pressure, electrocardiogram and pulse oxygen saturation are monitored and oxygen is given. Anesthesia and postoperative analgesia will be performed in combination with esketamine. PONV will be actively observed, and a rating scale will be used at 1, 24 and 48 h after surgery to record the amount of postoperative relief analgesia. Venous blood will be taken before surgery, 1 h during surgery, and 1, 24 and 72 h after surgery to detect inflammation indicators. The occurrence of postoperative adverse reactions such as hallucinations or nightmares will be recorded.

What are the possible benefits and risks of participating?

The protocol of this clinical study has been reviewed by the Medical Ethics Committee of the hospital to protect the rights and interests of the subjects to the greatest extent. The research doctors will pay close attention to and deal with PONV and pain, to improve the comfort of the perioperative period to a certain extent.

The drug used in this study, esketamine hydrochloride, has been approved by the State Food and Drug Administration, and the adverse reactions are controllable. Common adverse reactions mainly include agitation, increased secretions, nausea, transient increased heart rate, transient elevated blood pressure, diplopia, dreaming, dizziness, etc., but the general degree is mild and tolerable.

Where is the study run from?

The Department of Anesthesiology at Changzhou First People's Hospital, China

When is the study starting and how long is it expected to run for?

September 2023 to September 2024

Who is funding the study?

The Department of Anesthesiology at Changzhou First People's Hospital, China. Zhao Xin, the attending doctor, was the main contact for this research and provided funds for the research.

Who is the main contact?

Zhao Xin, 644388187@qq.com

Contact information

Type(s)

Public, Principal investigator

Contact name

Mr Zhao Xin

Contact details

The First People's Hospital of Changzhou, 185 Guqian Street, Tianning District
Changzhou City, Jiangsu Province
China
213000
+86 13775127431
644388187@qq.com

Type(s)

Scientific

Contact name

Dr Qiu Lan

Contact details

The First People's Hospital of Changzhou, 185 Guqian Street, Tianning District
Changzhou City, Jiangsu Province

China
213000
+86 13915089989
772117973@qq.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Study information

Scientific Title

Effect of esketamine-based opioid reduction anesthesia on postoperative nausea and vomiting in patients undergoing laparoscopic radical resection of colorectal cancer

Study objectives

In this study, esketamine was used for anesthesia and analgesia during the perioperative period of laparoscopic radical resection of colorectal cancer, partially replacing the role of opioids. By observing the occurrence of postoperative nausea and vomiting (PONV) in patients, the influence of opioid reduction anesthesia based on esketamine on PONV in patients undergoing laparoscopic radical resection of colorectal cancer was evaluated, to guide clinical practice and optimize anesthesia medication plan.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 13/09/2023, Ethics Committee of Changzhou First People's Hospital (185 Guqian Street, Tianning District, Changzhou, Jiangsu Province, 213000, China; +86 0519-68870965; xyyirb@163.com), ref: (2023) Teach No. 056

Study design

Single-center double-blind randomized controlled case study

Primary study design

Interventional

Study type(s)

Efficacy

Health condition(s) or problem(s) studied

Postoperative nausea and vomiting

Interventions

Participants will be randomly divided into two groups, the esketamine group (Group K) and the control group (Group C). The drugs required for anesthesia induction will be formulated

according to the dosage in the trial protocol. Esketamine 50mg for anesthesia induction and maintenance in Group K will be diluted to 10 ml and 50 ml with normal saline, respectively, while normal saline will be used as a control in Group C.

Both groups will be given midazolam 0.05 mg/kg, propofol 1.5-2 mg/kg, and cisatracurium benzoate 0.2 mg/kg for anesthesia induction. The esketamine group (group K) will be given intravenous esketamine 0.15 mg/kg (0.03ml/kg) and sufentanil 0.2 µg/kg. The control group (group C) will be given intravenous sufentanil 0.4 µg/kg and normal saline 0.03 ml/kg.

Patients in both groups will be injected with propofol 4-8 mg/kg·h, remifentanyl 0.05-0.2 µg/kg·min, and intermittent intravenous injection of cisatracurium benzoate 0.03 mg/kg as needed to maintain anesthesia. Group K will be injected with 0.15 mg/kg·h (0.15 ml/kg·h) of esketamine until 30 minutes before the end of the operation. Group C will be pumped with saline at the same rate. When SPI > 50 for more than 1min or the value increased by more than 10, sufentanil will be added 0.1µg/kg to maintain SPI: 20-50. At the same time, the pumping rate of propofol will be adjusted to maintain BIS: 40-60.

In both groups, propofol and remifentanyl will be stopped and intravenous analgesia pumps connected during suture. PCIA formula Group K esketamine 1 mg/kg, sufentanil 1 µg/kg and palonosetron hydrochloride 0.25 mg, diluted to 100ml with normal saline; Group C Sufentanil 2 µg/kg, palonosetron hydrochloride 0.25mg, diluted to 100ml with normal saline. The intravenous analgesia pumps in both groups were set as follows: the first dose will be 2ml, the background dose will be 2.0 ml/h, the automatic dose will be 2ml/ time, and the locking time will be 15min.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Esketamine

Primary outcome(s)

The occurrence and intensity of postoperative nausea and vomiting (PONV) measured using the PONV Impact Scale at 1, 24, and 48 h after surgery

Key secondary outcome(s))

1. The intraoperative dosage of opiates and propofol measured using data recorded in patient medical records throughout the procedure
2. Blood pressure, heart rate, bispectral index (BIS) value and surgical pleth index (SPI) value measured using data recorded in patient medical records collected before the operation, immediately after intubation, pneumoperitoneum establishment, 0.5h during operation, 1h during operation, 1.5h during operation, and at the end of the operation
3. Serum IL-6, IL-10, TNF-α, gastrin, motilin and substance P levels measured using ELISA in venous blood collected from patients in T0 (before surgery), T1 (1h during surgery), T2 (1h after surgery), T3 (24h after surgery) and T5 (72h after surgery)
4. The quality of postoperative recovery measured using the recovery scale QoR-15 before and 48 hours after surgery
5. Pain level at rest and exercise measured using a visual analog scale (VAS) at 1h, 24h and 48h after surgery

6. Dizziness, diplopia, hallucination or nightmare and other adverse reactions measured using data collected in patient medical records at 1 h, 24 h and 48 h after surgery

Completion date

12/09/2024

Eligibility

Key inclusion criteria

1. Patients undergoing laparoscopic radical resection of colorectal cancer
2. Aged 18-80 years old, no gender limitation
3. ASA I-III
4. Postoperative agreed to use the PCIA
5. Voluntarily signed informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

80 years

Sex

All

Total final enrolment

84

Key exclusion criteria

1. People with severe consciousness disorder or mental system disease and cognitive dysfunction
2. Merge heart liver and kidney and other viscera function disorder
3. Arrhythmia (atrial fibrillation or atrioventricular conduction block between arrhythmia etc.) or after cardiac pacemaker implantation
4. BP or 180/100 MMHG, untreated or poor control of blood pressure
5. Increased intracranial pressure
6. Glaucoma
7. Without treatment, and treatment of hyperthyroidism
8. Known for esketamine injection, the active ingredient or complementary makings allergies
9. Alcohol or alcohol addiction
10. Have a vision, hearing, or language barrier and cannot communicate and cooperate

Date of first enrolment

01/10/2023

Date of final enrolment

01/08/2024

Locations

Countries of recruitment

China

Study participating centre

Changzhou First People's Hospital

185 Guqian Street, Tianning District

Changzhou City, Jiangsu Province

China

213000

Sponsor information

Organisation

Changzhou Science and Technology Bureau

ROR

<https://ror.org/00wgk5w17>

Funder(s)

Funder type

Government

Funder Name

Changzhou Municipal Science and Technology Bureau

Alternative Name(s)

, CZST

Funding Body Type

Government organisation

Funding Body Subtype

Local government

Location
China

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Data sharing statement to be made available at a later date

Study outputs					
Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results	version 2025	14/11/2025	14/11/2025	No	No
Plain English results			14/11/2025	No	Yes
Statistical Analysis Plan			14/11/2025	No	No