

Surgical Wounds Healing By Secondary Intention - 2

Submission date 15/02/2019	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 25/03/2019	Overall study status Completed	<input checked="" type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 23/10/2025	Condition category Surgery	<input checked="" type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

After an operation, most wounds are closed using stitches or staples. Some wounds cannot be closed in this way and are left open. Sometimes wounds that have been closed may open up again. These “open” wounds are usually left to heal, over time, from the bottom up rather than attempting to close them again by some other means (healing by secondary intention). The most common treatment for these wounds is plain dressings. Another type of treatment is Negative Pressure Wound Therapy (NPWT) which is a relatively new treatment for open surgical wounds. It uses a small machine to apply suction to a wound through a special dressing. Use of NPWT has become more common and is used in around one third of people with open surgical wounds. It is not known which of these (NPWT or wound dressings) is the most effective treatment for surgical wounds healing by secondary intention and which treatment is best value for money. The aim of this study is to compare NPWT to normal wound dressings to see if it makes any difference to how quickly these open wounds heal.

Who can participate?

Patients aged 16 years or older with a surgical wound healing by secondary intention

What does the study involve?

Participants receive one of the two treatments (NPWT or wound dressings), selected at random using a computer system. The two groups are compared over 12 months including how long it takes their wounds to heal, number of infections, hospital admissions and further operations, and how much both treatments cost.

What are the possible benefits and risks of participating?

The information from this study may help to treat people with open wounds more effectively in the future. As with any treatment there are always potential risks, although side effects in both treatments are very uncommon. Where negative pressure wound therapy machines are used, the device may present a trip hazard and so care should be taken when moving around. Where a wound is located on the stomach, there is an increased risk of developing a fistula, an abnormal opening between organs and the skin. Where normal wound dressings are used, there may be a

need for frequent dressing changes, sometimes daily. These are usually completed by a nurse. Being in this study will not harm or disadvantage participant's care in any way and participants will be monitored regularly as part of usual NHS care.

Where is the study run from?
University of York (UK)

When is the study starting and how long is it expected to run for?
November 2018 to January 2024

Who is funding the study?
National Institute for Health Research (NIHR) (UK)

Who is the main contact?
Catherine Arundel
catherine.arundel@york.ac.uk

Contact information

Type(s)
Scientific

Contact name
Ms Catherine Arundel

ORCID ID
<https://orcid.org/0000-0003-0512-4339>

Contact details
York Trials Unit
Lower Ground Floor
ARRC Building
Department of Health Science
University of York
Heslington
York
United Kingdom
YO10 5DD
+44 (0)1904 321 116
catherine.arundel@york.ac.uk

Type(s)
Public

Contact name
Ms Catherine Arundel

ORCID ID
<https://orcid.org/0000-0003-0512-4339>

Contact details

York Trials Unit
Lower Ground Floor
ARRC Building
Department of Health Science
University of York
Heslington
York
United Kingdom
YO10 5DD
+44 (0)1904 321 116
catherine.arundel@york.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Sponsor ID: R2319; HTA 17/42/94; 40908

Study information

Scientific Title

A pragmatic, multicentre, randomised controlled trial to assess the clinical and cost effectiveness of negative pressure wound therapy versus usual care for surgical wounds healing by secondary intention (SWHSI 2)

Acronym

SWHSI-2

Study objectives

Negative Pressure Wound Therapy (NPWT) is superior to usual care in the treatment of surgical wounds healing by secondary intention (SWHSI).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 05/04/2019, Yorkshire and The Humber - Leeds East Research Ethics Committee (NHSBT Newcastle Blood Donor Centre, Holland Drive, Newcastle upon Tyne, NE2 4NQ; +44(0) 207 1048 088; nrescommittee.yorkandhumber-leedseast@nhs.net), ref: 19/YH/0054

Study design

Pragmatic multi-centre randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Surgical wounds healing by secondary intention

Interventions

Current intervention as of 01/04/2022:

Patients with a potential planned SWHSI (pre-operatively) or a SWHSI occurring at any point following surgery will be screened for potential eligibility by their clinical care team or GP.

Patients who agree to take part will receive one of the two treatments selected at random using a computer system.

Intervention: Negative Pressure Wound Therapy (NPWT).

Participants will be treated with negative pressure wound therapy (NPWT). The patient's wound is filled with a suitable dressing, and a liner may also be used to protect the wound. A vacuum pump is then attached which applies suction to the wound. A disposable, plastic canister slots into the pump to collect wound exudate and this is removed and replaced either when it becomes full, or at least once a week.

The device is generally used as part of the SWHSI treatment pathway rather than to the point of healing and is administered by both nurses and clinicians.

The use of any CE marked NPWT device, providing pressure of 60-150mmHg, in use within the NHS will be permitted in this trial, given that the principles of any device are similar and there is no evidence to suggest clinical or cost-effectiveness differences between devices. The device will be used in accordance with manufacturer guidance, and the clinical care team, in conjunction with local treatment guidelines, will determine the duration of device use, and whether this includes continuous or intermittent pressure cycles.

Control: Usual Care (no NPWT)

Usual care will be that used locally, without NPWT. This is most likely to be other sorts of wound dressings and use of any dressing type will be permitted. The frequency of dressing changes will continue as per standard practice.

Participants in both groups will be followed up by telephone on a weekly basis until the participant's wound has healed. Following wound healing, a face to face visit will be completed to assess the wound, complete a wound infection assessment, and to collect a photograph of the healed wound. Healing status will then be confirmed by telephone for two further weeks.

Participants in both groups will be asked to complete short questionnaires at 3 months, 6 months and 12 months which include assessment of pain, wound infection, quality of life and resource use.

Previous intervention:

Patients with a potential planned SWHSI (pre-operatively) or a SWHSI occurring at any point following surgery will be screened for potential eligibility by their clinical care team or GP.

Patients who agree to take part will receive one of the two treatments selected at random using a computer system.

Intervention: Negative Pressure Wound Therapy (NPWT).

Participants will be treated with negative pressure wound therapy (NPWT). The patient's wound is filled with a suitable dressing, and a liner may also be used to protect the wound. A vacuum

pump is then attached which applies suction to the wound. A disposable, plastic canister slots into the pump to collect wound exudate and this is removed and replaced either when it becomes full, or at least once a week.

The device is generally used as part of the SWHSI treatment pathway rather than to the point of healing and is administered by both nurses and clinicians.

The use of any CE marked NPWT device, providing pressure of 60-150mmHg, in use within the NHS will be permitted in this trial, given that the principles of any device are similar and there is no evidence to suggest clinical or cost-effectiveness differences between devices. The device will be used in accordance with manufacturer guidance, and the clinical care team, in conjunction with local treatment guidelines, will determine the duration of device use, and whether this includes continuous or intermittent pressure cycles.

Control: Usual Care (no NPWT)

Usual care will be that used locally, without NPWT. This is most likely to be other sorts of wound dressings and use of any dressing type will be permitted. The frequency of dressing changes will continue as per standard practice.

Participants in both groups will be followed up by telephone on a weekly basis until the participant's wound has healed. Following wound healing, three face to face visits will be completed to assess the wound, complete a wound infection assessment, and to collect photographs of the healed wound.

Participants in both groups will be asked to complete short questionnaires at 3 months, 6 months and 12 months which include assessment of pain, wound infection, quality of life and resource use.

Intervention Type

Device

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Negative Pressure Wound Therapy; Usual wound dressings

Primary outcome(s)

Current primary outcome measure as of 18/11/2020:

Time to healing in days from randomisation (defined using the commonly used and clinically certified criteria 'complete epithelial cover in the absence of a scab (eschar)'). Confirmation of wound healing by a health care professional will initially be participant-reported through weekly telephone contact with the research nurse. Participants will be asked to report if their clinician or nurse has indicated that their wound is healed. In the event the participant reports their wound to be healed, but this has not been confirmed by a healthcare professional, the research nurse will contact the clinical care team to obtain this confirmation. Once wound healing has been confirmed by a healthcare professional and treatment has ceased, participants will undergo clinical assessments on three subsequent consecutive weeks. The first healing visit will be completed face to face wherever possible, during which standardised photographs will be taken of the wound for blinded outcome verification. We will also ask participants to take and return a

photograph of the wound themselves. Study-specific instructions will be provided to the participant.

The second and third post healing visits may be completed by telephone if required.

Previous primary outcome measure:

Time to healing in days from randomisation (defined using the commonly used and clinically certified criteria 'complete epithelial cover in the absence of a scab (eschar)'). Confirmation of wound healing by a health care professional will initially be participant-reported through weekly telephone contact with the research nurse. Participants will be asked to report if their clinician or nurse has indicated that their wound is healed. In the event the participant reports their wound to be healed, but this has not been confirmed by a healthcare professional, the research nurse will contact the clinical care team to obtain this confirmation. Once wound healing has been confirmed by a healthcare professional and treatment has ceased, participants will undergo clinical assessments on three subsequent consecutive weeks during which standardised photographs will be taken of the wound for blinded outcome verification.

Key secondary outcome(s)

Current Secondary outcome measures as of 11/03/2024:

1. Clinical events including antibiotic treatment, hospital admission or discharge, treatment status (including reasons for dressing or treatment failure or change), re-operation (including skin grafting and closure surgery*), amputation and death will be assessed using clinic records and patient-reported events on a weekly basis until the point of wound healing.

*The decision for closure surgery will be made blinded to treatment allocation as far as is possible

2. Wound infection assessed using the Bluebelle Wound Healing Questionnaire (WHQ). The questionnaire includes items to assess signs, symptoms and wound care interventions indicative of surgical site infection (SSI) and can be completed by patient self-report or by healthcare professionals. The tool may be used to assess wounds in hospital or after the patient has been discharged. The WHQ will be completed by the participants themselves at baseline, 3 month follow-up assessments, and will also be completed by the patient at the initial healing visit

3. Wound pain, assessed using a visual analogue scale (with anchors 0 'no pain' and 10 'worst imaginable pain'). The scale will be completed by the participants themselves at baseline, 3, 6 and 12 month follow-up assessments

4. Quality of life, measured using the EQ-5D-5L at baseline, 3 months, 6 months and 12 months follow-up assessments

5. Resource use: wound-related NHS consultations, support (e.g. occupational therapy, in-home adaptations) and out-of-pocket costs collected using a patient-reported questionnaire at baseline, 3, 6 and 12 months

Previous secondary outcome measures as of 01/04/2022:

1. Clinical events including antibiotic treatment, hospital admission or discharge, treatment status (including reasons for dressing or treatment failure or change), re-operation (including skin grafting and closure surgery*), amputation and death will be assessed using clinic records and patient-reported events on a weekly basis until the point of wound healing.

* The decision for closure surgery will be made blinded to treatment allocation as far as is possible

2. Wound infection assessed using the Bluebelle Wound Healing Questionnaire (WHQ). The questionnaire includes items to assess signs, symptoms and wound care interventions indicative of surgical site infection (SSI) and can be completed by patient self-report or by healthcare

professionals. The tool may be used to assess wounds in hospital or after the patient has been discharged. The WHQ will be completed by the participants themselves at baseline, 3, 6 and 12 month follow up assessments, and will also be completed by the patient at the initial healing visit

3. Wound pain, assessed using a visual analogue scale (with anchors 0 'no pain' and 10 'worst imaginable pain'). The scale will be completed by the participant themselves at baseline, 3, 6 and 12 month follow up assessments
4. Quality of life, measured using the EQ-5D-5L at baseline, 3 months, 6 months and 12 months follow up assessments
5. Resource use: wound-related NHS consultations, support (e.g. occupational therapy, in home adaptations) and out of pocket costs collected using a patient-reported questionnaire at baseline, 3, 6 and 12 months

Previous secondary outcome measures:

1. Clinical events including antibiotic treatment, hospital admission or discharge, treatment status (including reasons for dressing or treatment failure or change), re-operation (including skin grafting and closure surgery*), amputation and death will be assessed using clinic records and patient-reported events on a weekly basis until the point of wound healing. * The decision for closure surgery will be made blinded to treatment allocation
2. Wound infection assessed using the Bluebelle Wound Healing Questionnaire (WHQ). The questionnaire includes items to assess signs, symptoms and wound care interventions indicative of surgical site infection (SSI) and can be completed by patient self-report or by healthcare professionals. The tool may be used to assess wounds in hospital or after the patient has been discharged. The WHQ will be completed by the participants themselves at baseline, 3, 6 and 12 month follow up assessments, and will also be completed by the patient at the initial healing visit
3. Wound pain, assessed using a visual analogue scale (with anchors 0 'no pain' and 10 'worst imaginable pain'). The scale will be completed by the participant themselves at baseline, 3, 6 and 12 month follow up assessments
4. Quality of life, measured using the EQ-5D-5L at baseline, 3 months, 6 months and 12 months follow up assessments
5. Resource use: wound-related NHS consultations, support (e.g. occupational therapy, in home adaptations) and out of pocket costs collected using a patient-reported questionnaire at baseline, 3, 6 and 12 months

Completion date

31/01/2024

Eligibility

Key inclusion criteria

1. Aged 16 years or over
 2. Has an acute SWHSI (i.e. a wound left open as planned following surgery or a wound initially closed using sutures, clips, or other closure methods and dehisced along the whole or part of its length, and of less than 6 weeks in duration), arising from any surgical specialty and occurring on any part of the body, deemed appropriate to receive either NPWT or wound dressing treatment
 3. Has a SWHSI that is considered ready for NPWT treatment (i.e. contains at least 80% viable tissue or has only a very thin layer of slough requiring no further debridement)
 4. Patient is not deemed to be malnourished, as per NICE guidelines CG 32 (BMI <18.5 kg/m²; unplanned* weight loss >10% in the last 3-6 months; BMI <20kg/m² and unplanned* weight loss >5% in the last 3-6 months) or assessed as at high risk of malnutrition using the Malnutrition Universal Screening Tool (MUST)
- *Patients with weight loss arising either from underlying comorbidity (e.g. ulcerative colitis) or

from the reasons for surgery being completed (e.g. bowel cancer) may be included at the clinician's discretion

5. Willing and able to give informed consent and provide follow-up data

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

16 years

Sex

All

Total final enrolment

686

Key exclusion criteria

Current participant exclusion criteria as of 01/04/2022:

1. Life expectancy of less than 6 months e.g. undergoing end stage palliative care
2. Active systemic infection (including osteomyelitis) at baseline as defined by clinical and/or laboratory assessment. Note: Patients who have an active infection, but are improving following 1 week's duration of antibiotics may be included at the clinician's discretion
3. Inadequate haemostasis or patients who are at risk of bleeding
4. Chronic wounds non-surgical in origin (e.g. pressure ulcers or foot ulcers)*

*Note diabetic foot ulcers which have been incised and drained or debrided as an inpatient in theatre may be included given this constitutes a surgical wound.

5. Current wound has previously been, or is currently being, treated with NPWT
6. Planned delayed primary closure of the wound
7. Contraindication to NPWT including: presence of unclear undermining in the wound cavity; presence of necrotic tissue, malignant tissue or eschar; wounds involving exposed blood vessels and/or organs, anastomotic sites and/or nerves (including the "open abdomen" where the abdominal fascia is open); wounds situated where, in the opinion of the treating clinician, a vacuum seal cannot be obtained; presence of a non-enteric or unexplored fistula; people requiring emergency airway aspiration, pleural mediastinal or chest tube drainage or surgical suction (removed 07/11/2019: people with a sensitivity or allergy to silver)
8. Currently participating in another wound research study, where the primary outcome time point has not yet been reached

Previous participant exclusion criteria:

1. Life expectancy of less than 6 months e.g. undergoing end stage palliative care
2. Active systemic infection (including osteomyelitis) at baseline as defined by clinical and/or laboratory assessment. Note: Patients who have an active infection, but are improving following 1 week's duration of antibiotics may be included at the clinician's discretion
3. Inadequate haemostasis or patients who are at risk of bleeding
4. Chronic wounds non-surgical in origin (e.g. pressure ulcers or foot ulcers)

5. Current wound has previously been, or is currently being, treated with NPWT
6. Planned delayed primary closure of the wound
7. Contraindication to NPWT including: presence of unclear undermining in the wound cavity; presence of necrotic tissue, malignant tissue or eschar; wounds involving exposed blood vessels and/or organs, anastomotic sites and/or nerves (including the "open abdomen" where the abdominal fascia is open); wounds situated where, in the opinion of the treating clinician, a vacuum seal cannot be obtained; presence of a non-enteric or unexplored fistula; people requiring emergency airway aspiration, pleural mediastinal or chest tube drainage or surgical suction (removed 07/11/2019: people with a sensitivity or allergy to silver)
8. Currently participating in another wound research study, where the primary outcome time point has not yet been reached

Date of first enrolment

01/05/2019

Date of final enrolment

13/01/2023

Locations

Countries of recruitment

United Kingdom

England

Scotland

Wales

Study participating centre

Hull University Teaching Hospital NHS Trust

Hull Royal Infirmary

Anlaby Road

Hull

United Kingdom

HU3 2JZ

Study participating centre

University Hospitals of Birmingham NHS Foundation Trust

Queen Elizabeth Medical Centre

Edgbaston

Birmingham

United Kingdom

B15 2TH

Study participating centre

Royal Liverpool and Broadgreen University Hospitals NHS Trust

Royal Liverpool University Hospital

Prescot Street

Liverpool

United Kingdom

L7 8XP

Study participating centre

Doncaster and Bassetlaw Teaching Hospitals NHS Foundation Trust

Doncaster Royal Infirmary

Armthorpe Road

Doncaster

United Kingdom

DN2 5LT

Study participating centre

The Dudley Group NHS Foundation Trust

Russells Hall Hospital

Pensnett Road

Dudley

United Kingdom

DY1 2HQ

Study participating centre

Royal Cornwall Hospitals NHS Trust

Royal Cornwall Hospital

Treliske

Truro

United Kingdom

TR1 3LJ

Study participating centre

Imperial College Healthcare NHS Trust

St Mary's Hospital

Praed Street

London

United Kingdom

W2 1NY

Study participating centre

The Newcastle Upon Tyne Hospitals NHS Foundation Trust
The Freeman Hospital
High Heaton
Newcastle
United Kingdom
NE7 7DN

Study participating centre
Leeds Teaching Hospitals NHS Trust
St James' University Hospital
Beckett Street
Leeds
United Kingdom
LS9 7TF

Study participating centre
Norfolk and Norwich University Hospitals NHS Foundation Trust
Colney Lane
Norwich
United Kingdom
NR3 7UY

Study participating centre
University Hospitals of Birmingham NHS Foundation Trust
Heartlands Hospital
Bordesley Green East
Birmingham
United Kingdom
B9 5SS

Study participating centre
St George's University Hospitals NHS Foundation Trust
Blackshaw Road
Tooting
London
United Kingdom
SW17 0QT

Study participating centre
Mid Yorkshire Hospitals NHS Trust
Pinderfields Hospital

Aberford Road
Wakefield
United Kingdom
WF1 4DG

Study participating centre
Pennine Acute Hospitals NHS Trust
Royal Oldham Hospital
Rochdale Road
Oldham
United Kingdom
OL1 2JH

Study participating centre
Aneurin Bevan University Health Board
Royal Gwent Hospital
Cardiff Road
Newport
United Kingdom
NP20 2UB

Study participating centre
Queen Elizabeth Hospital Kings Lynn NHS Trust
Gayton Road
Kings Lynn
United Kingdom
PE30 4ET

Study participating centre
Northumbria Healthcare NHS Foundation Trust
Tissue Viability Office
Nursery Park
Ashington
United Kingdom
NE63 0HP

Study participating centre
University Hospitals of Derby and Burton
Uttoxeter Rd

Derby
United Kingdom
DE22 3NE

Study participating centre
Sunderland Royal Hospital
South Tyneside and Sunderland NHS Foundation Trust
Kayll Road
Sunderland
United Kingdom
SR4 7TP

Study participating centre
Manchester University NHS Foundation Trust
Manchester Royal Infirmary
Oxford Road
Manchester
United Kingdom
M13 9WL

Study participating centre
Frimley Health NHS Foundation Trust
Frimley Park Hospital
Portsmouth Road
Frimley
Camberley
United Kingdom
GU16 7UJ

Study participating centre
Bradford Teaching Hospitals NHS Foundation Trust
Bradford Royal Infirmary
Duckworth Lane
Bradford
United Kingdom
BD9 6RJ

Study participating centre
NHS Lanarkshire
University Hospital Hairmyres
Eaglesham Road

Glasgow
United Kingdom
G75 5RG

Study participating centre

NHS Lothian

Royal Infirmary of Edinburgh
51 Little France Crescent
Old Dalkeith Road
Edinburgh
United Kingdom
EH16 4SA

Study participating centre

North Bristol NHS Trust

Southmead Hospital
Bristol
United Kingdom
BS10 5NB

Study participating centre

Royal Free London NHS Foundation Trust

Royal Free Hospital
Pond Street
London
United Kingdom
NW3 2QG

Study participating centre

University Hospitals of Leicester NHS Trust

Glenfield Hospital
Groby Road
Leicester
United Kingdom
LE3 9QP

Study participating centre

The Shrewsbury and Telford Hospital NHS Trust

Mytton Oak Road

Shrewsbury
United Kingdom
SY3 8XQ

Study participating centre
Worcestershire Acute Hospitals NHS Trust
Worcestershire Royal Hospital
Charles Hastings Way
Worcester
United Kingdom
WR5 1DD

Sponsor information

Organisation
Hull University Teaching Hospital NHS Trust

ROR
<https://ror.org/01b11x021>

Funder(s)

Funder type
Government

Funder Name
Health Technology Assessment Programme

Alternative Name(s)
NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type
Government organisation

Funding Body Subtype
National government

Location
United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Current IPD sharing plan as of 23/10/2025:

Anonymised datasets generated and analysed during the current study will be stored in the publicly available open research repository OSF. Data is now available via this repository, following completion of analysis and subsequent publication via the following link: https://osf.io/8mehf/overview?view_only=07643d8f208f45f18b214594ca645103

Sharing of this anonymised data is covered by original participant consent for the SWHSI-2 trial which permits sharing of data to support future research via sharing anonymously.

Previous IPD sharing plan:

Anonymised datasets generated and analysed during the current study will be stored in a publicly available open research repository (<https://osf.io/echxv>). Data is anticipated to be available via this repository by end 2024, following completion of analysis and subsequent publication. Sharing of this anonymised data is covered by original participant consent for the SWHSI-2 trial which permits sharing of data to support future research via sharing anonymously.

IPD sharing plan summary

Stored in publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		15/04/2025	23/04/2025	Yes	No
Results article	outcomes and learning from the internal pilot phase and main trial	02/07/2025	03/07/2025	Yes	No
Protocol article		25/10/2021	19/01/2023	Yes	No
Dataset		13/05/2025	21/05/2025	No	No
HRA research summary			28/06/2023	No	No
Other files	Health economics analysis plan version 1.0	14/12/2022	03/02/2023	No	No
Statistical Analysis Plan	version 1.0	14/12/2022	03/02/2023	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes