Early or late drain removal after pancreatic surgery

Submission date	Recruitment status	[X] Prospectively registered
10/04/2018	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
11/05/2018	Ongoing	Results
Last Edited	Condition category	Individual participant data
08/06/2023	Surgery	Record updated in last year

Plain English summary of protocol

Background and study aims

Abdominal drains are sometimes placed during surgery in order to minimize the risk from bowel contents or other dangerous fluids collecting in the abdomen. This practice has been reduced in many branches of surgery during the last decades as the benefit of this treatment is questionable and the risk associated with having drains (plastic tubes) inside the abdomen is inevitable. When it comes to pancreatic surgery the risk from leakage of pancreatic juice into the abdominal cavity is high and the routine use of drains is therefore the standard of care. The drains are usually left in place for a minimum of 3 days after surgery and then removed if no signs of leakage are found. There are results available from earlier research indicating that drains can safely be removed as early as one day after surgery if no leakage is present. The aim of this study is to find out whether early drain removal after pancreatic surgery shortens hospital stay.

Who can participate?

Patients undergoing removal of the pancreatic head and associated tissue (Whipple operation) and with normal concentration of drain amylase (pancreatic enzyme) the day after surgery

What does the study involve?

Participants are randomly allocated to undergo either early drain removal (one day after surgery) or conventional drain removal (at earliest three days after surgery). Their treatment is otherwise according to hospital standards and the follow up (within the study) is 90 days. Length of hospital stay is measured at discharge.

What are the possible benefits and risks of participating?

The possible benefits of participating are that early drain removal may shorten hospital stay and reduce the risk of drain-associated complications. The main risk is that the normal drain amylase on the day after surgery may not completely exclude leakage and therefore longer drain treatment could be of value.

Where is the study run from?

- 1. Linköping University Hospital (Sweden)
- 2. Skånes University Hospital (Sweden)

- 3. Sahlgrenska University Hospital (Sweden)
- 4. Oslo University Hospital (Norway)

When is the study starting and how long is it expected to run for? April 2017 to December 2025

Who is funding the study? Investigator initiated and funded

Who is the main contact? Dr Bergthor Björnsson bergthor.bjornsson@liu.se

Contact information

Type(s)

Scientific

Contact name

Dr Bergthor Björnsson

Contact details

Linköping University Hospital Linköping Sweden 58185 +46 (0)703766890 bergthor.bjornsson@liu.se

Additional identifiers

Protocol serial number

DrainRemovalStudy1.1

Study information

Scientific Title

Post-operative drainage after pancreaticoduodenectomy: a randomized controlled trial of early drain removal vs late drain removal

Study objectives

Early drain removal after pancreatoduodenectomy shortens hospital stay.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Regional Ethical Review Board, Linköping, Sweden, 16/10/2017, ref: Dnr 2017/310-31

Study design

Prospective randomized study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Pancreatic disease treated with pancreatoduodenectomy

Interventions

The randomisation will be by online websolution after the results of drain amylase POD 1 are known and within the inclusion limits of the study. There are two treatment arms:

- 1. Early drain removal (POD 1)
- 2. Conventional drain removal earliest on POD 3 when no pancreatic fistula exist according to the definition of POPF (postoperative pancreatic fistula)

The treatment will otherwise be according to hospital standards and the follow up (within the study) is 90 days.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Hospital stay, measured at discharge

Key secondary outcome(s))

- 1. Overall morbidity rate classified as Clavien-Dindo ≥3a within 90 days
- 2. Mortality rate (in-hospital, 30 and 90 days)
- 3. Overall morbidity rate as classified by Clavien-Dindo within 90 days
- 4. Fistula rate according to ISGPF (grade B and C fistulas) within 90 days
- 5. Intraabdominal abscess rate within 90 days
- 6. Wound infection rate within 90 days
- 7. Delayed gastric emptying according to ISGPF within 90 days
- 8. Postpancreatectomy bleeding according to ISGPF within 90 days
- 9. Need of interventional radiology within 90 days
- 10. Need for reoperation within 90 days
- 11. Readmission rate within 90 days

Completion date

31/12/2025

Eligibility

Key inclusion criteria

Patients planned for pancreatoduodenectomy with high risk for anastomosis leakage

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Low risk for anastomosis leakage

Date of first enrolment

01/09/2018

Date of final enrolment

31/08/2025

Locations

Countries of recruitment

Norway

Sweden

Study participating centre Linköping University Hospital

Linköping Sweden 58185

Study participating centre Skånes University Hospital

Lund Sweden 22242

Study participating centre Sahlgrenska University Hospital

Gothenburg Sweden 41345

Study participating centre Oslo University Hospital Oslo Norway 0372

Sponsor information

Organisation

Region Östergotland

ROR

https://ror.org/0326gsy75

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date. Additional documents (such as study protocol, statistical analysis plan) will not be made available online at this time.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet 11/11/2025 No Yes