A two-part study to evaluate the safety, tolerability, and processing by the body (pharmacokinetics) of fenebrutinib (part A) and to evaluate the effect of fenebrutinib on the heartbeat (QT/QTC interval) (part B) in healthy subjects

Submission date	Recruitment status	Prospectively registered
21/04/2021	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
29/05/2021	Completed	Results
Last Edited	Condition category	Individual participant data
01/08/2022	Other	Record updated in last year

Plain English summary of protocol

Background and study aims

This is a two-part study of an investigational drug called fenebrutinib that will be performed in healthy volunteers. Part A will evaluate the safety and drug levels in the blood of single doses of fenebrutinib. This information will help with the selection of the doses to test in part B. Part B of the study will primarily evaluate whether fenebrutinib has any effect on the electrocardiogram, a measurement of the heart's electrical activity.

Who can participate? Healthy volunteers

What does the study involve?

In part A of the study, a single dose of oral fenebrutinib or placebo will be given to volunteers to evaluate the safety and drug levels in the blood. This information will help with the selection of the doses to test in part B. In part B of the study, on different days volunteers will be given a single dose of fenebrutinib (at two different dose levels), moxifloxacin (an antibiotic), and placebo. The objective of part B is to evaluate whether fenebrutinib has any effect on the electrocardiogram, a measurement of the heart's electrical activity. Part B will also evaluate the safety and blood levels of fenebrutinib. Volunteers will participate in either part A or B, but not both.

What are the possible benefits and risks of participating?

Study volunteers will not receive any medical benefit. They will be compensated for their time. One identified risk of fenebrutinib is potential increased blood level of liver enzymes. Other risks that have not been observed but are theoretically possible include infection, change in

levels of certain cells in the blood, liver injury, effect on vaccinations, bleeding, nausea, vomiting, diarrhea, abnormal heart rhythm, inflammation (swelling or redness) of blood vessels, ability to fight cancer, birth defects, or rash.

Where is the study run from? Genentech (USA)

When is the study starting and how long is it expected to run for? November 2020 to August 2022

Who is funding the study? F. Hoffmann-La Roche Ltd (USA)

Who is the main contact? global-roche-genentech-trials@gene.com, reference Protocol ID GP42654

Contact information

Type(s)

Public

Contact name

Dr Clinical Trials

Contact details

1 DNA Way South San Francisco United States of America 94080 +1 888-662-6728 global-roche-genentech-trials@gene.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

GP42654

Study information

Scientific Title

A two-part, phase 1, randomized, double-blind, single ascending dose study to evaluate the safety, tolerability, and pharmacokinetics of fenebrutinib in healthy subjects (part A) and a randomized, single-dose, placebo- and positive-controlled, crossover phase 1 study to evaluate the effect of fenebrutinib on the QT/QTC interval in healthy subjects (part B)

Study objectives

To evaluate the safety, tolerability and clinical activity of fenebrutinib in healthy subjects

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 10/03/2021, Salus IRB (2111 West Braker Lane, Suite 100, Austin, TX, USA; +1 512-380-1244; salus@salusirb.com), ref: 8441772

Study design

Single center phase I double-blinded two-part interventional study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Safety, tolerability and clinical activity of fenebrutinib in healthy subjects

Interventions

Part A: Single Ascending Dose Study

Participants will enroll in three sequential cohorts (A1, A2, and A3) that will test single ascending doses. In cohort A1, participants will receive a single dose of fenebrutinib or placebo, administered orally. Dosing of subsequent cohorts A2 and A3 will be contingent upon available preliminary fenebrutinib safety, tolerability, and PK data from the preceding dose. Participants in cohorts A2 and A3 will also receive a single dose of oral fenebrutinib or placebo; the dose will be determined based on analysis of the data from the prior cohort.

Duration: 1 day of treatment, 7 days of follow-up.

Part B: Thorough QT Interval Study

The fenebrutinib therapeutic dose and supratherapeutic dose in Part B will be selected based on PK and safety data from Part A. Each participant will receive a single dose of the following treatments:

- 1. Fenebrutinib therapeutic dose
- 2. Fenebrutinib supratherapeutic dose
- 3. Positive control, moxifloxacin 400-mg tablet
- 4. Placebo

Duration: Approximately 24 days of treatment and 7 days of follow-up.

The randomization is done using a randomization list and unblinded pharmacist since there is only one site for this study. In Part A of the study participants are randomized 3:1 to active drug or placebo. In part B, participants are randomized 1:1:1:1 to one of four sequences of receiving the study drugs.

Intervention Type

Drug

Phase

Phase I

Drug/device/biological/vaccine name(s)

fenebrutinib, RO7010939

Primary outcome(s)

Part A:

- 1. Incidence and severity of AEs and incidence of abnormalities measured using clinical laboratory evaluations, 12-lead ECGs, vital signs measurements, and physical examinations throughout the study
- 2. Pharmacokinetics measured using Cmax, tmax, AUC0- ∞ and/or AUC0-t, and t1/2 of fenebrutinib on days 1 and 2

Part B

1. Placebo-adjusted change from predose-averaged baseline QTcF ($\Delta\Delta$ QTcF) of fenebrutinib measured using ECG on days 1 and 2

Key secondary outcome(s))

Part B:

- 1. Incidence and severity of AEs and incidence of abnormalities measured using clinical laboratory evaluations, 12-lead ECGs, vital signs measurements, and physical examinations throughout the study
- 2. Pharmacokinetics measured using Cmax, tmax, AUC0-∞ and/or AUC0-t, and t1/2 of fenebrutinib and moxifloxacin on days 1 and 2
- 3. Absolute and change from baseline in heart rate and PR, RR, QRS, QT, and QTcB intervals; and change from baseline T-wave morphology and U-wave presence measured using ECG on days 1 and 2
- 4. $\Delta\Delta$ QTcF of moxifloxacin measured using ... at 1, 1.5, 2, 3, and 4 hours postdose
- 5. Change from baseline QTcF (Δ QTcF) at the individual tmax for each treatment (Δ QTcF, tmax) measured using ECG

Completion date

21/08/2022

Eligibility

Key inclusion criteria

- 1. Males or females of non-childbearing potential, between 18 and 60 years of age, inclusive;
- 2. Within body mass index (BMI) range 18 to 31 kg/m², inclusive, with a bodyweight >45 kg;
- 3. In good health, determined by no clinically significant findings from medical history, 12-lead ECG, and vital signs;
- 4. Clinical laboratory evaluations (including chemistry panel [fasted at least 10 hours], complete blood count [CBC], coagulation [prothrombin time {PT}, international normalized ratio {INR}, and activated partial thromboplastin time {aPTT}], and urinalysis [UA] with complete microscopic analysis) within the reference range for the test laboratory, unless deemed not clinically significant by the investigator;
- 5. Negative test for selected drugs of abuse and cotinine at Screening (does not include alcohol) and at Check-in (Day -1 for Part A, Period 1 Day -1 for Part B; does include alcohol);
- 6. Negative hepatitis panel (hepatitis B virus core antibody, hepatitis B surface antigen, and hepatitis C virus antibody) and negative human immunodeficiency virus (HIV) antibody screens;
- 7. Females will not be pregnant or breastfeeding, and must be either postmenopausal (at least

12 months without a period [i.e., amenorrhea]; in a woman at least 50 years of age and documented by a serum follicle-stimulating hormone [FSH] level consistent with postmenopausal status [i.e., ≥40 IU/L] in the absence of a reversible medical iatrogenic cause) or surgically sterile (e.g., tubal ligation, bilateral salpingectomy, or hysterectomy) for at least 90 days. For all females, the pregnancy test result must be negative at Screening and Check-in (Day -1 for Part A, Period 1 Day -1 for Part B).

8. Males with partners of childbearing potential will either be sterile (confirmed by documentation in addition to agreeing to using a condom from Check-in [Day -1 for Part A, Period 1 Day -1 for Part B] until 90 days following study completion) or agree to use from Check-in (Day -1 for Part A, Period 1 Day -1 for Part B) until 90 days following study completion, one of the following approved methods of contraception: male condom with spermicide; sterile sexual partner; use by female sexual partner of an intrauterine device with spermicide; a female condom with spermicide; a contraceptive sponge with spermicide; an intravaginal system [e.g., NuvaRing®]; a diaphragm with spermicide; a cervical cap with spermicide; or oral, implantable, transdermal, or injectable contraceptives. Subjects will refrain from sperm donation from Check-in (Day -1 for Part A, Period 1 Day -1 for Part B) until 90 days or 5 half-lives plus 74 days (a spermatogenesis cycle), whichever is longer, following study completion. Male subjects (including men who have had vasectomies) whose partners are currently pregnant should use a barrier method for the duration of the study and for 90 days or 5 half-lives plus 74 days (a spermatogenesis cycle), whichever is longer, after the study completion. This is to ensure that the fetus is not exposed to the drug in the ejaculate;

9. Receive an explanation of the mandatory WGS component of the study; 10. Able to comprehend and willing to sign an Informed Consent Form (ICF); 11. Able to comply with the study restrictions.

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

- 1. Significant history or clinical manifestation of any metabolic, allergic, dermatological, hepatic, renal, hematological, pulmonary, cardiovascular, gastrointestinal (GI), neurological, or psychiatric disorder (as determined by the investigator)
- 2. Evidence of active infection (with the exception of fungal nail infections or oral herpes); history of recurrent bacterial, viral, mycobacterial, or fungal infections (defined as >2 similar episodes requiring anti-microbial treatment within the previous 12 months), with the exception of recurrent oral or genital herpes (herpes simplex virus 1/herpes simplex virus 2) or uncomplicated urinary tract infections in females; or history of infection requiring hospitalization within 8 weeks prior to Screening
- 3. History of significant hypersensitivity, intolerance, or allergy to any drug compound, food, or

other substance, unless approved by the investigator

- 4. Any personal or family history of bleeding disorders and any personal use of drugs known to affect blood clotting within 30 days prior to dosing
- 5. Family history of intracranial bleed (berry aneurysm, hemorrhagic stroke) or recent personal history of head trauma
- 6. History of vasculitis
- 7. Having one or more of the following clinical laboratory evaluations at Screening:
- Estimated glomerular filtration rate (eGFR) <60 mL/min/1.73 m 2 as calculated using the Cockcroft-Gault or Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation (may be repeated if 45 to 59 mL/min/1.73 m 2
- ALT or AST >2 × ULN (may be repeated if 2 to 3 × ULN)
- Total bilirubin >1.5 × ULN (may be repeated if 1.6 to 3 × ULN)
- Abnormalities in hepatic synthetic function tests (e.g., PT, INR, aPTT, albumin) judged by the investigator to be clinically significant
- 8. Sustained (i.e., three consecutive occasions of assessments) systolic blood pressure measurements <85 or >140 mmHg or diastolic blood pressure measurements <45 or >90 mmHg at Screening
- 9. Sustained (i.e., confirmed upon repeat) pulse >100 or <45 beats per minute at Screening
- 10. Personal or family history of congenital long QT syndrome or family history of sudden death
- 11. History or presence of an abnormal ECG (including marked mean QTcF prolongation ≥450 msec) at Screening or Check-in (Day -1 for Part A, Period 1 Day -1 for Part B) that, in the investigator's opinion, is clinically significant
- 12. Short QTc (mean QTcF <300 msec) at Screening or Check-in (Day -1 for Part A, Period 1 Day -1 for Part B)
- 13. Subject had dyspepsia for which he/she had recently taken (within 2 weeks prior to Check-in [Day -1 Part A, Period 1 Day -1 for Part B]) prescription and/or over-the-counter medicinal products for the control of gastric acidity (e.g., proton-pump inhibitors, H2 blockers, antacids)
- 14. History of stomach or intestinal surgery or resection or any GI disorder that would potentially alter absorption and/or excretion of orally administered drugs, except that appendectomy, hernia repair, and/or cholecystectomy will be allowed
- 15. History of alcoholism or drug addiction within 1 year prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B)
- 16. History of tuberculosis (TB) or treatment with TB prophylaxis within 12 months prior to study enrollment
- 17. Use of oral antibiotics within 4 weeks or IV antibiotics within 8 weeks prior to the Screening evaluation
- 18. Any vaccination within 6 weeks prior to dosing
- 19. Participation in any other investigational study drug trial in which receipt of an investigational study drug occurred within 30 days or 5 half-lives, whichever is longer, prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B)
- 20. Use of any moderate or strong CYP3A inhibitor or inducer within 30 days or 5 half-lives, whichever is longer, prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B)
- 21. Use of any prescription medications/products within 14 days or 5 half-lives, whichever is longer, prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B), unless deemed acceptable by the investigator
- 22. Use of acetaminophen within 24 hours prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B)
- 23. Use of any over-the-counter, non-prescription preparations (including vitamins; minerals; and phytotherapeutic-, herbal-, and plant-derived preparations) within 14 days or 5 half-lives, whichever is longer, prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B), unless deemed acceptable by the investigator
- 24. Use of tobacco- or nicotine-containing products (including, but not limited to, cigarettes, e-

cigarettes, pipes, cigars, chewing tobacco, nicotine patches, nicotine lozenges, or nicotine gum) within 6 months prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B)

- 25. Use of grapefruit-containing foods or beverages within 7 days prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B), unless deemed acceptable by the investigator
- 26. Strenuous exercise from 7 days prior to Check-in (Day -1 Part A, Period 1 Day -1 for Part B)
- 27. Use of alcohol- or caffeine-containing foods or beverages within 48 hours prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B), unless deemed acceptable by the investigator
- 28. Poor peripheral venous access
- 29. History of malignancy, except for appropriately treated carcinoma in situ of the cervix or nonmelanoma skin carcinoma with 3-year disease-free follow-up
- 30. Donation or loss of blood in excess of approximately 450 mL from 30 days prior to Screening through study completion, inclusive, or of plasma from 2 weeks prior to Screening through study completion, inclusive
- 31. Receipt of blood products within 2 months prior to Check-in (Day -1 for Part A, Period 1 Day -1 for Part B)
- 32. Any acute or chronic condition that, in the opinion of the investigator, would limit the subject' s ability to complete and/or participate in this clinical study.

Date of first enrolment 27/04/2021

Date of final enrolment 18/07/2022

Locations

Countries of recruitment United States of America

Study participating centre Covance Research Unit Dallas United States of America 75247

Sponsor information

Organisation

Roche (United States)

ROR

https://ror.org/011qkaj49

Funder(s)

Funder type

Industry

Funder Name

Roche

Alternative Name(s)

F. Hoffmann-La Roche Ltd, F. Hoffmann-La Roche & Co, F. Hoffmann-La Roche AG, Roche Holding AG, Roche Holding Ltd, Roche Holding, Roche Holding A.G., Roche Holding, Limited, F. Hoffmann-La Roche & Co., Roche Holdings, Inc.

Funding Body Type

Government organisation

Funding Body Subtype

For-profit companies (industry)

Location

Switzerland

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due to this not being a regulatory requirement.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet 11/11/2025 No Yes