# Health and development effectiveness of integrated home-based interventions in rural Andean communities: a randomised trial

Submission date	Recruitment status No longer recruiting	<ul><li>Prospectively registered</li></ul>		
30/11/2017		☐ Protocol		
Registration date 15/01/2018	Overall study status Completed	Statistical analysis plan		
		[X] Results		
<b>Last Edited</b> 07/06/2022	<b>Condition category</b> Other	[] Individual participant data		
0110012022	Other			

#### Plain English summary of protocol

Background and study aims

Children and women in low- and middle-income countries are frequently exposed to accumulating health and developmental risks often rooted in unhealthy environments. Nearly 3 billion people worldwide use solid fuels, which are a major source of household air pollution (HAP). Exposure to HAP drastically increases the risk of acute respiratory infections and cognitive deficits in children. In addition, a considerable fraction of the world population lacks access to secure drinking water sources (11%) or adequate sanitation (36%). Exposure to unsafe drinking water, sanitation and inadequate personal hygiene (WASH) increases the risk of several negative health outcomes such as diarrhoea, malnutrition or trachoma. In 2011, HAP and WASH accounted for 1.3 million child deaths, more than AIDS, malaria and measles combined. Early child development (ECD) is one of the main social determinants of health – opportunities that are provided to young children are crucial in shaping their lifelong health and development status. However, today 200 million children worldwide are not developing their full cognitive potential. This disparity has lifetime adverse consequences and impacts the wellbeing of future generations. Poor cognitive stimulation has been identified as one of the key risk factors hampering ECD in low-income countries. Poor health and development require integrated approaches to address underlying the risks and structural determinants. Many investigations have demonstrated that low-cost interventions addressing common health risk factors at the same time have a powerful combined effect, leading to significant health and social improvements. Health and ECD interventions have demonstrated to be effective alone, so integrating environmental, health and ECD interventions can create safe, stimulating and responsive homes where children can play and eat without being exposed to factors that threaten their health and development. Furthermore, this approach can reduce the use of resources and provide a more holistic approach towards prevention. Although Peru is now classified as upper-middle income country economic growth it maintains large inequalities between urban and rural areas with poverty and extreme poverty, infant mortality, chronic malnutrition and illiteracy. The home-based interventions proposed in this study can alleviate the negative effects of poverty on child health and development and thereby prevent the widening gap between the rich and the poor. This includes health and non-health benefits such as better nutritional status, empowerment of women or the community, capacity building,

increased security and time savings. The aim of this study is to assess the effectiveness of a package of home-environmental health and ECD interventions in rural Andean communities.

#### Who can participate?

Families with at least one child aged under 20 months, who use solid fuels as their main energy source for cooking/heating, have access to piped water in the yard or the community where the kitchen connections can be made, and do not plan to move within the next 24 months

#### What does the study involve?

Communities in two poor rural provinces of the Peruvian Andes are randomly allocated to one of four groups. Households in the three intervention groups receive either improved biomass cookstoves, kitchen sinks and hygiene education, the Peruvian national program on ECD, or both interventions combined. The control group do not receive any of the interventions during the study and follow-up period, but receive all the interventions (kitchen stove and sink) at the end of study. The researchers monitor the occurrence of acute respiratory infections, diarrhoea and changes in ECD in children under 36 months of age over one year through weekly visits and start and end of study comparisons. In addition, they study changes in exposure to air pollution in households and individuals in a small group of participants to assess cookstove efficiency, and also analyse the presence of E.coli (one of the main causes of diarrhoea) in drinking water of all the participants at the beginning and end of the study and regularly during the study in a small group of participants.

What are the possible benefits and risks of participating?

Households receive a cookstove, a kitchen sink and a piped water supply into the kitchen, which may improve health and development. The study will provide much needed evidence on health and non-health impacts and factors influencing the effectiveness of the interventions. There are no known risks of participating in this study. All study households receive all of the interventions at the end of the study.

Where is the study run from?

The study takes place in two Peruvian provinces (San Marcos and Cajabamba)

When is the study starting and how long is it expected to run for? November 2014 to December 2017

Who is funding the study?

- 1. UBS Optimus Foundation (Switzerland)
- 2. Grand Challenges Canada (Canada)

Who is the main contact?

- 1. Stella Hartinger
- 2. Daniel Mausezahl

# Contact information

**Type(s)**Scientific

#### Contact name

Mrs Stella Hartinger

#### Contact details

Av. Reynaldo de Vivanco 410, Dept 405B, Chacarilla, Surco Lima Peru 33

#### Type(s)

Scientific

#### Contact name

Mr Daniel Mausezahl

#### Contact details

Socinstrasse 57 Basel Switzerland 4051

# Additional identifiers

#### Protocol serial number

1.0

# Study information

#### Scientific Title

A 2x2 factorial 12-month randomised trial to evaluate the effectiveness of two integrated home-based interventions in environmental health and early child development for improving health and child development in children under 36 months living in rural Andean communities (IHIP-2)

#### Acronym

IHIP-2

#### Study objectives

Hypothesis 1: Improved biomass cookstoves and improvements in access to drinking-water and hygiene education lead to reductions of the morbidity of the most important diseases in children.

Hypothesis 2: Home-based early child stimulation leads to measurable improvements in major child development areas.

Hypothesis 3: Integration of environmental health and early child education interventions leads to synergistic effects.

## Ethics approval required

Old ethics approval format

### Ethics approval(s)

The Cayeteno Heredia University ethics review board, 02/11/2014

# Study design

Community-level controlled 2x2 factorial cluster randomised non-blinded trial

#### Primary study design

Interventional

#### Study type(s)

Prevention

#### Health condition(s) or problem(s) studied

Diarrhoea, acute respiratory infections and early child development

#### **Interventions**

The interventions will be implemented as a randomised controlled trial with a 2x2 full factorial design. This design applies two intervention packages individually and in combination:

- 1. An environmental health intervention comprised of improved biomass cookstoves, kitchen sinks and hygiene education (IHIP)
- 2. An early child development program applied by the Peruvian state (ECD)

This design leads to four potential experimental conditions with a 1:1:1:1 allocation ratio. Communities in two poor rural provinces of the Peruvian Andes will be randomly allocated to the three intervention and one control group:

- 1. IHIP + ECD
- 2. IHIP
- 3. ECD
- 4. Control: the control group did not receive any of the interventions during the trial and follow-up period. However, participants in the control group will receive all the interventions (kitchen stove and sink) at the end of study.

The four intervention arms are assessed simultaneously during 12 months for a primary endpoint of changes in ECD and health morbidity in children under 36 months of age.

#### Intervention Type

Mixed

# Primary outcome(s)

- 1. Diarrhoea incidence: cases recorded weekly during 12 months of follow-up with a paper-based questionnaire using the WHO protocol/definition of diarrhoea (at least three loose stools passed within 24 hours)
- 2. Age standardized mean score of psychomotoric assessment: cognitive, language and motor development will be assessed twice, at baseline and end of study in children <3 years of age. The assessment will be carried out by trained local fieldworkers and trained psychologists using nationally and internationally standardised validated and comparable tools. ECD status is measured with a Peruvian validated tool (ESDI) at the baseline and end of study and with an international validate tool (Bayley Scales of Infant Development) at the end of study.

# Key secondary outcome(s))

- 1. Compliance to use the interventions, recorded weekly during 12 months of follow-up
- 2. Acute respiratory infections incidence during 12 months of follow-up: ARI cases are recorded weekly with a paper-based questionnaire using the WHO/IMCI protocol for diagnosis. ARI will be diagnosed in children <3 years of age by trained local fieldworkers using the IMCI protocol: presence of cough and fever and observable signs: difficult breathing (>60 breaths per minute and >50 breaths per minute in children < 1 years and > 1 year respectively), chest in-drawing,

stridor or other danger signs (i.e. vomiting, being lethargic)

- 3. Incidence of severe cases of diarrhoea such as persistent diarrhoea (lasts 14 days or longer) and bloody diarrhoea
- 4. Household and personal exposure to carbon monoxide (CO) and particle matter (PM2.5) during 12 months of follow-up. Household Air Pollution (HAP) assessment is conducted at various time points (4 time points) in a sub-sample of participants (N=40)
- 5. Microbiological contamination (E.coli) in drinking water. This analysis is carried out by a trained biologist for all study participants at the baseline and end of study, and in a sub-sample of participants (N=40) at various time points over 12 months (4 measurements). The method used for the microbiological analysis is the membrane filtration method (delAgua)

#### Completion date

31/12/2017

# **Eligibility**

#### Key inclusion criteria

Families eligible for the trial must comply with all of the following at recruitment:

- 1. Have at least one child <20 months living in the home
- 2. Use solid fuels as main energy source for cooking/heating
- 3. Have access to piped water in the yard or the community where the kitchen connections can be made
- 4. Do not plan to move within the next 24 months

### Participant type(s)

Other

#### Healthy volunteers allowed

No

#### Age group

Mixed

#### Sex

All

#### Total final enrolment

317

#### Key exclusion criteria

- 1. Participate in the Peruvian national programme on early child development (PNCM)
- 2. The child has any congenital or chronic disease
- 3. Refusal to participate

#### Date of first enrolment

02/07/2015

#### Date of final enrolment

24/02/2016

# **Locations**

#### Countries of recruitment

Peru

Study participating centre
UPCH-SwissTPH San Marcos Research Station

Jr. José Galvez 757 San Marcos Peru 51

# Sponsor information

## Organisation

Universidad Peruana Cayetano Heredia

#### Organisation

Swiss Tropical and Public Health Institute

# Funder(s)

# Funder type

Charity

#### Funder Name

**UBS Optimus Foundation** 

## Alternative Name(s)

# **Funding Body Type**

Private sector organisation

#### **Funding Body Subtype**

Trusts, charities, foundations (both public and private)

#### Location

Switzerland

#### Funder Name

**Grand Challenges Canada** 

#### Alternative Name(s)

Grands Défis Canada, gchallenges, Grand Challenges Canada / Grands Défis Canada, grandchallengescanada, GCC

## **Funding Body Type**

Government organisation

# **Funding Body Subtype**

National government

#### Location

Canada

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Daniel Mausezahl and Stella Hartinger.

## IPD sharing plan summary

Available on request

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	02/04/2020	06/04/2020	Yes	No
Results article		06/06/2022			No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes