

Does wearing a sling after keyhole tendon repair surgery in the shoulder give a better outcome than not wearing a sling?

Submission date 04/04/2019	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 17/04/2019	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 10/09/2021	Condition category Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

A rotator cuff tear is a damaged tendon around the shoulder. It can cause pain and weakness when moving the shoulder. It is commonly treated using keyhole surgery to reattach the torn tendon to the bone. After surgery, people go through a rehabilitation programme with a physiotherapist that lasts several months. This may involve wearing a sling after surgery. The aim of this study is to see whether a sling is beneficial or detrimental to your recovery after having your torn rotator cuff tendon repaired. Some surgeons prefer to rest the shoulder in a sling for several weeks after surgery to give the repaired tendon some time to heal. This may lead to stiffness due to lack of movement, and weakness of the other muscles around your shoulder, both of which require additional treatment from a physiotherapist. Modern tendon repair techniques are now stronger than ever and a period of rest may no longer be necessary. It is also known that the quality of healing may be improved by early movement.

Who can participate?

Patients undergoing a keyhole repair of a torn tendon in the shoulder

What does the study involve?

Participants' shoulder movements are measured and they complete some questionnaires. They also have a scan 6 months after surgery to check whether the repaired tendon has healed. After surgery, if the surgeon is satisfied that the repair is sound, participants are given an individualised exercise programme by the physiotherapist. This is selected at random from one of two options. One involves wearing a sling for 1-2 days (maximum 1 week), and the other involves using a sling for 4 weeks after surgery. Participants have regular visits to the physiotherapist to monitor their progress over the following 12 months using standard outcome scores, measure their range of movement, and also measure the electrical activity in their shoulder muscles. A scan of their shoulder is also performed after 6 months to determine whether the tendon has healed.

What are the possible benefits and risks of participating?

It is not yet known whether patients benefit more from shorter- or longer-term sling use after

rotator cuff repair. Participation will help to determine the answer to this and if participants are allocated to the treatment that is found to be more successful, then they may derive a benefit.

Where is the study run from?

Manchester University NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for?

January 2017 to August 2020

Who is funding the study?

British Elbow and Shoulder Society (UK)

Who is the main contact?

Chris Peach

chris.peach@mft.nhs.uk

Contact information

Type(s)

Scientific

Contact name

Mr Chris Peach

ORCID ID

<http://orcid.org/0000-0002-2516-4760>

Contact details

Department of Shoulder and Elbow Surgery

Manchester University Foundation Trust

Manchester

United Kingdom

M23 9LT

+44 (0)161 291 6150

chris.peach@mft.nhs.uk

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

NCT03913611

Secondary identifying numbers

4.1

Study information

Scientific Title

Traditional versus accelerated rehabilitation programmes following double-row rotator cuff repair: a randomised trial to compare patient outcomes and structural failure rates

Acronym

S-START

Study objectives

There is a difference in Oxford Shoulder Score 3 months after rotator cuff repair in patients who have an accelerated rehabilitation programme versus traditional rehabilitation.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 01/04/2019, West of Scotland REC 5 (West of Scotland Research Ethics Service, West Glasgow Ambulatory Care Hospital, Dalnair Street, Glasgow, G3 8SW; Tel: +44 (0)141 232-1804; Email: WoSREC5@ggc.scot.nhs.uk), ref: 19/WS/0008

Study design

Randomised clinical trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet.

Health condition(s) or problem(s) studied

Rotator cuff tears

Interventions

The study will involve patients who are having keyhole surgery to repair their torn rotator cuff tendons. If the surgeon achieves a secure repair, patients will be randomly allocated by sealed envelope to traditional rehabilitation or accelerated rehabilitation.

Traditional rehabilitation will require use of a sling for 4-6 weeks, and those in the accelerated group will be asked to wear it for comfort, for up to 1 week. Over the following 12 months, the researchers shall assess patients' outcomes using standard outcome scores (e.g. Oxford Shoulder Score), measure their range of movement, and also measure the electrical activity in

their shoulder muscles. They shall also perform a scan of their shoulder after 6 months to determine whether the tendon has healed to see whether accelerated rehabilitation affects healing.

Intervention Type

Device

Phase

Not Applicable

Primary outcome measure

Primary functional outcome measured using Oxford Shoulder Score (OSS) at 3 months post op (at patient consultation)

Secondary outcome measures

1. Pain and function measured using Shoulder Pain and Disability Index (SPADI) at 6 weeks, 3 months, 6 months and 12 months
2. Quality of life measured using EQ5D-5L at 6 weeks, 3 months, 6 months and 12 months
3. Pain measured using VAS at 6 weeks, 3 months, 6 months and 12 months
4. Range of movement measured by clinical assessment with long arm goniometer at 6 weeks, 3 months, 6 months and 12 months at patient consultation
5. Integrity of the repair assessed using MRI scan at 6 months post op

Overall study start date

08/01/2017

Completion date

01/08/2020

Eligibility**Key inclusion criteria**

All patients awaiting arthroscopic rotator cuff tear surgery

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

120

Key exclusion criteria

1. Non-English speaker
2. Concomitant musculoskeletal disease in ipsilateral arm
3. Unfit for surgical intervention

4. Subscapularis tear
5. Massive cuff tear
6. Intra-operatively, if the tear is found not to be repairable
7. If only a partial repair was possible
8. If the repair was under tension

Date of first enrolment

01/10/2019

Date of final enrolment

01/10/2021

Locations

Countries of recruitment

England

United Kingdom

Study participating centre

Manchester University NHS Foundation Trust

Southmoor Road

Manchester

United Kingdom

M23 9LT

Sponsor information

Organisation

Manchester University NHS Foundation Trust

Sponsor details

Southmoor Road

Manchester

England

United Kingdom

M23 9LT

+44 (0)161 291 6150

chris.peach@mft.nhs.uk

Sponsor type

Hospital/treatment centre

Website

mft.nhs.uk

ROR

<https://ror.org/00he80998>

Funder(s)

Funder type

Other

Funder Name

British Elbow and Shoulder Society; Grant Codes: Ltr014PPG/PO. No. CPRD063001

Funder Name

NIHR Central Commissioning Facility (CCF)

Results and Publications

Publication and dissemination plan

1. Results to be presented at national and international scientific meetings
2. Results will be disseminated to trial participants in plain English form
3. Results will be published in peer-reviewed scientific journals
4. The protocol will be available on the study website

Intention to publish date

01/04/2021

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No