A comparison of the effects of Lisinopril and Bendrofluazide on cardiac baroreceptor sensitivity and large artery function following recent ischaemic cerebral stroke

Submission date	Recruitment status	Prospectively registered
30/09/2004	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
30/09/2004	Completed	Results
Last Edited	Condition category	[] Individual participant data
02/05/2018	Circulatory System	[] Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr David Eveson

Contact details

University Hospitals of Leicester c/o Research and Development Office Leicester General Hospital NHS Trust Leicester United Kingdom LE1 4PW +44 (0)116 258 4109 david.eveson@uhl-tr.nhs.uk

Additional identifiers

Protocol serial number N0123138457

Study information

Scientific Title

A comparison of the effects of Lisinopril and Bendrofluazide on cardiac baroreceptor sensitivity and large artery function following recent ischaemic cerebral stroke

Study objectives

The aim is to investigate the effects on the cardiovascular system of two, commonly-used, blood pressure-lowering treatments following a recent stroke. The objective is to establish the effects of the thiazide diuretic, Bendrofluazide, and the angiotensin Lisinopril

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Stroke

Interventions

Clinical Study - Human Physiology -Inpatient Admission| - Multicentre Study: National|NHS Patients|Study of Non-Therapeutic Procedure

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Bendrofluazide, lisinopril

Primary outcome(s)

Cardiac baroreceptor sensitivity

Key secondary outcome(s))

Not provided at time of registration

Completion date

05/05/2004

Eligibility

Key inclusion criteria

- 1. The onset of first ischaemic stroke between 2 and 8 weeks prior to randomisation.
- 2. A mean brachial blood pressure (BP) level measured on 2 different occasions a week apart between 140/90 mmHg and 180/110 mmHg.
- 3. The presence of sinus rhythm.
- 4. Age >18 years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Not Specified

Key exclusion criteria

Does not match inclusion criteria

Date of first enrolment

27/05/2003

Date of final enrolment

05/05/2004

Locations

Countries of recruitment

United Kingdom

England

Study participating centre University Hospitals of Leicester

Leicester United Kingdom LE1 4PW

Sponsor information

Organisation

Department of Health

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

University Hospitals of Leicester NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration