

Pre-treatment drops or spray for managing earwax

Submission date 28/10/2024	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 23/12/2024	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 19/12/2025	Condition category Ear, Nose and Throat	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Earwax is a naturally occurring substance that forms part of the ear's self-cleaning mechanism. Normally it causes no problems however earwax can build up in the ear canal and become dry and impacted. This build-up could be due to overproduction of earwax, use of cotton buds or reduced self-cleaning mechanism due to age. Excessive earwax causes problematic symptoms such as hearing loss, pain and dizziness and can have a significant negative impact on quality of life. It is thought to affect 2.3 million people per year in the UK.

Earwax that is causing symptoms should be removed as recommended in healthcare guidelines. Earwax can be removed mechanically but research has shown that wax should be softened first to ease its removal and reduce the chances of discomfort. This is done by the use of wax softeners. Many different wax softeners are available but the most commonly used is olive oil. The Audiology service at BCUHB offers a wax removal service ran by Audiology practitioners for patients in North Wales in Primary Care locations. Patients can self-refer or be referred by a GP or other healthcare professional for wax removal. Currently patients are advised to use either olive oil drops or olive oil spray for 7 days prior to their wax removal appointment.

Many previous research studies have previously compared the use of different wax softeners but none have compared different delivery methods (drops or spray). We aim to do this by randomly allocating GP practices that offer the wax removal pathway to either using olive oil spray or olive oil drops prior to their wax removal appointment and comparing the outcomes between the two groups. The outcomes we will investigate are the success of wax removal by microsuction, rate of adverse events and improvement in self-reported symptoms following wax removal.

Who can participate?

Adult patients aged 18 years old or over attending the BCUHB Audiology Primary Care wax removal service delivered in GP Practices across North Wales.

What does the study involve?

Each GP practice delivering the BCUHB Audiology Primary Care wax removal service will be randomly assigned to instruct their patients to use either olive oil drops or olive oil spray for 7 days before attending their wax removal appointment. As per current service delivery patients will be responsible for purchasing/sourcing and administering the olive oil solution themselves.

Patients will then attend their wax removal appointment with the Audiology Practitioner. At the appointment patients will be asked about the symptoms they are experiencing in relation to the wax in their ears. The Audiology Practitioner will then attempt to remove the earwax with microsuction. Following microsuction the Audiology Practitioner will ask the patient to:

1. Rate the improvement in the symptoms they reported at the start of the appointment
2. Record the amount of wax remaining in the patient's ear/s
3. Arrange another appointment for wax removal if it hasn't been fully successful.

The study will be looking at comparing the above outcomes between the two groups (olive oil drops and olive oil spray) to see whether there are any differences.

All data recorded as part of the study is currently being recorded as part of service evaluation and no patient identifiable information will be collected.

What are the possible benefits and risks of participating?

Taking part in the study may not directly benefit the patient but the research will hopefully help us understand whether the method of delivering olive oil to the ear has any difference on the outcome of wax removal. There are no known associated risks to the patient and also participating in the study does not affect the patient's normal clinical care in any way.

Where is the study run from?

Betsi Cadwaladr University Health Board (UK)

When is the study starting and how long is it expected to run for?

January 2022 to December 2025

Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Linor Llwyd Jones, linor.ll.jones@wales.nhs.uk

Contact information

Type(s)

Public, Scientific, Principal investigator

Contact name

Mrs Linor Jones

Contact details

Audiology Department

Glan Clwyd Hospital

Rhyl

United Kingdom

LL18 5UJ

+44 (0)3000843864

Linor.Ll.Jones@wales.nhs.uk

Additional identifiers

Integrated Research Application System (IRAS)

Study information

Scientific Title

A comparison of the effectiveness of pre-treatment olive oil used as drops versus spray prior to earwax removal by microsuction

Study objectives

Olive oil spray is more effective than olive oil drops when used as a pre-treatment wax softener prior to wax removal by microsuction.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 15/07/2024, London - Riverside Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 (0)20 7104 8150; riverside.rec@hra.nhs.uk), ref: 24/LO/0420

Study design

Single-centre two-arm parallel-group interventional cluster randomized control trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Pre-treatment of wax in patients accessing a wax removal service

Interventions

The research study is a two-arm cluster randomised control trial. The clusters refer to GP practices and the two arms refer to the pre-treatment of olive oil drops and olive oil spray. The pre-treatment of choice will be randomised to the GP practices prior to the start of the study. The randomisation will involve assigning to either olive oil spray or olive oil drops to the practice in a prior allocation.

The research is a between-subjects design with one factor (olive oil delivery method) and two levels (olive oil drops and olive oil spray). As per normal clinical care patients will source and administer the olive oil themselves. Patients will be recommended to apply three drops or three sprays of olive oil every day for 7 days as per current BCUHB Audiology Primary Care wax removal service guidelines as stated by the Welsh Government 'Ear Wax Management Primary and Community Care Pathway'.

Randomisation will happen at the GP surgery (cluster) level with a 1:1 allocation ratio to olive oil drops or olive oil spray using stratified block randomisation based on prognostic value. Stratifying based on site size (small, large) using block sizes 2 and 4. Randomisation will be done using randomisation software.

Intervention Type

Device

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Olive oil drops, olive oil spray

Primary outcome(s)

The success and effectiveness of wax removal will be measured by comparing the proportion of patients where wax removal is successful at the first attempt, not needing a further visit, with those who need another visit for a re-attempt at removal. This is determined by the Audiology Practitioner immediately after microsuction.

Key secondary outcome(s)

1. Self-reported needs and symptoms assessed immediately following wax removal. The wax removal pathway is a needs-based pathway and each patient being seen for wax removal is asked about their main need/symptom for accessing the service. These needs/symptoms are documented by the Audiology Practitioner. Following the wax removal, the Audiology Practitioner asks the patient to rate how much improvement there is in their reported need /symptom on a five-point scale (much better, slightly better, about the same, slightly worse, much worse).
2. Post-microsuction visualisation ratings for the two groups as rated on an analogue scale immediately after microsuction is complete, assessed using the ratings below:
0 No or very minimal earwax, complete visualisation of tympanic membrane
1 Minor amount of earwax with majority of tympanic membrane visible
2 Moderate amount of earwax with only part of the tympanic membrane visible
3 Fully occluding wax, no visualisation of tympanic membrane
3. Adverse events of microsuction recorded as per normal clinical practice, any that occur in clinic during or immediately after microsuction and any reported by patients up to 1 week after microsuction.

Completion date

31/12/2025

Eligibility**Key inclusion criteria**

1. Adults over 18 years of age
2. Able to give informed consent for microsuction
3. Presence of wax in one or both ears giving rise to symptoms or need that warrant removal

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

18 years

Upper age limit

99 years

Sex

All

Total final enrolment

0

Key exclusion criteria

1. Under 18 years of age
2. Active ear infection
3. Active dermatitis
4. Perforated or non-intact tympanic membrane
5. Foreign body present in ear canal
6. Any contraindication for microsuction or use of wax softeners – previous ear surgery including mastoid obliteration; grommets in situ

Date of first enrolment

01/11/2024

Date of final enrolment

01/06/2025

Locations

Countries of recruitment

United Kingdom

Wales

Study participating centre

Betsi Cadwaladr University Lhb

Executive Offices, Ysbyty Gwynedd

Penrhosgarnedd

Bangor

Wales

LL57 2PW

Sponsor information

Organisation

Betsi Cadwaladr University Health Board

ROR

<https://ror.org/03awsb125>

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article		18/12/2025	19/12/2025	Yes	No
Protocol (preprint)		14/07/2025	18/12/2025	No	No