Community-based rehabilitation for the elderly following hip fracture

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered	
20/11/2018		[X] Protocol	
Registration date	Overall study status	Statistical analysis plan	
23/11/2018	Completed	[X] Results	
Last Edited	Condition category	Individual participant data	
06/06/2025	Injury, Occupational Diseases, Poisoning		

Plain English summary of protocol

Background and study aims

Hip fracture is a common, major health problem in old age, especially for people with other health problems or who are frail. Some patients who suffer this type of fracture need surgery to repair it. They take a long time to recover, and others may not recover fully. Once patients are discharged, the routine care they receive can vary depending on local NHS policy. Some may not find it as easy to live independently afterwards.

An enhanced rehabilitation package has been designed for patients who are recovering from this surgery, which is delivered in addition to standard NHS care. FEMuR III will compare the enhanced package with standard NHS care to see if it can improve recovery for patients. The study will collect information that will hopefully show the best way to treat patients who are recovering from hip repair surgery after a fracture.

The enhanced rehabilitation package is made specifically for each patient and it is thought that this should improve their recovery. It is considered that the package should work better if it includes physiotherapy (to help patients recover movement), occupational therapy (to help patients with activities associated with daily living) and also provides tools to help build confidence and mood.

Who can participate?

People aged 60 years or older who have recently had a hip fracture repaired by replacement arthroplasty or hip fixation

What does the study involve?

This study will compare the enhanced rehabilitation package to standard NHS care and participants will be randomly allocated to receive one of these care packages. The enhanced rehabilitation package mixes extra therapy with self-help tools, which aim to help patients improve aspects of their own recovery, for example, building confidence in trying exercises by themselves. The enhanced rehabilitation package that has been designed involves additional rehabilitation at follow up visits. Participants will be given a workbook and a goal-setting diary to complete during the first few months of recovery. These follow up visits and the diary will help to collect information about their recovery.

Participants will have their progress followed-up in the 12 months after surgery and information will also be collected during that time so that the study can see how participants are.

What are the benefits and risks of taking part?

It is not known whether standard NHS care or the enhanced rehabilitation package is best, but it is anticipated that both will aid patients recovery following surgery. The aim is that the results from the study will help doctors, therapists, patients and their carers in the future when making decisions about treatment.

There are no foreseeable significant risks involved in taking part in FEMuR III. All of the physical exercises suggested are used in normal rehabilitation after hip fracture and will be supervised by trained healthcare professionals to minimise any risk. The enhanced rehabilitation package will take up more of participants time due to additional therapy sessions and having to complete the diary.

Where is the study run from?

The study is run from the Clinical Trials Research Centre at the University of Liverpool (UK) and will take place at 12 hospitals across the UK

When is the study starting and how long is it expected to run for? August 2018 to January 2022

Who is funding the study? National Institute for Health Research (NIHR) (UK)

Who is the main contact? Claire Soady femur3@liverpool.ac.uk

Contact information

Type(s)

Public

Contact name

Ms Claire Soady

Contact details

Liverpool Clinical Trials Centre
2nd Floor
Institute in the Park
Alder Hey Children's NHS Foundation Trust
Eaton Road
Liverpool
United Kingdom
L12 2AP
+44 (0)151 794 9765
femur3@liverpool.ac.uk

Additional identifiers

Integrated Research Application System (IRAS)

246828

Protocol serial number

38492, IRAS 246828

Study information

Scientific Title

Fracture in the Elderly Multidisciplinary Rehabilitation - Phase III (FEMuR III): a definitive randomised controlled trial and economic evaluation of a community-based rehabilitation package following hip fracture

Acronym

FEMuR III

Study objectives

The primary objective is to determine the effectiveness of an enhanced rehabilitation programme following surgical repair of proximal femoral fracture in older people compared with usual care, in terms of the performance of activities of daily living at the 52 week follow-up.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 31/10/2018, North East – Tyne & Wear South Research Ethics Committee (NHSBT Newcastle Blood Donor Centre, Holland Drive, Newcastle upon Tyne, NE2 4NQ, United Kingdom; +44 (0)207 1048084; tyneandwearsouth.rec@hra.nhs.uk), ref: 18/NE/0300

Study design

Randomised; Interventional; Design type: Treatment, Education or Self-Management, Psychological & Behavioural, Complex Intervention, Physical, Rehabilitation

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Hip fracture

Interventions

Participants will be randomly allocated to the enhanced rehabilitation arm or the usual care arm. Randomisation is performed using a secure 24 hour web based randomisation program. Participants will be asked to complete a range of questionnaires at the baseline.

The intervention will be delivered by community teams either at the participants home or in a community hospital up to 4 months post start of intervention. Participants allocated to the usual care arm will receive usual care as per standard local NHS practice. Participants in the enhanced rehabilitation arm will receive standard rehabilitation care as per local NHS practice, plus up to 6 extra rehabilitation sessions.

All participants will be asked to record hospital admissions, concomitant medications and other visits to healthcare professionals. Participants in the enhanced rehabilitation arm will be

provided with a workbook and a goal setting diary. The objectives of the workbook and diary are to:

- 1. Give patients better understanding of what has happened physically to them and broadly what to expect during their recovery
- 2. Provide information and contact details on rehabilitation services that are available to them as they progress in their rehabilitation (e.g. Intermediate care teams, social services enablement teams, outpatient physiotherapy, falls prevention groups, national exercise referral services). The information will enable patients to ask the therapist they are working with or their GP about available services and what benefit they might offer, and at what stages they would be most beneficial, and to contact the services themselves for more information
- 3. Enable them to work collaboratively with their therapist to set goals and monitor progress of their rehabilitation in order to improve the quality and the quantity of the physical and activities of daily living exercises they are performing
- 4. Improve patients' self-efficacy:
- 4.1. Encourage the patient to set goals they want to achieve and to discuss them with their therapist
- 4.2. Monitor the progress towards/attainment of these goals through keeping a diary of progress. This will provide feedback in the form of both self-reflection and reflection with the therapist. Feedback is recognised as an important component for improving self-efficacy 5. Improve communication between hospital and community services, and between the patient and all the different professionals and services they come into contact with during their rehabilitation.
- 6. Reduce patients fear of falling by improving self-efficacy for avoiding falls/ exercising, and providing information about local falls prevention services
- 7. Signpost patients to local follow-on community programmes such as exercise referral and falls prevention services with contact details.

The RSPO will visit each participant at the 4 month point and participants will be asked to complete the same questionnaires as at the baseline.

After the 4 month follow up visit and at a convenient date at time for trial participants, 30 participants from the usual care arm and 30 participants from the enhanced rehabilitation arm (together with up to 30 of their carers for both arms) will be contacted by telephone by a qualitative researcher to ask patients and carers questions about their experiences and views about their involvement in the trial, these interviews will be recorded so that they can be analysed following the call, consent for this will be obtained from each participant and their carer.

The RSPO will visit each participant at the 12 month point and participants will be asked to complete the same questionnaires as at the baseline.

The 12 month visit will conclude trial participation in both arms for participants and their carers. Throughout the trial safety monitoring will take place, adverse events will be reported routinely and serious adverse events will be reported to the Clinical Trials Research Centre (CTRC).

Intervention Type

Behavioural

Primary outcome(s)

Difference in performance of activities of daily living, assessed using the Nottingham Extended Activities of Daily Living (NEADL) scale at the baseline, after 4 months and after 12 months

Key secondary outcome(s))

Current secondary outcome measures as of 21/03/2025:

The following will be compared between the enhanced rehabilitation programme and usual care:

- 1. Cost-effectiveness of the enhanced rehabilitation programme compared with usual care, assessed using a cost-utility analysis from a health service and personal social care perspective using the following:
- 1.1. EuroQol EQ-5D-3L, completed at the baseline and after 17 and 52 weeks of follow-up
- 1.2. Bespoke Client Service Receipt Inventory (CSRI)
- 2. Effectiveness in terms of performance of activities of daily living after 17 weeks, assessed using the NEADL scale at the baseline and after 17 weeks follow-up
- 3. Effectiveness in terms of anxiety and depression, assessed using the Hospital Anxiety and Depression Scale (HADS) at the baseline and after 17 and 52 weeks of follow-up
- 4. Changes in self-efficacy, hip pain, cognitive function measured using the Abbreviated Mental Test Score (AMTS), fear of falling and physical function as potential mediators for improving activities of daily living:
- 4.1. Self-efficacy, assessed using the International Self-Efficacy Scale at the baseline and after 17 and 52 weeks of follow-up
- 4.2. Hip pain, assessed using a Visual Analogue Scale (VAS) at the baseline and after 17 and 52 weeks of follow-up
- 4.3. Fear of falling, assessed using a Visual Analogue Scale (VAS) at the baseline and after 17 and 52 weeks of follow-up
- 4.4. Physical function, assessed using grip strength with a hand dynamometer and the Short Physical Performance Battery (SPPB) at the baseline and after 17 and 52 weeks of follow-up 5. Change in caregiver strain, anxiety and depression in carers, assessed using the following at
- the baseline and after 17 and 52 weeks of follow-up:
- 5.1. Carer Strain Index (CSI)
- 5.2. Hospital Anxiety and Depression Scale (HADS)
- 6. Mechanisms and processes that explain the implementation and impacts of the enhanced rehabilitation programme, assessed using a process evaluation including:
- 6.1. Qualitative interviews with a purposive sample of participants in each trial arm after the 17-week assessment
- 6.2. Qualitative interviews with the therapists delivering the enhanced rehabilitation programme after the 17-week assessment
- 6.3. Routinely collected data that therapists complete on their information management systems

Previous secondary outcome measures:

The following will be compared between the enhanced rehabilitation programme and usual care:

- 1. Cost-effectiveness of the enhanced rehabilitation programme compared with usual care, assessed using a cost-utility analysis from a health service and personal social care perspective using the following:
- 1.1. EuroQol EQ-5D-3L, completed at the baseline and after 17 and 52 weeks of follow-up
- 1.2. Bespoke Client Service Receipt Inventory (CSRI)
- 2. Effectiveness in terms of performance of activities of daily living after 17 weeks, assessed using the NEADL scale at the baseline and after 17 weeks follow-up
- 3. Effectiveness in terms of anxiety and depression, assessed using the Hospital Anxiety and Depression Scale (HADS) at the baseline and after 17 and 52 weeks of follow-up
- 4. Changes in self-efficacy, hip pain, cognitive function, fear of falling and physical function as potential mediators for improving activities of daily living:
- 4.1. Self-efficacy, assessed using the International Self-Efficacy Scale at the baseline and after 17 and 52 weeks of follow-up
- 4.2. Hip pain, assessed using a Visual Analogue Scale (VAS) at the baseline and after 17 and 52 weeks of follow-up
- 4.3. Fear of falling, assessed using a Visual Analogue Scale (VAS) at the baseline and after 17 and 52 weeks of follow-up
- 4.4. Physical function, assessed using the Short Physical Performance Battery (SPPB) at the

baseline and after 17 and 52 weeks of follow-up

- 5. Change in caregiver strain, anxiety and depression in carers, assessed using the following at the baseline and after 17 and 52 weeks of follow-up:
- 5.1. Carer Strain Index (CSI)
- 5.2. Hospital Anxiety and Depression Scale (HADS)
- 6. Mechanisms and processes that explain the implementation and impacts of the enhanced rehabilitation programme, assessed using a process evaluation including:
- 6.1. Qualitative interviews with a purposive sample of participants in each trial arm after the 17-week assessment
- 6.2. Qualitative interviews with the therapists delivering the enhanced rehabilitation programme after the 17-week assessment
- 6.3. Routinely collected data that therapists complete on their information management systems

Completion date

31/01/2022

Eligibility

Key inclusion criteria

- 1. Aged 60 years or older
- 2. Recent proximal hip fracture including intracapsular, extracapsular (peri-trochanteric, intertrochanteric, reverse oblique or sub-trochanteric)
- 3. Surgical repair by replacement arthroplasty or internal fixation
- 4. Living in their own home prior to hip fracture
- 5. Living and receiving rehabilitation from the NHS in the area covered by the trial centres

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Αll

Key exclusion criteria

- 1. Living in residential or nursing homes prior to hip fracture
- 2. Unable to understand English or Welsh
- 3. Lacking mental capacity to give informed consent

Date of first enrolment

01/04/2019

Date of final enrolment

31/05/2020

Locations

Countries of recruitment

United Kingdom

England

Wales

Study participating centre Norfolk Community Health & Care NHS Trust

Ground Floor 137 West Pottergate Earlham Road Norwich United Kingdom NR2 4BX

Study participating centre BETSI Cadwaladr University

LHB Executive Offices Ysbyty Gwynedd Penrhosgarnedd Bangor Gwynedd United Kingdom CF72 8XR

Study participating centre Prince Charles Hospital CWN TAF University Health Board Gurnos Merthyr Tydfil United Kingdom

Study participating centre St Helens and Knowsley Hospital Services NHS Trust

Whiston Hospital Warrington Road Prescot Merseyside Prescot United Kingdom L35 5DR

CF47 9DT

Study participating centre

Wirral University Teaching Hospital NHS Foundation Trust

Arrowe Park Hospital Arrowe Park Road Upton Wirral Merseyside Wirral United Kingdom CH49 5PE

Sponsor information

Organisation

University of Liverpool

ROR

https://ror.org/04xs57h96

Funder(s)

Funder type

Government

Funder Name

NIHR Evaluation, Trials and Studies Co-ordinating Centre (NETSCC); Grant Codes: 16/167/09

Results and Publications

Individual participant data (IPD) sharing plan

Anonymised data collected during this trial will be available to access. Proposals should be directed to Professor Williams, on behalf of the FEMuR III Trial Management Group (Nefyn.Williams@liverpool.ac.uk). Access will be provided to researchers after the proposal has been reviewed and agreed by the trial data sharing committee.

IPD sharing plan summary

Available on request

Study outputs

Output type
Results article

Details

Date created Date added Peer reviewed? Patient-facing?

18/12/2024 19/12/2024 Yes

No

Results article	Effectiveness	12/05/2025	06/06/2025 Yes	No
Protocol article	protocol	16/10/2020	20/10/2020 Yes	No
HRA research summary			28/06/2023 No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025 No	Yes
Study website	Study website	11/11/2025	11/11/2025 No	Yes