# Does the use of a biliblanket prevent the need for phototherapy in newborns over 24 hours of age with a gestation over 35 weeks who have developed physiological jaundice near to the treatment requirement line?

Submission date	<b>Recruitment status</b> No longer recruiting	<ul><li>Prospectively registered</li></ul>		
05/01/2022		[X] Protocol		
Registration date	Overall study status	Statistical analysis plan		
08/01/2022	Completed	Results		
Last Edited	Condition category	Individual participant data		
07/01/2022	Neonatal Diseases	Record updated in last year		

#### Plain English summary of protocol

Background and study aims

Jaundice is a common and usually harmless condition in newborn babies that causes the skin and the whites of the eyes to turn yellow. In most cases, there is no underlying cause and it is therefore called 'physiological jaundice'. It occurs in well, term infants due to the accumulation of bilirubin (an orange-yellow pigment). Some infants must undergo treatment for jaundice with the use of phototherapy (light therapy) due to the risk of long-term consequences of high levels of bilirubin. Treatment of jaundice requires admission to the neonatal intensive care unit and therefore separating mother and baby. The biliblanket is a phototherapy device that is available for use on the ward. However, there is no evidence as to whether the use of a biliblanket on the ward before the requirement for phototherapy decreases the risk of needing admission for phototherapy. There is also concern that the use of a biliblanket may slow the rise of bilirubin, resulting in prolonged inpatient stays. The aim of this study is to find out whether the use of a biliblanket prevents the need for phototherapy in infants with jaundice.

## Who can participate?

Infants over 35 weeks gestational age and 24 hours of life who have developed neonatal jaundice but have not reached a level requiring treatment with overhead phototherapy.

#### What does the study involve?

Infants are randomly allocated to receive treatment with a biliblanket or to not receive treatment. Both groups will be monitored closely for the need for phototherapy treatment. Both groups will have their bilirubin levels monitored every 12 hours until they require overheard phototherapy treatment or they reach a safe level for their age and gestation.

What are the possible benefits and risks of participating? There are no benefits to individual participants. There is some risk of prolonged length of stay with the use of a biliblanket. No other risks are known.

Where is the study run from?
Coombe Women and Infants University Hospital (Ireland)

When is the study starting and how long is it expected to run for? April 2020 to December 2022

Who is funding the study? Investigator initiated and funded

Who is the main contact? Dr Aoife Branagan branagaa@tcd.ie

## **Contact information**

#### Type(s)

Public

#### Contact name

Dr Aoife Branagan

#### **ORCID ID**

https://orcid.org/0000-0001-5220-5372

#### Contact details

Dublin 8 Dublin Ireland D08W2T0 +353 (0)872218141 branagaa@tcd.ie

#### Type(s)

Scientific

#### Contact name

Dr Aoife Branagan

#### Contact details

Dublin 8
Dublin
Ireland
Dublin 8
+353 (0)872218141
Branagaa@tcd.ie

## Additional identifiers

#### Clinical Trials Information System (CTIS)

Nil known

#### ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Nil known

# Study information

#### Scientific Title

Utilization of a biliblanket to decrease the need for overhead phototherapy and admission to a special care baby unit - a randomised control trial (BiB trial)

#### **Acronym**

BIB

#### **Study objectives**

Neonatal jaundice is a common condition affecting newborn infants, occuring in well, term infants due to the accumulation of bilirubin. Some infants must undergo treatment for jaundice with the use of phototherapy due to the risk of long-term consequences of high levels of bilirubin. Treatment of jaundice requires admission to the neonatal intensive care unit and therefore separating mother and baby. The biliblanket is a phototherapy device which is available for use on the ward. The hypothesis is that treatment with a biliblanket on the postnatal ward, in infants who have a serum bilirubin level approaching the treatment line on a standardised nomogram, will decrease the need for overhead phototherapy.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 12/08/2020, Research Ethics Committee, Coombe Women and Infants University Hospital (Cork St, Dublin 8, Ireland; +353 (0)1-4082000; rec@coombe.ie), ref: 21-2020

## Study design

Single-centre two-armed randomized controlled trial

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Neonatal physiological jaundice

#### **Interventions**

Randomisation will occur on receipt of informed consent. A computer-generated random number list will be used, prepared by an investigator with no clinical involvement in the trial. Sequence generation will be stratified by gestational age – under 37 weeks and over 37 weeks. Once generated the randomisation lists will be sealed in opaque envelopes. Once the patient is consented to enter the trial, the investigator will open the next sequential opaque envelope in the correct strata and provide the allocated interventions. In the case of multiple births, each infant will be individually randomised. Blinding of parents or research staff will not be possible due to the nature of the intervention.

The intervention group will be treated with a biliblanket system on the postnatal ward. The control group will have standard care (no biliblanket). Both groups will have serum bilirubin levels monitored every 12 hours until they require overheard phototherapy treatment or they reach a safe level for their age and gestation.

After receipt of a serum bilirubin level near the treatment line (less than 35 umol/l below treatment line for age taken) when compared with age on an appropriate treatment nomogram, the parents or guardians of the infant will be approached for informed consent by a member of the research team or a neonatal doctor.

#### Intervention Type

Device

#### Phase

Not Applicable

#### Primary outcome(s)

Need for overhead phototherapy (binary [yes/no]): admission to the neonatal unit for phototherapy measured by a chart review at discharge from hospital and time of maximum follow-up (6 weeks of age)

## Key secondary outcome(s))

Length of hospital stay (continuous) measured in days by chart review at time of maximum follow-up (6 weeks of age)

#### Completion date

01/12/2022

## **Eligibility**

## Key inclusion criteria

- 1. Infant greater than 24 hours of age on post natal ward
- 2. Greater than 35 weeks gestation
- 3. Suspected physiological jaundice
- 4. Serum bilirubin less than 35 umol/l under the phototherapy line

## Participant type(s)

**Patient** 

#### Healthy volunteers allowed

No

#### Age group

Neonate

#### Sex

All

#### Key exclusion criteria

- 1. Jaundice under 24 hours
- 2. Infants admitted to the neonatal unit for any reason
- 3. Infant less than 35 weeks gestation
- 4. Known Direct Coombs Test (DCT) positive jaundice at time of recruitment

#### Date of first enrolment

01/09/2020

#### Date of final enrolment

01/08/2022

## Locations

#### Countries of recruitment

Ireland

## Study participating centre Coombe Women and Infants University Hospital

Cork St Dublin 8 Dublin Ireland D08 XW7X

# Sponsor information

#### Organisation

Coombe Women & Infants University Hospital

#### **ROR**

https://ror.org/00bx71042

# Funder(s)

#### Funder type

Other

#### Funder Name

Investigator initiated and funded

## **Results and Publications**

#### Individual participant data (IPD) sharing plan

Data will be available on request from the investigators (Dr A Branagan [branagaa@tcd.ie] or Prof. J Miletin [jmiletin@coombe.ie]). Anonymised data will be made available after analysis in Excel format. The data will be made available on request for the purpose of meta-analysis. As it is fully anonymised it can be shared by email.

#### IPD sharing plan summary

Available on request

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	version 1	17/06/2020	07/01/2022	No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file			07/01/2022	No	No