

Contraception in type 1 diabetes

Submission date 21/06/2017	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 27/06/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 22/07/2025	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Type 1 Diabetes (T1D) is one of the most common chronic diseases of childhood, causing blood sugar levels to become too high. When children with T1D become teenagers or young adults, managing their diabetes is very important due to risky behaviours like unplanned pregnancies. Most young adults are not aware that there can be complications with both the mother and baby due to diabetes, such as birth malformations. Conception should be planned when the mother has normal blood sugar levels. It is important to provide education, counselling and contraception (methods to prevent pregnancy) in young women both with and without T1D. Contraception usually involves hormones (chemical messengers that control the bodies functions), such as birth control or an implantable medical device. Research is needed to examine what the metabolic effects (how food is processed to energy) of hormonal contraception in teenagers with T1D. Previous research has shown that hyperandrogenism (too much male sex hormones) and inflammation (swelling) are frequent problems in women with T1D. Another consideration in this study is to examine the impact of T1D and the use of contraceptives on telomere length. Telomere is the ending portion of the chromosome (part of the genes that carry out genetic information). Telomere length decreases with advancing age and is shortened in cases of type 2 diabetes. Recently, it was shown that the use of COC in adolescents with polycystic ovary syndrome (a syndrome that affects how a womens ovaries work causing irregular periods and other symptoms) decreases telomere length. Whether telomere length is affected by T1D in young women or by the use of contraceptives in these young women, is unknown. The aim of this study is to examine the impact of different types of contraception on young women with T1D when compared to young women without T1D.

Who can participate?

Young women aged 16 to 25 who have T1D and healthy women aged 16-25 years old.

What does the study involve?

Participants are provided counselling about contraception and pregnancy prevention. They are then offer either a contraceptive pill (taken by mouth) or the implantable rods as birth control methods. Participants are followed up for two years and provide blood samples before taking the contraception and at three, six, 12 and 24 months. The women with T1D and the women without are compared to see how the contraception impacts their metabolic control, bone mass, insulin sensitivity and telomere length.

What are the possible benefits and risks of participating?

Participants may benefit from receiving the contraceptive methods and medical control without cost for them. The only risk of the study is the already reported as well-known possible side effects of hormonal contraception.

Where is the study run from?

1. University of Chile (Chile)
2. Instituto Chilena de Medicina Reproductiva (ICMER) (Chile)

When is the study starting and how long is it expected to run for?

May 2017 to April 2021

Who is funding the study?

Fondecyt (Chile)

Who is the main contact?

Professor Ethel Codner

Contact information

Type(s)

Scientific

Contact name

Prof Ethel Codner

ORCID ID

<https://orcid.org/0000-0002-2899-2705>

Contact details

Institute of Maternal and Child Research

Casilla 226-3, Santiago

Santiago

Chile

8360160

Additional identifiers

Protocol serial number

FONDECYT 1170895

Study information

Scientific Title

Metabolic effect of the long-acting reversible contraceptive method (implant) compared to combined oral contraceptive in young women with type 1 diabetes

Study objectives

Adolescents/young women with T1D receiving hormonal combined oral contraception or a long-acting reversible method (implantable progesterone rod) exhibit a detrimental profile of

markers associated with inflammation, insulin resistance, and telomere length compared with healthy non-diabetic young women receiving the same type of contraception. As well, T1D young women will show lower bone mass than healthy young women.

Ethics approval required

Old ethics approval format

Ethics approval(s)

IRB of the Servicio Salud Metropolitano Central (SSMC) of the Health Ministry of Chile, 29/05/2017, ref: N°431/2017

Study design

Interventional non randomised study

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Young women with type 1 diabetes or healthy controls.

Interventions

Healthy participants and participants with T1D (N:40) seeking contraception are recruited. Counseling about contraception and pregnancy prevention is given to both groups of women.

Participants are offered either a combined oral contraceptive pill containing ethinyl estradiol 30 ug/desogestrel 0.15mg, or the implantable rod with etonogestrel, the active metabolite of desogestrel. Both methods are currently available for women of this age group and have been considered as first-line options for young women.

Participants are followed for two years and blood samples are obtained at baseline, three, six, 12, and 24 months.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Implanon (R) contraceptive implant and desogestrel/ethinyl estradiol

Primary outcome(s)

1. Inflammatory markers is measured using C-Reactive Protein Levels at baseline, three, six, and 12 months
2. Metabolic control (HbA1c) is measured using HbA1c with DCA-2000 (Siemens) equipment at month three, six, nine and 12
3. Bone Mass is measured using DXA (Lunar Prodigy GE) at baseline, 12 and 24 months
4. Insulin sensitivity is measured using estimated insulin sensitivity index, which includes the

measurement of adiponectin and was validated for subjects with and without T1D6 at baseline, three, six, and 12 months

5. Telomere length is measured using blood tests at baseline and at 12 months

Key secondary outcome(s)

1. Metabolic control will be assessed at 18 and 24 months in women with type 1 diabetes
2. Bone mass DXA (Lunar Prodigy GE) and insulin sensitivity in all the participants is assessed using DXA (Lunar Prodigy GE) and insulin sensitivity index at 24 months of treatment

Completion date

30/04/2021

Eligibility

Key inclusion criteria

T1DM patients:

1. Clinical diagnosis of T1D is clear (patient received treatment with insulin from the time of diagnosis with clinical evidence of severe insulin deficiency).
2. Age: 16-25 years old.
3. Seeking contraception.
4. HbA1c lower than 13%
5. Has not used hormonal contraception for the last three months

Healthy subjects:

1. Age 16-25 years old
2. Seeking contraception
3. Has not used hormonal contraception for the last three months

Participant type(s)

Mixed

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

16 years

Upper age limit

25 years

Sex

Female

Total final enrolment

80

Key exclusion criteria

Both groups of subjects (patients with type 1 diabetes and healthy controls):

1. Chronic conditions such as celiac sprue, epilepsy, cardiopulmonary or gastrointestinal conditions are present.
2. Use of steroidal medication.
3. Contraindications for using hormonal contraception (WHO criteria: migraine with aura, prothrombotic problems, etc)

Persons with type 1 diabetes:

1. HbA1c higher than 12.9%
2. Honeymoon period
3. Other type of diabetes

Healthy controls:

1. Menstrual cycle abnormalities according to the American Academy of Pediatrics.

Date of first enrolment

01/07/2017

Date of final enrolment

30/03/2020

Locations

Countries of recruitment

Chile

Study participating centre

University of Chile

Institute of Maternal and Child Research (IDIMI)

Santa Roa 1234

Santiago

Chile

8360160

Study participating centre

Instituto Chilena de Medicina Reproductiva (ICMER)

José Victorino Lastarria 29 dp 101

Santiago

Chile

8320165

Sponsor information

Organisation

University of Chile

ROR

<https://ror.org/04teye511>

Funder(s)

Funder type

Research council

Funder Name

Fondo Nacional de Desarrollo Científico y Tecnológico

Alternative Name(s)

National Fund for Scientific and Technological Development, El Fondo Nacional de Desarrollo Científico y Tecnológico, FONDECYT

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Chile

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Ethel Codner at ecodner@med.uchile.cl

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		25/11/2023	30/01/2024	Yes	No
Results article		07/05/2024	22/07/2025	Yes	No