

A placebo-controlled trial of anti-TNF α chimeric monoclonal antibody (infliximab, remicade) in the modification of vascular disease markers in active rheumatoid arthritis

Submission date 05/03/2007	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 23/11/2007	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 14/06/2011	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Bruce Kirkham

Contact details

Rheumatology Department

4th floor

Thomas Guy House

Guy's Hospital

London

United Kingdom

SE1 9RT

bruce.kirkham@gstt.nhs.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

RJI 03/0139

Study information

Scientific Title

Acronym

DIVERT - Defining Infliximab Vascular Effects Rheumatoid arthritis Trial

Study objectives

That surrogate measures of vascular disease (pulse wave velocity, flow mediated dilatation, carotid-intimal media thickness), will improve after infliximab therapy in patients with active rheumatoid arthritis.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Guy's Hospital Research Ethics Committee, approved on 28 November 2003 (ref: RJI - 03/0139)

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Rheumatoid Arthritis

Interventions

Placebo controlled 2:1 randomisation, active infliximab (3 mg/kg intravenous) vs placebo infusion for 26 weeks, then open label until week 56, with placebo escape arm at week 14.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

infliximab

Primary outcome measure

The following will be assessed at baseline (Week 0), Week 24 and Week 56, unless indicated otherwise:

1. Endothelial function (Flow Mediated Dilatation [FMD]). This will be assessed at Week 8 and Week 16 in addition to the timepoints stated above.
2. Vascular structure:
 - 2.1. Pulse Wave Velocity [PWV]
 - 2.2. Augmentation Index [Aix]
 - 2.3. Carotid Intimal Medial Thickening [CIMT]

Secondary outcome measures

The following will be assessed at baseline (Week 0), Week 24 and Week 56, unless indicated otherwise:

1. RA disease activity:
 - 1.1. Modified Health Assessment Questionnaire (HAQ). This will be assessed at Week 8 and Week 16 in addition to the timepoints stated above.
 - 1.2. 28 swollen and tender joint counts. This will be assessed at Week 8 and Week 16 in addition to the timepoints stated above.
 - 1.3. Erythrocyte Sedimentation Rate (ESR)
 - 1.4. Patient Global Assessment (PGA) using a 100 mm visual analogue scale and Disease Activity Score 28 (DAS 28). This will be assessed at Week 8 and Week 16 in addition to the timepoints stated above.
2. CV risk factors:
 - 2.1. Systolic and diastolic Blood Pressure (BP)
 - 2.2. Body Mass Index (BMI)
 - 2.3. High sensitivity C-Reactive Protein (HsCRP)
 - 2.4. Serum fasting lipid profile (total cholesterol, High and Low Density Lipoprotein fractions [HDL, LDL] and triglycerides)
 - 2.5. Oxidised LDL sub-fractions
 - 2.6. Insulin resistance measured by log homeostasis model assessment (HOMA)
 - 2.7. Serum levels of soluble Intracellular Adhesion Molecules (ICAM)
 - 2.8. Vascular Cell Adhesion Molecules (VCAM) and adiponectin

Overall study start date

15/05/2003

Completion date

15/05/2005

Eligibility

Key inclusion criteria

1. RA defined by American College of Rheumatology criteria
2. Referred for TNF-blocking therapy according to the British Society of Rheumatology (BSR) criteria

3. Patients giving written informed consent
4. Patients failed two DMARDs including methotrexate
5. Disease Activity Score 28 (DAS 28) greater than 5.1 on two occasions four weeks apart
6. Patients taking methotrexate (≤ 25 mg/week)

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

30

Key exclusion criteria

1. Age < 18 years
2. History of ischemic heart disease, cerebrovascular disease, peripheral vascular disease, diabetes mellitus
3. Previous treatment with infliximab or any therapeutic agent targeted at reducing TNFa
4. Treatment with aspirin
5. Patients with evidence of current or previous infection with tuberculosis (TB)

Date of first enrolment

15/05/2003

Date of final enrolment

15/05/2005

Locations**Countries of recruitment**

England

United Kingdom

Study participating centre

Rheumatology Department

London

United Kingdom

SE1 9RT

Sponsor information

Organisation

Guy's & St Thomas' NHS Foundation Trust (UK)

Sponsor details

R&D Department
Connybeare House
Guy's Hospital
St Thomas Street
London
England
United Kingdom
SE1 9RT
jackie.pullen@gstt.nhs.uk

Sponsor type

Hospital/treatment centre

Website

<http://www.guysandstthomas.nhs.uk/>

ROR

<https://ror.org/00j161312>

Funder(s)**Funder type**

Industry

Funder Name

Centocor BV (The Netherlands)

Results and Publications**Publication and dissemination plan**

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2009		Yes	No