

# Is knee-cap taping feasible and acceptable as part of a treatment package for painful knee-cap arthritis?

<b>Submission date</b>	<b>Recruitment status</b>	<input checked="" type="checkbox"/> Prospectively registered
14/11/2024	Recruiting	<input checked="" type="checkbox"/> Protocol
<b>Registration date</b>	<b>Overall study status</b>	<input type="checkbox"/> Statistical analysis plan
20/11/2024	Ongoing	<input type="checkbox"/> Results
<b>Last Edited</b>	<b>Condition category</b>	<input type="checkbox"/> Individual participant data
04/02/2026	Musculoskeletal Diseases	<input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Knee-cap arthritis (wear and tear) causes pain and disability for millions of adults in the UK. There are few effective treatments. Knee braces can help but they can be expensive and two-thirds of patients stop wearing them because either their leg is the wrong shape or it slips down. Therefore, this study will test the feasibility of a long-term simpler, cheaper intervention - knee-cap taping. If this study shows that taping is feasible and acceptable to patients, the ultimate goal is to carry out a large Health Technology Assessment-funded trial to test the extended effectiveness of self-managed knee-cap taping in painful knee-cap arthritis. This study aims to see if knee-cap taping is feasible and acceptable as part of a treatment package for painful knee-cap arthritis.

### Who can participate?

People with painful knee-cap arthritis

### What does the study involve?

Participants will receive the treatment package of best practice care and taping for 3 months. The number of participants willing to enrol will be recorded, their adherence to the intervention, how precisely they applied the tape, if taping had any side effects, and how many people did not complete the trial. Then, 15 participants will participate in one-to-one semi-structured interviews about their experiences of using the tape. Participants will provide advice regarding the treatment package and give insights to inform the development of a large, long-term trial of the effectiveness of taping.

### What are the possible benefits and risks of participating?

Participants will benefit from improvement in their knee pain over the 12-week trial period, the aesthetics of using the tape over wearing a bulky knee brace, the comfort of wearing tape over the knee brace, and taping method training (for future use). The clinical follow-up appointments at 2, 6 and 12 weeks, mean being seen more frequently than routine clinic visits.

There are no foreseen ethical, legal or management issues arising from this study. There may be minimal risk to the participants in Work Package 2 (WP2) and Work Package 3 (WP3), these have been fully disclosed in the Patient Information Sheet.

WP2 risks:

1. Reaction to tape - will have an exclusion criterion that removes people with known allergies to tape, psoriasis, broken skin or lesions in the area. If they have an unknown allergy and react to the tape they will be withdrawn and referred to their GP for treatment.
2. No improvement/worsening of pain - participants will be referred back to their GP or referring clinician at the end of the study for the next step of treatment.
3. Incorrect self-application of tape - this would just lead to inefficacy. The week two visit is in place to check the participants are using the correct method.

WP3 risks:

Participants may find discussing their experiences of taping or knee pain upsetting, and they are asked to let the team know if they wish to take a break or end the interview. If during the interview the participant reveals that their safety or the safety of others may be at risk, this information will be discussed with individuals outside the study team. Lastly, there may be a risk of a breach of confidentiality. Extensive systems are in place to try to prevent this and are detailed this in the PIS. Two PPIE workshops were held in Work Package 1. The purpose of these workshops was to co-design and refine the treatment package using patellar taping for kneecap arthritis, for use in a subsequent clinical trial feasibility study (WP2). It is believed that risks have been minimised from these workshops.

Where is the study run from?

The University of Manchester, Manchester Metropolitan University and the Sponsor is Manchester University NHS Foundation Trust (MFT).

When is the study starting and how long is it expected to run for?

December 2023 to February 2026

Who is funding the study?

NIHR Research for Patient Benefit (RfPB) programme

Who is the main contact?

Prof Michael Callaghan (Chief Investigator), Michael.callaghan@mmu.ac.uk  
Deborah Maskell (Project Manager), Deborah.Maskell@manchester.ac.uk

## Contact information

**Type(s)**

Scientific, Principal investigator

**Contact name**

Prof Michael Callaghan

**ORCID ID**

<https://orcid.org/0000-0003-3540-2838>

**Contact details**

Brooks Building Manchester Metropolitan University, Birley Campus  
Manchester  
United Kingdom  
M15 6GX  
+44 (0)1613060542  
Michael.callaghan@mmu.ac.uk

#### Type(s)

Public

#### Contact name

Ms Deborah Maskell

#### Contact details

TAPE-it Project Manager  
Division of Musculoskeletal and Dermatological Sciences  
The University of Manchester, Room 2.805 Stopford Building, Oxford Road  
Manchester  
United Kingdom  
M13 9PT  
-  
Deborah.Maskell@manchester.ac.uk

## Additional identifiers

**Integrated Research Application System (IRAS)**  
322953

**National Institute for Health and Care Research (NIHR)**  
204993

**Central Portfolio Management System (CPMS)**  
54632

## Study information

#### Scientific Title

The feasibility of knee taping in painful patellofemoral joint osteoarthritis (PFJOA): TAPE-it

#### Acronym

TAPE-it

#### Study objectives

The hypothesis is that medical sports taping of the knee will be both feasible and acceptable to participants as part of a treatment package for painful patellofemoral osteoarthritis.

#### Ethics approval required

Ethics approval required

#### Ethics approval(s)

## **Study design**

Non-randomised study

## **Primary study design**

Interventional

## **Study type(s)**

Treatment

## **Health condition(s) or problem(s) studied**

Medical sports taping of the knee for painful patellofemoral osteoarthritis

## **Interventions**

Before the original application: Five patients were invited to be introduced to the project concept and design. Their feedback helped inform the Work Package 1 (WP1) part of the study.

WP1: Before this protocol, two workshops were held in January and February 2024. This involved 11 participants being taken through the study concept, and then shown the taping methods and planned training materials for the actual intervention study. This was a PPIE exercise to ensure the right questions were being asked and, the best methods used and were meaningful. The findings from WP1 fully informed the WP2 methods described below.

## WP2

1. Screening: Manchester Local Care Organisation (MLCO) and Salford Royal Hospital: Clinicians refer potentially eligible patients with PFJOA to the study team either directly to CI or Study Team via secure NHS or NHS.net email address or online Trust system. Participants are given a Participant Information Sheet (PIS) for at least 24 hours to digest the information and ask questions. A member of the research team calls the participant to book a baseline visit.
2. Baseline Visit (30 - 60 minutes): face-to-face in a private room in the NIHR Manchester Clinical Research Facility.
  - 2.1. Aims explained and any questions from PIS answered.
  - 2.2. Provide written informed consent for trial.
  - 2.3. Clinical examination and eligibility questions asked to diagnose PFJOA and confirm eligibility (ineligible patients will be screened fail and referred back to GP or referring clinician).
  - 2.4. Determine aggravating activity, assess the best taping method, show the participant how to apply tape, and provide a Patient Treatment Package.
  - 2.5. Supply two weeks pre-cut tape.
- 2.6. Baseline Assessments: electronic or paper copy knee pain (VASNA) and knee function (KOOS-PF) questionnaires
- 2.7. Two-week assessment (20 minutes): Check taping treatment accuracy and correct if required. Supply 10 weeks pre cut tape. Knee pain (VASNA) and knee function (KOOS-PF) questionnaires and adverse events recording
- 2.8. Six-week assessment (15 minutes): Telephone assessment - knee pain (VASNA), function (KOOS-PF) questionnaires and adverse event recording.
- 2.9. Twelve-week assessment (15 minutes): Telephone assessment - knee pain (VASNA), function (KOOS-PF) questionnaires and adverse event recording

End of WP2: all completing participants will be given a PIS for WP3 and asked if they might want to take part. Then a sample of fifteen consenting participants will be purposefully selected (to show a range of age, sex, and ethnicity) to participate in the one-to-one interviews in WP3 below:

WP3:

1. Participants were given a PIS and 24 hours to consider the study and whether they would like to take part.
2. Full Informed Consent was taken and demographic information was collected. A semi-structured interview (approx 60 minutes) will be conducted to gain an in-depth understanding of the participant's experience of WP2.
3. Interviews will be conducted by a trained Research Associate (RA) either in person, online via Teams or over the phone depending on participants' preference. In-person interviews will be held in a private room at the Brooks Building, Birley Campus, Manchester Metropolitan University.
4. Discussions will be from an interview question list but also guided by each patient's response to allow exploration of unanticipated issues/feedback. Emerging findings from each interview will iteratively feed into subsequent interviews.
5. Interviews will be digitally recorded, professionally transcribed and then analysed by the Research Associate.

### **Intervention Type**

Other

### **Primary outcome(s)**

Current primary outcome(s)

Participant uptake is assessed as follows:

1. Recruitment numbers measured using study data at baseline
2. % dropout/withdrawal numbers measured using study data at 2, 6 and 12 weeks

---

Previous primary outcomes as of 20/01/2026:

1. Recruitment rate: number of participants recruited over the recruitment period.
2. Intervention adherence: determined by self-report at 2, 6 and 12 weeks
3. Self-report number of tape strips used at weeks 2, 6, 12.
4. Retention: proportion of participants who were contactable by telephone at 12 weeks.
5. Understanding participants' experience of taping, the acceptability of the intervention and follow-up methods through semi-structured interviews conducted over the post intervention follow-up period.
6. Willingness to be randomised: proportion of eligible patients approached who would be willing to be randomised to a 'no-tape' group in a future RCT.

---

Previous primary outcomes:

Participant uptake is assessed as follows:

1. Recruitment numbers measured using study data at baseline
2. % dropout/withdrawal numbers measured using study data at 2, 6 and 12 weeks

## **Key secondary outcome(s))**

Current key secondary outcome(s) as of 04/02/2026:

- . Correct tape application measured using study observation by the CI at the 2-week visit. Advice and adjustments to the taping method if required by the CI.
2. Adherence measured using a count of the remaining tape strips at weeks 2,6,12: Approximate amount of time of tape wearing measured using patient self-reported hours per day and days per week of tape wearing at weeks 2, 6 and 12.
3. Reduction in knee pain measured using the 11-point (0-10) Visual Analogue Scale during a Nominated Activity (VASNA) and knee pain and function measured using the Knee Injury and Osteoarthritis Outcome Score for patellofemoral pain and osteoarthritis (KOOS-PF) at baseline, 2, 6 and 12 weeks

---

Previous secondary outcomes as of 20/01/2026:

Reduction in knee pain measured using the 11-point (0-10) Visual Analogue Scale during a Nominated Activity (VASNA) and knee pain and function measured using the Knee Injury and Osteoarthritis Outcome Score for patellofemoral pain and osteoarthritis (KOOS-PF) at baseline, 2, 6 and 12 weeks.

---

Previous secondary outcomes:

1. Correct tape application measured using study observation by the CI at the 2-week visit. Advice and adjustments to the taping method if required by the CI.
2. Adherence measured using a count of the remaining tape strips at weeks 2,6,12: Approximate amount of time of tape wearing measured using patient self-reported hours per day and days per week of tape wearing at weeks 2, 6 and 12.
3. Reduction in knee pain measured using the 11-point (0-10) Visual Analogue Scale during a Nominated Activity (VASNA) and knee pain and function measured using the Knee Injury and Osteoarthritis Outcome Score for patellofemoral pain and osteoarthritis (KOOS-PF) at baseline, 2, 6 and 12 weeks

## **Completion date**

28/02/2026

## **Eligibility**

### **Key inclusion criteria**

1. Adults over 40 years of age
2. Pain predominantly over the patella (anterior knee pain) and greater than medial or lateral knee pain
3. Clinically significant patellofemoral pain/anterior knee pain on weight-bearing activity such as stair ascent/descent, sit-stand scored at 4 or above on an 11-point (0-10) visual analogue scale (VAS)
4. Already on a stable/regular dose of analgesia
5. Able to understand English (or English speaking) and give full informed consent
6. Willing to participate

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Mixed

**Lower age limit**

40 years

**Upper age limit**

100 years

**Sex**

All

**Total final enrolment**

0

**Key exclusion criteria**

1. Symptoms not attributable to predominant patellofemoral OA or attributable only to medial or lateral knee compartment OA
2. Previous major surgery in the knee to be treated (partial/total knee replacement, knee fracture or knee realignment surgery, high tibial osteotomy)
3. Diagnosis of rheumatoid arthritis, gout or other forms of inflammatory arthritis
4. Cancer within the last 5 years (except non-melanoma skin cancers)
4. Planned knee replacement surgery or other knee surgery in the next 6 months
5. Responded to a steroid or viscosupplementation injection to the painful knee in the last 3 months
6. Those with a known allergy to elasticated tape, with fragile or very sensitive skin, psoriasis, or with lesions/rash/open wound in the area where the tape will be applied
7. Significant neurological disorder (e.g. stroke, dementia, MS, Parkinson's) affecting cognitive ability

**Date of first enrolment**

16/12/2024

**Date of final enrolment**

28/02/2026

## Locations

**Countries of recruitment**

United Kingdom

England

**Study participating centre**  
**Manchester Metropolitan University**  
All Saints  
Grosvenor Square  
Manchester  
England  
M15 6BH

**Study participating centre**  
**University of Manchester**  
Oxford Road  
Manchester  
England  
M13 9WL

**Study participating centre**  
**Manchester University NHS Foundation Trust**  
Cobbett House  
Oxford Road  
Manchester  
England  
M13 9WL

## **Sponsor information**

**Organisation**  
Manchester University NHS Foundation Trust

**ROR**  
<https://ror.org/00he80998>

## **Funder(s)**

**Funder type**  
Government

**Funder Name**  
Research for Patient Benefit Programme

**Alternative Name(s)**

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

## Results and Publications

### Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Data sharing statement to be made available at a later date

#### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>	version 1.1	31/10/2024	20/01/2026	No	Yes
<a href="#">Protocol file</a>	version 1.0	02/10/2024	19/12/2024	No	No