

Effects of a Gyejibongnyeong-hwan on dysmenorrhea caused by blood stagnation

Submission date 19/04/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 30/04/2010	Overall study status Completed	<input checked="" type="checkbox"/> Protocol
Last Edited 07/11/2013	Condition category Urological and Genital Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Seong-Gyu Ko

Contact details
1 Hoegi-dong
Dongdaemun-gu
Seoul
Korea, South
130-701

Additional identifiers

Protocol serial number
CCRG_08_03

Study information

Scientific Title
Effects of a Gyejibongnyeong-hwan on dysmenorrhea caused by blood stagnation: a randomized, double-blinded, placebo-controlled, multicenter study

Study objectives

Gyejibongnyeong-hwan reduces dysmenorrhea caused by blood stagnation more than placebo control.

As of 21/04/2011 the anticipated end date for this trial has been extended from 30/06/2010 to 30/06/2011.

Ethics approval required

Old ethics approval format

Ethics approval(s)

1. Institutional Review Board (IRB) of Kyung Hee Oriental Medical Center approved on the 18/08/2008 (ref: KOMC IRB 2008-07)
2. IRB of Wonkwang University Sanbon oriental medical center approved on the 24/02/2009 (ref: WONSBBH IRB 2009-02)
3. IRB of Kyungwon Gil Oriental Medical Hospital approved on the 02/02/2009 (ref: 09-101)

Study design

Randomised double-blind parallel group, placebo-controlled trial phase IV study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Dysmenorrhea (menstrual pain) caused by blood stagnation

Interventions

Patients will be randomised to receive:

1. Gyejibongnyeong-hwan, a Korean herbal remedy consisting of Cinnamomi ramulus, Poria, Moutan cortex, Persicae semen (peach seed) and Paeoniae radix. 1200 mg/day (400 mg x 3 daily)
2. Placebo control (x 3 daily)

Participants will receive the intervention for approx 8 weeks (two menstrual cycles) and be followed on week 4 and 12 after the end of the treatment. These times will vary depending on the duration of the period/cycle.

Participants will visit six times, screening, visit 1 (week 0), visit 2 (week 4), visit 3 (week 8), visit 4 (week 12), visit 5 (week 20).

The total duration of the trial will be 20 weeks.

Intervention Type

Other

Phase

Phase IV

Primary outcome(s)

Pain, assessed by Visual Analogue Scale (VAS), measured at screening, baseline and weeks 4, 8, 12 and 20

Key secondary outcome(s)

1. Blood Stagnation Scale (based on Korean Medicine), measured at screening, baseline and weeks 4 and 8
2. Quantity of pain killer pills during period, recorded at screening, baseline and weeks 4, 8, 12 and 20
3. Short Form McGill Pain Questionnaire, measured at screening, baseline and weeks 4, 8, 12 and 20
4. Cox Menstrual Symptom Scale, measured at screening, baseline and weeks 4, 8, 12 and 20
5. Heart rate variability (HRV) outcome, measured at screening, baseline and weeks 4 and 8

Completion date

30/06/2011

Eligibility

Key inclusion criteria

1. Female aged 18 to 35 years
2. Women with a period cycle of 30 ± 3 days during last 3 months
3. Women who have menstrual pain (dysmenorrhea) over 6 degrees by Visual Analogue Scale (VAS)
4. Women who are diagnosed with blood stagnation by two oriental medical gynaecologic specialists
5. Given written informed consent form
6. Given written informed consent form of genetic study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Key exclusion criteria

1. Women who have major neuro-psychiatric disorder or have history of major neuro-psychiatric disorder (schizophrenia, epilepsy, alcohol abuse, anorexia etc)
2. Women who are planning to have baby or do not agree to use appropriate contraception (oral pill, hormone contraception, intrauterine device, condom etc)
3. Women who are taking anti-depressant, anti-serotonin barbiturate, psychotropic drugs

Date of first enrolment

19/05/2009

Date of final enrolment

30/06/2011

Locations

Countries of recruitment

Korea, South

Study participating centre

1 Hoegi-dong

Seoul

Korea, South

130-701

Sponsor information

Organisation

Korea Health Industry Development Institute (South Korea)

ROR

<https://ror.org/00fdzyk40>

Funder(s)

Funder type

Research organisation

Funder Name

Korea Health Industry Development Institute (South Korea)

Alternative Name(s)

KHIDI

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Korea, South

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/01/2013		Yes	No
Protocol article	protocol	05/01/2012		Yes	No