A comparison of response to treatment in patients with Myelodysplastic Syndrome /Myeloproliferative Neoplasm (MDS/MPN) Overlap Syndromes taking ASTX727 versus best supportive care

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
11/05/2022		Protocol		
Registration date	Overall study status Ongoing	Statistical analysis plan		
13/05/2022		Results		
Last Edited 17/06/2024	Condition category Cancer	Individual participant data		
		Record updated in last year		

Plain English summary of protocol

Background and study aims

MDS/MPN Overlap Syndromes are rare bone marrow cancers. For most patients there are no effective treatments available. Patients experience bone marrow failure, leading to fatigue, infections and bleeding, with a damaging overgrowth of white blood cells. Most patients don't live for more than two years from diagnosis. AMMO will test a new drug called ASTX727 to see how effective it is at treating and extending/improving the lives of people with these diseases. ASTX727 is a new tablet "hypomethylating drug". These work differently from typical chemotherapy, by altering how genes are switched on and off. They do this by blocking a protein whose job it is to "mark" which genes should be switched on in any particular cell. The hypomethylating drug azacitidine has been used for several years, including for some MDS/MPN patients. It can improve blood counts and symptoms, even putting the cancer into remission and helping patients live longer. However, it must be given as injections on repeated visits to hospital, and isn't available for most MDS/MPN patients. We want to know if ASTX727 can provide similar or possibly better results across all MDS/MPN Overlap syndromes, safely and more conveniently, and whether it overall improves the lives of patients.

Who can participate?

Adults over 18 years, with a diagnosis of MDS/MPN overlap syndromes

What does the study involve?

Patients will be allocated at random to receive either best supportive care or ASTX727. For patients on best supportive care they will go on to be treated as they would in normal routine care which would differ between patients and may include hydroxycarbamide, transfusional support and management of infections. Patients allocated to receive ASTX727 will take ASTX727 orally once daily on days 1-5 of 28 day cycles for up to 6 cycles. If the patient experiences a benefit from the drug they may continue to take it until disease progression,

unacceptable toxicity or patient chooses to discontinue. Patients will come into hospital on day 1 of each cycle (and day 15 of cycle 1) for standard tests including blood tests and a physical exam and patients will have a bone marrow aspirate at cycles 3 and 6 (and every 6 cycles for patients continuing ASTX727) to monitor their progress. Patients will be followed up annually until the end of the study.

What are the possible benefits and risks of participating?

There is no guaranteed benefit of taking part in this study because we do not yet know which of the two treatment approaches is better. ASTX727 has shown promise in early trials and so may be a useful treatment for these diseases; it may, however, not provide any overall benefit compared with best supportive care. The information gained from this study will help inform treatment for other people in the future. The main risk to taking part is side effects from ASTX727 or hydroxycarbamide. These will be outlined in the patient information sheet.

Where is the study run from? University of Birmingham (UK)

When is the study starting and how long is it expected to run for? February 2020 to December 2025

Who is funding the study? BLOODWISE (Blood Cancer UK)

Who is the main contact? AMMO@trials.bham.ac.uk

Contact information

Type(s)

Scientific

Contact name

Ms Louise Hopkins

Contact details

Centre for Clinical Haematology Queen Elizabeth Hospital Birmingham United Kingdom B15 2TH +44 121 371 7861 AMMO@trials.bham.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS) 2021-004585-35

Integrated Research Application System (IRAS) 1004656

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

CPMS 52390, Grant Code 20001, IRAS 1004656

Study information

Scientific Title

A randomised phase 2 study of ASTX727 versus best supportive care in(MDS/MPN) Overlap Syndromes

Acronym

AMMO

Study objectives

To compare the overall response rate of ASTX727 vs best supportive care +/- hydroxycarbamide, according to IWG 2015 MDS/MPN proposed response criteria. ORR includes patients who achieve best response of complete remission (CR), partial remission (PR), marrow response (MR) or clinical benefit (CB).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 22/02/2022, West of Scotland REC 1 (West of Scotland Research Ethics Service, Ward 11, Dykebar Hospital, Grahamston Road, Paisley, PA2 7DE, UK; +44 (0)141 3140212; WoSREC1@ggc.scot.nhs.uk), ref: 22/WS/0011

Study design

Interventional randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Myelodysplastic Syndrome/Myeloproliferative Neoplasm (MDS/MPN) Overlap Syndromes

Interventions

Patients will be randomised via a computer system, the site staff will log on to the study portal in order to randomise a patient between best supportive care and the experimental arm (ASTX727).

Patients on best supportive care will receive standard of care treatments which may include hydroxycarbamide, transfusional support and management of infections. We will collect this information for 6 months from trial entry (or 6 months + 28 days for those on hydroxycarbamide). Best supportive care will differ between patients and will be at the Investigators discretion. Patients allocated to receive ASTX727 will take ASTX727 orally once

daily on days 1-5 of 28 day cycles for up to 6 cycles. If the patient experiences a benefit from the drug they may continue to take it until disease progression, unacceptable toxicity or patient chooses to discontinue.

All patients will come into hospital on day 1 of each cycle (and day 15 of cycle 1) for standard tests including blood tests and a physical exam and patients will have a bone marrow aspirate at cycles 3 and 6 (and every 6 cycles for patients continuing ASTX727) to monitor their progress. Patients will be followed up annually until the end of the study.

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

ASTX727

Primary outcome(s)

Best overall response at end of Cycle 6, defined as the patient experiencing Complete Remission (CR), Partial Remission (PR), Marrow Response (MR) or Clinical Benefit (CB) according to the IWG MDS/MPN response criteria. Any patient for whom neither response assessment (post Cycle 2; post Cycle 6) is recorded will be classed as a non-responder.

Key secondary outcome(s))

- 1. Response will be assessed according to the IWG MDS/MPN response criteria at the end of cycles 2 and 6.. Patients will be classed as responding if they achieve complete remission (CR), partial remission (PR), marrow response (MR) or clinical benefit (CB). Patients with no response assessment available will be classed as a non-responder.
- 2. Transformation-free survival (TFS) defined as time from entry to the trial to time of transformation to AML or death. Patients who are alive and have not transformed to AML at the time of the analysis will be censored at date last seen. TFS will be assessed as time from study entry to time of transformation to AML or death.
- 3. Progression-free survival (PFS) defined as time from entry to the trial to first documentation of progressive disease (PD), as defined by the 2015 IWG MDS/MPN response criteria, or death from any cause. Patients who are alive and progression free at the time of analysis will be censored at their date last seen. PFS will be assessed from time of study entry to time of progressive disease.
- 4. Overall Survival (OS) defined as time between trial entry and date of death from any cause. Patients who are alive at the time of analysis will be censored at date last seen. OS will be assessed as time from study entry to date of death.
- 5. Treatment-related toxicity defined as the number of patients who experience one or more grade 3 or greater adverse event or serious adverse event of any grade. Toxicity will be assessed from start of trial treatment to 28 days post treatment discontinuation.
- 6. Quality of life assessed using EQ5D-5L and the MPN-SAF tools at baseline, cycle 3, cycle 6, 1 year post randomisation and thereafter every 6 months in patients continuing on treatment.

Completion date

30/12/2025

Eligibility

Key inclusion criteria

- 1. ≥18 years of age at the time of trial entry
- 2. Morphologically confirmed diagnosis of MDS/MPN (excluding JMML), in accordance with WHO 2016 diagnostic criteria, with any of the following characteristics:
- 2.1. CMML-2 disease stage [CMML only]
- 2.2. CPSS [Such et al Blood 2013] or CPSS-Mol [Elena et al Blood 2016] score of intermediate-2 or high risk [CMML only]
- 2.3. Other patients with one or more of the following:
- 2.3.1. Bone Marrow blasts >10% (including promonocytes)
- 2.3.2. Adverse risk cytogenetics (as defined by CPSS or MDS R-IPSS)
- 2.3.3. WCC \geq 50 (or \geq 30 with symptoms attributable to myeloproliferation)
- 2.3.4. RBC transfusion dependence with pre-transfusion Hb <90 g/L
- 2.3.5. Symptomatic anaemia (with Hb \leq 100 g/L)
- 2.3.6. Thrombocytopenia (Plt \leq 50 x 10^9/L)
- 2.3.7. Symptomatic splenomegaly
- 2.3.8. Systemic symptoms with no alternative explanation (including weight loss ≥10% of baseline over previous 6 months)
- 2.3.9. Symptomatic extramedullary involvement (e.g. skin infiltration, serous effusions)
- 3. Treatment-naïve for prior hypomethylating agent, intensive chemotherapy or other disease-modifying anti-neoplastic therapy (e.g. lenalidomide); patients may have received prior hydroxycarbamide, recombinant erythropoietin, danazol, interferon or anagrelide.
- 4. ECOG performance status of 0,1 or 2 at trial entry (Appendix 3).
- 5. Life expectancy of \geq 3 months at trial entry, as assessed by the treating physician.
- 6. Must have adequate hepatic, renal and endocrine function during screening as demonstrated by the following:
- 6.1. ALT and/or AST ≤3x upper limit of normal (ULN);
- 6.2. Total bilirubin \leq 1.5x ULN or \leq 2x ULN if upon judgement of the treating investigator the hyperbilirubinaemia is due to extramedullary haematopoiesis related to the underlying MDS /MPN or to Gilbert's disease;
- 6.3. Serum creatinine $\leq 1.5 \times \text{ULN}$ or estimated creatinine clearance $>/=30 \text{ml/min}/1.73 \text{m}^2$.
- 7. Patient willing and able to comply with scheduled visits, treatment plan and other study procedures.
- 8. Patient able to provide written informed consent for the trial

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Αll

Total final enrolment

Key exclusion criteria

- 1. Patients eligible for intensive chemotherapy and/or allogeneic haematopoietic stem cell transplantation (HSCT).
- 2. CMML with eosinophilia and 5q33 abnormality.
- 3. Previous cytotoxic chemotherapy for MDS/MPN, except hydroxycarbamide.
- 4. Prior hypomethylating agent exposure.
- 5. Transformation to AML (≥20% myeloid blasts in bone marrow or peripheral blood at screening).
- 6. Prior organ transplantation, including allogeneic haematopoietic stem cell transplant (HSCT).
- 7. Known or suspected central nervous system disease involvement.
- 8. Known history of clinically significant or uncontrolled cardiac disease, including recent history (within 6 months) of unstable angina, acute myocardial infarction, NYHA class III or IV congestive cardiac failure, or clinically significant arrhythmia.
- 9. Other active malignancy, not including localized non-melanoma skin cancer, cervical carcinoma in situ, breast ductal carcinoma in situ of the breast, or localized prostate cancer controlled with hormone therapy. Patients with history of other cancers should be free of disease without ongoing anti-neoplastic therapy for at least 2 years.
- 10. Receipt of wide-field radiotherapy (including therapeutic radioisotopes) \leq 28 days or limited field radiation for palliation \leq 14 days prior to starting any study medications (or has not recovered from side effects of such therapy).
- 11. Active, uncontrolled infection. Patients with infection under control with active treatment are eligible.
- 12. Pregnant and lactating patients (patients of childbearing potential must have a negative pregnancy test prior to study entry).
- 13. Females of childbearing potential, and their partners, not willing to use adequate contraception during and for up to 6 months after treatment.
- 14. Any other concurrent serious or unstable medical, psychiatric, familial, geographic or sociological condition that in the investigator's opinion would jeopardise the patient's ability to comply with the protocol.

Date of first enrolment 31/05/2022

Date of final enrolment 22/05/2024

Locations

Countries of recruitment

United Kingdom

England

Northern Ireland

Scotland

Wales

Study participating centre The Christie NHS Foundation Trust

550 Wilmslow Road Withington Manchester United Kingdom M20 4BX

Study participating centre Oxford University Hospitals NHS Foundation Trust

John Radcliffe Hospital Headley Way Headington Oxford United Kingdom OX3 9DU

Study participating centre University College London Hospitals NHS Foundation Trust

250 Euston Road London United Kingdom NW1 2PG

Study participating centre University Hospitals Birmingham NHS Foundation Trust

Queen Elizabeth Hospital Mindelsohn Way Edgbaston Birmingham United Kingdom

B15 2GW

Study participating centre NHS Greater Glasgow and Clyde

J B Russell House Gartnavel Royal Hospital 1055 Great Western Road Glasgow Glasgow United Kingdom G12 0XH

Study participating centre Belfast Health and Social Care Trust

Trust Headquarters A Floor - Belfast City Hospital Lisburn Road Belfast United Kingdom BT9 7AB

Study participating centre

Nottingham University Hospitals NHS Trust - Queen's Medical Centre Campus

Nottingham University Hospital Derby Road Nottingham United Kingdom NG7 2UH

Study participating centre Sheffield Teaching Hospitals NHS Foundation Trust

Northern General Hospital Herries Road Sheffield United Kingdom S5 7AU

Study participating centre University Hospital Southampton NHS Foundation Trust

Southampton General Hospital Tremona Road Southampton United Kingdom SO16 6YD

Study participating centre Leeds Teaching Hospitals NHS Trust

St. James's University Hospital Beckett Street Leeds United Kingdom LS9 7TF

Study participating centre Cardiff & Vale University Lhb

Woodland House Maes-y-coed Road Cardiff United Kingdom CF14 4HH

Study participating centre NHS Grampian

Summerfield House 2 Eday Road Aberdeen United Kingdom AB15 6RE

Study participating centre Kings College Hospital

King's College Hospital NHS Foundation Trust Denmark Hill London United Kingdom SE5 9RS

Sponsor information

Organisation

University of Birmingham

ROR

https://ror.org/03angcq70

Funder(s)

Funder type

Charity

Funder Name

Blood Cancer UK

Alternative Name(s)

Blood Cancer UK Research

Funding Body Type

Private sector organisation

Funding Body Subtype

Trusts, charities, foundations (both public and private)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from the trial management group by contacting ammo@trials.bham.ac.uk following the end of the study. Each request will be considered on a case by case basis and any research must have the relevant approvals in place.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes