

International study of caesarean section surgical techniques: a randomised fractional factorial trial and follow-up study

Submission date 29/11/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 17/01/2007	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 10/05/2016	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Study website
<http://www.npeu.ox.ac.uk/coronis>

Contact information

Type(s)
Scientific

Contact name
Prof Peter Brocklehurst

Contact details
National Perinatal Epidemiology Unit
University of Oxford
Old Road Campus
Oxford
United Kingdom
OX3 7LF
-
peter.brocklehurst@npeu.ox.ac.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

G0500959/G1000469

Study information

Scientific Title

International study of caesarean section surgical techniques: a randomised fractional factorial trial and follow-up study

Acronym

CORONIS

Study objectives

Study hypothesis amended as of 02/08/2007:

The CORONIS Trial aims to evaluate alternative surgical techniques for five specific aspects of caesarean section technique in a large pragmatic randomised controlled trial, to help determine which methods lead to an optimum outcome for women and their infants.

During the trial development phase it was known as the International CAESAR study.

Ethics approval required

Old ethics approval format

Ethics approval(s)

1. Oxford Tropical Research Ethics Committee (OXTREC), 05/03/2007, ref: 013-06
2. Follow-up study: OXTREC, 17/11/2010, ref: 013-06

Study design

Multicentre factorial randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Information leaflet in English for main trial <http://www.npeu.ox.ac.uk/downloads/coronis/CORONIS-Information-for-Women.pdf> and for the follow-up study <https://www.npeu.ox.ac.uk/files/downloads/coronis-follow-up/Information-for-women-V1-June-2011.pdf>

Health condition(s) or problem(s) studied

Pregnancy and Childbirth

Interventions

There are five pairs of interventions being tested; however, each participating hospital will only take part in three of these five possible comparisons. The interventions being compared are:

1. Blunt versus sharp abdominal entry
2. Exteriorisation of the uterus versus intra-abdominal repair
3. Single versus double layer closure of the uterus
4. Closure versus non-closure of the peritoneum (pelvic and parietal)
5. Chromic catgut versus Vicryl for uterine closure

Intervention Type

Procedure/Surgery

Primary outcome measure

Current information as of 29/09/10:

Death or maternal infectious morbidity (one or more of the following: antibiotic use for maternal febrile morbidity during postnatal hospital stay, antibiotic use for endometritis, wound infection or peritonitis) or further operative procedures; or blood transfusion of >1 unit of whole blood or packed cells.

Initial information at time of registration:

Death or serious maternal morbidity (one or more of the following: antibiotic use for maternal febrile morbidity during postnatal hospital stay, antibiotic use for endometritis or wound infection; further operative procedures on the wound; blood transfusion).

Secondary outcome measures

All within six weeks of delivery unless stated otherwise:

1. Death
2. Febrile morbidity
3. Endometritis
4. Wound infection treated with antibiotics
5. Operative procedures on wound
6. Pain
7. Blood transfusion
8. Interventions used for severe primary Post-Partum Haemorrhage (PPH)
9. Stillbirth after trial entry
10. Apgar score less than three at five minutes
11. Laceration of baby at time of caesarean section
12. Death of the baby by six weeks of age
13. Other severe maternal morbidity

Health service utilisation:

1. Duration of operation (from incision to closure)
2. Duration of hospital stay post-caesarean section
3. Duration of stay in Intensive Care Unit post-caesarean section
4. Number and duration of re-admissions to hospital within six weeks of the caesarean section

Added as of 13/03/2013: Follow-up Study Outcomes

Womens health and mortality

1. Following the CORONIS birth and before any subsequent pregnancy, any new onset or worsening of:

- 1.1. pelvic pain
- 1.2. dysmenorrhoea
- 1.3. deep dyspareunia
- 1.4. urinary symptoms of poor stream and/or frequency which did not respond to antibiotics
2. Diagnostic laparoscopy or diagnostic laparotomy (not related to pregnancy)
3. Hysterectomy or tubal/ovarian surgery (not related to pregnancy)
4. Bladder or bowel damage in those women who have had surgery, excluding diagnostic laparoscopy and diagnostic laparotomy (not related to pregnancy).
5. Following the CORONIS birth, any new onset of:
 - 5.1. abdominal hernia
 - 5.2. bowel obstruction
6. Womans death
- Reproductive status
7. Number of women with no subsequent pregnancy
 - 7.1. Voluntary infertility
 - 7.2. Involuntary infertility
8. Use of fertility treatments
- Subsequent pregnancies
9. Number of women having any subsequent pregnancy and for these women, the following outcomes will be measured:
 - 9.1. Inter-pregnancy interval from the CORONIS birth to the end of the subsequent pregnancy (regardless of loss or birth)
 - 9.2. Miscarriage of the pregnancy subsequent to the CORONIS birth
 - 9.3. Ectopic pregnancy
 - 9.4. Gestation at delivery (by best estimate) of the first viable pregnancy (gestational age > 24 or >28 weeks depending on country specific definition)
 - 9.5. Stillbirth
 - 9.6. Neonatal death
 - 9.7. Mode of delivery:
 - 9.7.1. Non-instrumental vaginal
 - 9.7.2. Instrumental vaginal
 - 9.7.3. Pre-labour caesarean section
 - 9.7.4. In labour caesarean section
 - 9.8. Other pregnancy complications including: uterine rupture, uterine scar dehiscence, placenta praevia, morbidly adherent placenta, abruption, postpartum haemorrhage requiring transfusion, severe infection within 6 weeks postpartum, hysterectomy up to 6 weeks postpartum, manual removal of placenta,
 - 9.9. Bladder or bowel damage at the time of subsequent caesarean section
- CORONIS children morbidity and mortality
10. Death or serious morbidity of the child who was born at the time of CORONIS participation (Although no difference in death is expected, there is likely to be a time difference between sharp and blunt abdominal entry and this may, in theory, lead to more neonatal encephalopathy which may lead to a greater risk of later death).

Overall study start date

01/09/2006

Completion date

30/06/2015

Eligibility

Key inclusion criteria

Women will be eligible for trial entry if they are undergoing delivery by lower segment caesarean section through a transverse abdominal incision.

Note: women are eligible if they have had no previous, or one previous, caesarean section. Fever in labour is not a contra-indication to trial entry.

Added as of 13/03/2013: All women recruited into the CORONIS Trial and eligible for three year follow-up will have a health assessment conducted by a study assessment doctor.

Participant type(s)

Patient

Age group

Adult

Sex

Female

Target number of participants

15,729

Key exclusion criteria

Women will not be eligible for trial entry if there is clear indication for a particular surgical technique or material to be used, that interferes with any of the allocated interventions, e.g. in a previous vertical incision.

Date of first enrolment

01/09/2006

Date of final enrolment

30/06/2015

Locations**Countries of recruitment**

Argentina

Chile

England

Ghana

India

Kenya

Pakistan

Sudan

United Kingdom

Study participating centre

University of Oxford

Oxford

United Kingdom

OX3 7LF

Sponsor information

Organisation

University of Oxford (UK)

Sponsor details

Clinical Trials and Research Governance Office

Manor House

John Radcliffe Hospital

Headington

Oxford

England

United Kingdom

OX3 9DZ

-

Heather.House@admin.ox.ac.uk

Sponsor type

University/education

Website

<http://www.ox.ac.uk/>

ROR

<https://ror.org/052gg0110>

Funder(s)

Funder type

Research council

Funder Name

Medical Research Council (UK) (Grant ref: G0500959)

Alternative Name(s)

Medical Research Council (United Kingdom), UK Medical Research Council, MRC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	22/10/2007		Yes	No
Results article	results	20/07/2013		Yes	No
Protocol article	follow-up protocol	21/11/2013		Yes	No
Results article	3-year follow-up results	02/07/2016		Yes	No