

# International study of caesarean section surgical techniques: a randomised fractional factorial trial and follow-up study

<b>Submission date</b> 29/11/2006	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 17/01/2007	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 10/05/2016	<b>Condition category</b> Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

Not provided at time of registration

## Study website

<http://www.npeu.ox.ac.uk/coronis>

## Contact information

### Type(s)

Scientific

### Contact name

Prof Peter Brocklehurst

### Contact details

National Perinatal Epidemiology Unit  
University of Oxford  
Old Road Campus  
Oxford  
United Kingdom  
OX3 7LF

-

[peter.brocklehurst@npeu.ox.ac.uk](mailto:peter.brocklehurst@npeu.ox.ac.uk)

## Additional identifiers

EudraCT/CTIS number

IRAS number

**ClinicalTrials.gov number**

**Secondary identifying numbers**

G0500959/G1000469

## **Study information**

### **Scientific Title**

International study of caesarean section surgical techniques: a randomised fractional factorial trial and follow-up study

### **Acronym**

CORONIS

### **Study objectives**

Study hypothesis amended as of 02/08/2007:

The CORONIS Trial aims to evaluate alternative surgical techniques for five specific aspects of caesarean section technique in a large pragmatic randomised controlled trial, to help determine which methods lead to an optimum outcome for women and their infants.

During the trial development phase it was known as the International CAESAR study.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

1. Oxford Tropical Research Ethics Committee (OXTREC), 05/03/2007, ref: 013-06
2. Follow-up study: OXTREC, 17/11/2010, ref: 013-06

### **Study design**

Multicentre factorial randomised controlled trial

### **Primary study design**

Interventional

### **Secondary study design**

Randomised controlled trial

### **Study setting(s)**

Hospital

### **Study type(s)**

Treatment

### **Participant information sheet**

Information leaflet in English for main trial <http://www.npeu.ox.ac.uk/downloads/coronis/CORONIS-Information-for-Women.pdf> and for the follow-up study <https://www.npeu.ox.ac.uk/files/downloads/coronis-follow-up/Information-for-women-V1-June-2011.pdf>

### **Health condition(s) or problem(s) studied**

## Pregnancy and Childbirth

### Interventions

There are five pairs of interventions being tested; however, each participating hospital will only take part in three of these five possible comparisons. The interventions being compared are:

1. Blunt versus sharp abdominal entry
2. Exteriorisation of the uterus versus intra-abdominal repair
3. Single versus double layer closure of the uterus
4. Closure versus non-closure of the peritoneum (pelvic and parietal)
5. Chromic catgut versus Vicryl for uterine closure

### Intervention Type

Procedure/Surgery

### Primary outcome measure

Current information as of 29/09/10:

Death or maternal infectious morbidity (one or more of the following: antibiotic use for maternal febrile morbidity during postnatal hospital stay, antibiotic use for endometritis, wound infection or peritonitis) or further operative procedures; or blood transfusion of >1 unit of whole blood or packed cells.

Initial information at time of registration:

Death or serious maternal morbidity (one or more of the following: antibiotic use for maternal febrile morbidity during postnatal hospital stay, antibiotic use for endometritis or wound infection; further operative procedures on the wound; blood transfusion).

### Secondary outcome measures

All within six weeks of delivery unless stated otherwise:

1. Death
2. Febrile morbidity
3. Endometritis
4. Wound infection treated with antibiotics
5. Operative procedures on wound
6. Pain
7. Blood transfusion
8. Interventions used for severe primary Post-Partum Haemorrhage (PPH)
9. Stillbirth after trial entry
10. Apgar score less than three at five minutes
11. Laceration of baby at time of caesarean section
12. Death of the baby by six weeks of age
13. Other severe maternal morbidity

Health service utilisation:

1. Duration of operation (from incision to closure)
2. Duration of hospital stay post-caesarean section
3. Duration of stay in Intensive Care Unit post-caesarean section
4. Number and duration of re-admissions to hospital within six weeks of the caesarean section

Added as of 13/03/2013: Follow-up Study Outcomes

Womens health and mortality

1. Following the CORONIS birth and before any subsequent pregnancy, any new onset or worsening of:

- 1.1. pelvic pain
- 1.2. dysmenorrhoea
- 1.3. deep dyspareunia
- 1.4. urinary symptoms of poor stream and/or frequency which did not respond to antibiotics
2. Diagnostic laparoscopy or diagnostic laparotomy (not related to pregnancy)
3. Hysterectomy or tubal/ovarian surgery (not related to pregnancy)
4. Bladder or bowel damage in those women who have had surgery, excluding diagnostic laparoscopy and diagnostic laparotomy (not related to pregnancy).
5. Following the CORONIS birth, any new onset of:
  - 5.1. abdominal hernia
  - 5.2. bowel obstruction
6. Womans death
- Reproductive status
7. Number of women with no subsequent pregnancy
  - 7.1. Voluntary infertility
  - 7.2. Involuntary infertility
8. Use of fertility treatments
- Subsequent pregnancies
9. Number of women having any subsequent pregnancy and for these women, the following outcomes will be measured:
  - 9.1. Inter-pregnancy interval from the CORONIS birth to the end of the subsequent pregnancy (regardless of loss or birth)
  - 9.2. Miscarriage of the pregnancy subsequent to the CORONIS birth
  - 9.3. Ectopic pregnancy
  - 9.4. Gestation at delivery (by best estimate) of the first viable pregnancy (gestational age > 24 or >28 weeks depending on country specific definition)
  - 9.5. Stillbirth
  - 9.6. Neonatal death
  - 9.7. Mode of delivery:
    - 9.7.1. Non-instrumental vaginal
    - 9.7.2. Instrumental vaginal
    - 9.7.3. Pre-labour caesarean section
    - 9.7.4. In labour caesarean section
  - 9.8. Other pregnancy complications including: uterine rupture, uterine scar dehiscence, placenta praevia, morbidly adherent placenta, abruption, postpartum haemorrhage requiring transfusion, severe infection within 6 weeks postpartum, hysterectomy up to 6 weeks postpartum, manual removal of placenta,
  - 9.9. Bladder or bowel damage at the time of subsequent caesarean section
- CORONIS children morbidity and mortality
10. Death or serious morbidity of the child who was born at the time of CORONIS participation (Although no difference in death is expected, there is likely to be a time difference between sharp and blunt abdominal entry and this may, in theory, lead to more neonatal encephalopathy which may lead to a greater risk of later death).

#### **Overall study start date**

01/09/2006

#### **Completion date**

30/06/2015

## **Eligibility**

**Key inclusion criteria**

Women will be eligible for trial entry if they are undergoing delivery by lower segment caesarean section through a transverse abdominal incision.

Note: women are eligible if they have had no previous, or one previous, caesarean section. Fever in labour is not a contra-indication to trial entry.

Added as of 13/03/2013: All women recruited into the CORONIS Trial and eligible for three year follow-up will have a health assessment conducted by a study assessment doctor.

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Female

**Target number of participants**

15,729

**Key exclusion criteria**

Women will not be eligible for trial entry if there is clear indication for a particular surgical technique or material to be used, that interferes with any of the allocated interventions, e.g. in a previous vertical incision.

**Date of first enrolment**

01/09/2006

**Date of final enrolment**

30/06/2015

**Locations****Countries of recruitment**

Argentina

Chile

England

Ghana

India

Kenya

Pakistan

Sudan

United Kingdom

**Study participating centre**

**University of Oxford**

Oxford

United Kingdom

OX3 7LF

## **Sponsor information**

**Organisation**

University of Oxford (UK)

**Sponsor details**

Clinical Trials and Research Governance Office

Manor House

John Radcliffe Hospital

Headington

Oxford

England

United Kingdom

OX3 9DZ

-

Heather.House@admin.ox.ac.uk

**Sponsor type**

University/education

**Website**

<http://www.ox.ac.uk/>

**ROR**

<https://ror.org/052gg0110>

## **Funder(s)**

**Funder type**

Research council

**Funder Name**

Medical Research Council (UK) (Grant ref: G0500959)

**Alternative Name(s)**

Medical Research Council (United Kingdom), UK Medical Research Council, MRC

**Funding Body Type**

Government organisation

**Funding Body Subtype**

National government

**Location**

United Kingdom

## Results and Publications

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Protocol article</a>	protocol	22/10/2007		Yes	No
<a href="#">Results article</a>	results	20/07/2013		Yes	No
<a href="#">Protocol article</a>	follow-up protocol	21/11/2013		Yes	No
<a href="#">Results article</a>	3-year follow-up results	02/07/2016		Yes	No