The preventive and therapeutic effects of electroacupuncture on blood pressure and heart rate reduction in carotid artery stenting

Submission date 24/02/2025	Recruitment status Recruiting	Prospectively registeredProtocol
Registration date	Overall study status	Statistical analysis plan
05/03/2025	Ongoing	☐ Results
Last Edited	Condition category	☐ Individual participant data
03/03/2025	Circulatory System	[X] Record updated in last year

Plain English summary of protocol

Background and study aims

Carotid sinus reflex (CSR) is a common complication during and after carotid artery stenting (CAS), clinically manifested as bradycardia (slow heart rate) and hypotension (low blood pressure), which poses significant risks to patient safety. This study aims to investigate the effects of electroacupuncture on carotid sinus reaction occurring during and after carotid artery stenting.

Who can participate?

Patients aged 18-75 years diagnosed with carotid artery stenosis requiring carotid artery stenting

What does the study involve?

Participants will receive electroacupuncture treatment. The treatment is designed to stabilize hemodynamic changes (heart rate and blood pressure) during and after the carotid artery stenting procedure. The study will compare the outcomes of patients who receive electroacupuncture with those who do not, to evaluate its effectiveness in preventing or reducing carotid sinus reaction.

What are the possible benefits and risks of participating? Possible benefits:

- 1. Participants may experience improved stabilization of heart rate and blood pressure during and after the procedure.
- 2. The study may contribute to a better understanding of how electroacupuncture can help manage carotid sinus reflex, potentially benefiting future patients with similar conditions. Possible risks:
- 1. Normal reactions during acupuncture: You may experience sensations such as soreness, numbness, heaviness, or swelling, which are normal reactions to acupuncture. Additionally, dizziness may occur due to your physical condition or emotional tension, but it can be alleviated by stopping the acupuncture and resting appropriately.
- 2. Adverse reactions after acupuncture: Bleeding or hematoma may occur, but these can be

resolved with local pressure. If an infection develops at the acupuncture site, your doctor will address it promptly.

Where is the study run from?
Beijing Hospital of Traditional Chinese Medicine (China)

When is the study starting and how long is it expected to run for? November 2024 to June 2026

Who is funding the study? Beijing Hospital of Traditional Chinese Medicine (China)

Who is the main contact?

- 1. Dr Shaosong Wang, wangssmail@163.com
- 2. Dr Chuanjin Song, songcjmail@163.com

Contact information

Type(s)

Public, Scientific

Contact name

Ms Chuanjin Song

ORCID ID

http://orcid.org/0009-0001-0976-7676

Contact details

Beijing Hospital of Traditional Chinese Medicine Beijing China 100010 +86 (0)17812003007 songcjmail@163.com

Type(s)

Principal Investigator

Contact name

Dr Shaosong Wang

Contact details

Beijing Hospital of Traditional Chinese Medicine Beijing China 100010 +86 (0)13269211249 wangssmail@163.com

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

Nil known

Study information

Scientific Title

Vagal regulation-mediated effects of electroacupuncture on prevention and treatment of carotid sinus reflex in patients undergoing carotid artery stenting

Study objectives

Electroacupuncture can prevent and treat carotid sinus reflex induced by carotid artery stenting

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 21/01/2025, Beijing Traditional Chinese Medical Hospital Ethics Committee (Beijing Hospital of Traditional Chinese Medicine, No. 23, Art Museum Back Street, Dongcheng District, Beijing, 100010, China; +86 (0)10 8790 6734; website@bjzhongyi.com), ref: 2024BL02-139-01

Study design

Multicenter interventional single-blinded randomized controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment, Safety, Efficacy

Participant information sheet

Not available in web format, please use the contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Prevention and treatment of carotid sinus reflex in patients undergoing carotid artery stenting

Interventions

The randomization sequence will be generated by a statistician not involved in the trial using the random number table method. The allocation sequence will be concealed in sequentially numbered, sealed, opaque (non-recoverable) envelopes to ensure sequence confidentiality.

Participants will be randomly allocated (1:1) to either the treatment group or control group. One group will receive a single continuous electroacupuncture treatment during the surgical procedure on the operation day. The other group will receive a single continuous non-acupoint shallow needling on the operation day. The treatment duration will span from the beginning to the end of the surgical procedure. Assessments will be conducted at baseline (pre-intervention), on the operation day, daily from surgery completion to discharge, and at 4 weeks post-discharge.

Intervention Type

Procedure/Surgery

Primary outcome measure

The incidence of carotid sinus reflex (CSR) incidence, calculated by comparing baseline blood pressure/heart rate with the lowest intraoperative and postoperative values recorded via continuous non-invasive monitoring and ECG during and after surgery

Secondary outcome measures

- 1. Incidence of persistent carotid sinus reflex (CSR), measured using continuous hemodynamic monitoring data and anesthesia records intraoperatively and postoperatively until hospital discharge
- 2. Vasoactive drug utilization rate and postoperative pharmacological rescue rate: the percentage of patients requiring vasopressors/inotropes measured using anesthesia medication records and postoperative nursing charts during the procedure and postoperatively until discharge
- 3. Adverse event incidence rate: the frequency of complications (e.g., arrhythmias, stroke) measured using medical records and investigator-reported events from procedure initiation until hospital discharge
- 4. Neurological deficit measured using the NIH Stroke Scale (NIHSS) score at preoperative baseline (1 day before surgery), 1 hour postoperatively, 24 hours postoperatively, on the day of discharge, and 4 weeks post-discharge
- 5. Length of hospital stay: days from admission to discharge measured using hospital medical records at the time of discharge

Overall study start date

01/11/2024

Completion date

01/06/2026

Eligibility

Key inclusion criteria

- 1. European Society for Vascular Surgery (ESVS) 2023-defined carotid stenosis
- 2. CAS-indicated per Chinese Guidelines 2015
- 3. Willingness for endovascular intervention
- 4. Aged 18-75 years inclusive
- 5. Signed informed consent

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Upper age limit

75 Years

Sex

Both

Target number of participants

128

Key exclusion criteria

- 1. Presence of cardiac pacemaker
- 2. Documented severe dysfunction in any major organ system (cardiac, pulmonary, hepatic, or renal)
- 3. Identified coagulation disorders
- 4. Active skin infection at proposed acupuncture sites
- 5. Indication for emergency surgical intervention or additional neurointerventional procedures
- 6. Diagnosed psychiatric conditions or documented non-compliance that may compromise treatment protocol adherence
- 7. Documented intolerance to electroacupuncture therapy or history of severe adverse reactions
- 8. Anatomically unsuitable vascular access confirmed by imaging studies

Date of first enrolment

22/01/2025

Date of final enrolment

01/05/2026

Locations

Countries of recruitment

China

Study participating centre Beijing Hospital of Traditional Chinese Medicine

Capital Medical University
Department of Neurology
No. 23, Art Museum Back Street
Dongcheng District
Beijing

Study participating centre Peking University First Hospital

No. 8 Xishiku Street, Xicheng District Beijing China 100034

Sponsor information

Organisation

Beijing Hospital of Traditional Chinese Medicine

Sponsor details

No. 23, Art Museum Back Street, Dongcheng District Beijing China 100010 +86 (0)10 52176677 website@bjzhongyi.com

Sponsor type

Hospital/treatment centre

Website

https://www.bjzhongyi.com/

ROR

https://ror.org/057vq6e26

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Beijing Hospital of Traditional Chinese Medicine

Results and Publications

Publication and dissemination plan

Planned publication in a high-impact peer-reviewed journal

Intention to publish date

06/01/2027

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study during this study will be included in the subsequent results publication

IPD sharing plan summary

Other