

Randomised, double-blinded placebo controlled trial of the use of tocolysis for a repeat external cephalic version after a failed version for breech presentation at term

Submission date 23/01/2004	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 23/01/2004	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 12/01/2010	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mr Lawrence Impey

Contact details
Level 4
The Women's Centre
John Radcliffe Hospital
Headley Way
Headington
Oxford
United Kingdom
OX3 9DU
+44 (0)1865741166 Ex 21408/21624
LWMI@compuserve.com

Additional identifiers

Protocol serial number
SEO093

Study information

Scientific Title

Study objectives

To determine the optimum protocol for performing external cephalic version for breech presentation at term. Null hypothesis: that tocolysis for a repeat external cephalic version (ECV) after a failed ECV does not increase the success rates of ECV

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Childbirth

Interventions

1. 50 mg ritodrine hydrochloride infusion
2. 17 ml dextrose saline solution infusion

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

1. Incidence of cephalic presentation at delivery
2. Subjective patient assessment of discomfort
3. Incidence of caesarian section
4. Length of hospital stay
5. Incidence of neonatal Apgar scores <7 at 5 minutes
6. Incidence of neonatal intensive care admission

Key secondary outcome(s)

Not provided at time of registration

Completion date

06/04/2002

Eligibility

Key inclusion criteria

1. Women with suspected or confirmed breech presentation at >36 weeks gestation
2. Have undergone a single (<15 mins) unsuccessful attempt at external cephalic version without tocolysis at 37 or more weeks' gestation

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

1. Abnormal cardiotocograph
2. Cardiac disease
3. Pre-existing or gestational diabetes
4. Hypertension;
5. Pre-existing indication for caesarian section
6. Suspected unstable lie
7. Pre-eclampsia
8. recent (<4 weeks) antepartum haemorrhage
9. suspected fetal compromise
10. rhesus immunisation

Date of first enrolment

06/04/2000

Date of final enrolment

06/04/2002

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Level 4
Oxford
United Kingdom
OX3 9DU

Sponsor information

Organisation
NHS R&D Regional Programme Register - Department of Health (UK)

Funder(s)

Funder type
Government

Funder Name
NHS Executive South East (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/05/2005		Yes	No