

# The effect of air or enriched air nitrox breathing during simulated diving on intravascular bubble formation following decompression

**Submission date**  
24/02/2016

**Recruitment status**  
No longer recruiting

Prospectively registered

Protocol

**Registration date**  
25/03/2016

**Overall study status**  
Completed

Statistical analysis plan

Results

**Last Edited**  
25/04/2023

**Condition category**  
Injury, Occupational Diseases, Poisoning

Individual participant data

## Plain English summary of protocol

### Background and study aims

Decompression sickness is an uncommon but serious complication of diving. It occurs when hyperbaric (high pressure) conditions during diving lead to bubbles of nitrogen forming in the blood vessels upon decompression (when the diver ascends from depth). Decompression sickness can produce many symptoms, ranging from joint pain and rashes to paralysis and death. Breathing gas mixtures with less nitrogen and higher oxygen content may decrease bubble formation after decompression. This study aims to investigate whether enriched air nitrox reduces bubble formation during simulated diving as compared to breathing normal air.

### Who can participate?

Healthy volunteers aged over 18 with a diving license and without a history of a decompression accident.

### What does the study involve?

Participants perform a simulated dive breathing air in the hyperbaric chamber of the CHU d'Angers. Intravascular bubble formation is assessed after the dive using cardiac (heart) ultrasound. Twelve participants prone to bubbling are then selected to perform two more simulated dives, one dive breathing air and the other dive breathing enriched air nitrox. Cardiac ultrasound is performed after each dive.

### What are the possible benefits and risks of participating?

There are no benefits for the volunteers. Possible side-effects include decompression symptoms, similar to a non-simulated dive. All volunteers will be accompanied by a physician certified in hyperbaric medicine during a simulated dive.

### Where is the study run from?

Centre Hospitalier Universitaire d'Angers (France)

### When is the study starting and how long is it expected to run for?

January 2001 to August 2007

Who is funding the study?  
Investigator initiated and funded (France)

Who is the main contact?  
Prof Pierre Asfar

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Pierre Asfar

**Contact details**  
Department of Medical Intensive Care and Hyperbaric Medicine  
Centre Hospitalier Universitaire d'Angers  
4 rue Larrey  
Angers  
France  
49933

## Additional identifiers

**Protocol serial number**  
N/A

## Study information

**Scientific Title**  
Comparison of bubble formation after diving with air or nitrox using cardiac ultrasound  
(Comparaison par échocardiographie Doppler du phénomène bulleux à l'issue d'une plongée à l'air et au nitrox)

**Acronym**  
NITROX

**Study objectives**  
Enriched air nitrox reduces venous bubble scores as compared to air breathing during diving, and thereby the risk of decompression sickness in volunteers selected for high post-decompression bubble formation.

**Ethics approval required**  
Old ethics approval format

**Ethics approval(s)**  
Committee for protection of human subjects in biomedical research of Pays de la Loire (Comité consultatif de protection des personnes dans la recherche biomédical des Pays de la Loire), 23/11 /2001, Protocol number 2001/17

**Study design**

Single-center prospective double-blind study with crossover design

**Primary study design**

Interventional

**Study type(s)**

Prevention

**Health condition(s) or problem(s) studied**

Decompression illness in hyperbaric medicine

**Interventions**

The 47 divers underwent an initial simulated air dive to exclude subjects with low intravascular bubble production (Bubble score  $\leq 1$ ) in order to select divers with high bubble production (Bubble score  $\geq 2$ ). Twelve divers (10 men and two women) demonstrating high intravascular bubble production completed the study protocol in a randomized, double-blinded crossover setup. Divers were assigned to undergo one simulated dive while breathing air (Air; 21% oxygen) and one simulated dive breathing enriched air nitrox (EAN) with 36% oxygen in a randomized order.

**Intervention Type**

Other

**Primary outcome(s)**

Pulsed Doppler measurements of the trunk of the pulmonary artery in order to quantify intravascular bubble formation. Measurements of at least 1 minute duration were performed by a certified cardiologist, blinded for the FiO<sub>2</sub> at 0, 30, 60 and 90 minutes after decompression. Images were recorded and analyzed for bubble scores offline. A modified bubble score was used, based on the Doppler score system of Spencer. Bubbles scores were analyzed independently by two blinded reviewers.

**Key secondary outcome(s)**

Decompression incidents by questionnaire 90 minutes after completion of the simulated dive.

**Completion date**

10/08/2007

**Eligibility****Key inclusion criteria**

1. Forty-seven human volunteers
2. Age >18 years
3. Diving experience as confirmed by possession of a French recreational diver license
4. Absence of contraindication to dive

**Participant type(s)**

Healthy volunteer

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Total final enrolment**

12

**Key exclusion criteria**

1. History of decompression accident(s)
2. Oxygen administration or diving within 24 hours before simulated dive
3. Absent or low intravascular bubble production (Bubble score  $\leq$  1) after initial simulated test dive

**Date of first enrolment**

03/07/2002

**Date of final enrolment**

10/08/2007

**Locations****Countries of recruitment**

France

**Study participating centre**

Centre Hospitalier Universitaire d'Angers

4 Rue Larrey

Angers

France

49933

**Sponsor information****Organisation**

Centre Hospitalier Universitaire d'Angers (France)

**ROR**

<https://ror.org/0250ngj72>

# Funder(s)

## Funder type

Other

## Funder Name

Investigator initiated and funded (France)

# Results and Publications

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request

## IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	10/05/2016		Yes	No
<a href="#">Protocol (other)</a>		10/05/2016	25/04/2023	No	No