To compare the pain relief from combined intrathecal bupivacaine-morphine with bupivacaine alone in lower limb surgery

Submission date	Recruitment status	Prospectively registered
25/04/2022	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
12/05/2022	Completed	Results
Last Edited	Condition category	Individual participant data
05/05/2022	Surgery	Record updated in last year

Plain English summary of protocol

Background and study aims

Inadequate pain medication during and after surgery may lead to long term pain after surgery.

This study was designed to ascertain the effect of morphine addition to bupivacaine in the management of postoperative pain in patients undergoing lower limb surgeries at the Tamale Teaching Hospital, Ghana.

Who can participate?

Patients aged 16 – 60 years, undergoing lower limb surgery.

What does the study involve?

Participants will be randomly allocated to receive bupivicane alone or bupivicane plus morphine during surgery. Follow up lasts 72 hours.

What are the possible benefits and risks of participating?

Benefits: There were no immediate benefits for participation in the study. No gifts or money were given in exchange for the information obtained. However, data generated from this study will help policy makers in the Ministry of Health to improve the management of postoperative pain after Orthopaedic and Trauma surgeries.

Risks: Participants were informed that they may experience slight pain during the sample collection process. Should they suffer any injury during the collection process, members of the research team will take full responsibility and cover all expenses pertaining to treatment.

Where is the study run from? Tamale Teaching Hospital, Ghana.

When is the study starting and how long is it expected to run for? April 2021 to November 2021

Who is funding the study? Investigator initiated and funded

Who is the main contact? Sylvanus Kampo, sylvanuskampo@yahoo.com Fidelis Bayor, bayorfidelis@gmail.com

Contact information

Type(s)

Scientific

Contact name

Mr Fidelis Bayor

ORCID ID

https://orcid.org/0000-0002-2703-8856

Contact details

University of Ghana Medical School Accra Ghana 24 +233 249229949 bayorfidelis@gmail.com

Type(s)

Principal investigator

Contact name

Dr Sylvanus Kampo

Contact details

CK Tedam University of Technology and Applied Sciences Navrongo Ghana 19 +233 559355229 sylvanuskampo@yahoo.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Analgesic efficiency of intrathecal bupivacaine with morphine and bupivacaine alone in lower limb orthopaedic surgery. A comparative study

Acronym

IBM

Study objectives

Bupivacaine with morphine induces better postoperative pain relief and improve patient outcomes in lower limb orthopaedic surgery

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 30/09/2021, Ethical review board of the Kwame Nkrumah University of Science and Technology (University Post Office, Kumasi, Ghana; +233(0)205453785; chrpe@knust.edu.gh), ref: CHRPE/AP/416/21

Study design

Randomised comparative controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Postoperative pain management in trauma patients

Interventions

After meeting the inclusion criteria, eligible patients received 15 mg preservative free hyperbaric bupivacaine and 0.2 mg morphine or 15 mg bupivacaine alone. Patients were followed up 72 hrs for postoperative complications and patient outcome. Each recruited patient was randomly assigned to one of two groups using a computer-generated random number table. Group A (n=30) represented those who received bupivacaine alone and

Group B (n=30) represented those who received bupivacaine with morphine.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

bupivacaine, morphine

Primary outcome(s)

- 1. Vital signs (pulse rate, blood pressure, oxygen saturation, and respiratory rate) of the patient were monitored and recorded for every 5 minutes for the first 30 minutes and then for every 15 minutes during surgery
- 2. Pain severity was measured every 4 hours for 72 hours immediately after the surgery on a 100-mm VAS, 0 mm = no pain, and 100 mm = intolerable pain.

Key secondary outcome(s))

- 1. The episodes of postoperative nausea and vomiting (PONV) were identified by direct scheduled assessments or by a spontaneous complaint by the patients after the surgery. The incidence of PONV was recorded hourly for the first 4 hours and then 4 hourly for the next 24 hours using a 3 point ordinal scale (0 = none, 1 = nausea, 2 = vomiting). The incidence of PONV was calculated and categorized as early (0 4 hours) or delayed (5 24 hours).
- 2. The incidence of pruritus was recorded every 4 hours for 48 hours after surgery on a four-point categorical scale as; 0 = no pruritus, 1 = mild, 2 = moderate, 3 = severe pruritus.
- 3. Overall perioperative satisfaction was evaluated on the day of discharged during an interview as; 4 = excellent, 3 = good, 2 = satisfactory, 1 = poor.

Completion date

30/11/2021

Eligibility

Key inclusion criteria

- 1. Patients undergoing elective lower limb orthopaedic surgery
- 2. ASA status class I and II
- 3. Patients between 16 60 years of age
- 4. Patients who were on admission for 72 hours or more postoperatively

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

60

Key exclusion criteria

- 1. Pregnant or breastfeeding mothers
- 2. Patients with polytrauma
- 3. Glasgow coma scale <15
- 4. Opioids abusers
- 5. Expected post-surgical stay <24 hours

- 6. ASA physical status class III and IV
- 7. Patients allergic to bupivacaine and/morphine
- 8. Contraindication to spinal anaesthesia

Date of first enrolment

10/09/2021

Date of final enrolment

29/11/2021

Locations

Countries of recruitment

Ghana

Study participating centre Tamale Teaching Hospital

Box 1350TL Tamale Ghana

_

Sponsor information

Organisation

Platinum Orthopaedic and Spine Clinic

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

Availability of data and materials: The datasets generated and/or analysed during the current study are not publicly available due to patient confidentiality but are available from the

corresponding author on reasonable request. sylvanuskampo@yahoo.com

IPD sharing plan summary

Available on request

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet
Participant information sheet
11/11/2025 No Yes