Cognitive behaviour therapy for epilepsy: improving seizure control and quality of life

Submission date	Recruitment status No longer recruiting	Prospectively registered		
26/11/2008		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
10/12/2008		[X] Results		
Last Edited	Condition category	Individual participant data		
07/08/2020	Nervous System Diseases			

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Laura H Goldstein

Contact details

Department of Psychology Institute of Psychiatry King's College London PO77 De Crespigny Park London United Kingdom SE5 8AF

Additional identifiers

Protocol serial number

1.0

Study information

Scientific Title

Cognitive behaviour therapy for epilepsy: improving seizure control and quality of life - a pilot randomised controlled trial

Study objectives

This is a pilot/feasibility study investigating the use of cognitive behaviour therapy (CBT) plus standard medical care vs standard medical care alone in improving seizure control and quality of life in adults with refractory epilepsy. Specifically this feasibility study will enable us to:

- 1. Evaluate the applicability of our CBT techniques to a broader sample of patients than in our previous work
- 2. Determine the acceptability of this randomisation and treatment approach in a sample of patients attending specialist epilepsy clinics, thereby informing recruitment rates when designing a later, larger randomised controlled trial (RCT)
- 3. Obtain information on treatment effect sizes to undertake power calculations for a larger RCT

Ethics approval required

Old ethics approval format

Ethics approval(s)

Joint South London and Maudsley and the Institute of Psychiatry NHS Research Ethics Committee, 04/07/2008, ref: LREC 08/H0807/44

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Epilepsy

Interventions

Following baseline recording of seizure frequency participants will be randomly assigned to one of two arms:

CBT (plus standard medical care):

CBT will comprise 12 weekly/fortnightly individual therapy sessions with the CBT therapist, over 4 months. Treatment will follow our prepared treatment manual, with homework tasks and seizure recordings reviewed at each session. In addition to seizure-specific interventions (e.g. development of countermeasures), more general cognitive behavioural therapeutic techniques will be employed to address the management of times of increased vulnerability to seizures and to improve patients overall physical and emotional well-being, with a view to relapse prevention. Handouts will be given to participants.

Standard medical care:

Participants will be seen by their epilepsy specialist in their regular clinic, depending on clinical need. As with the CBT group, we will monitor the number of appointments that participants have with their epilepsy specialist over the 4-month period and whether any medication changes are implemented during that time.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Monthly seizure frequency, monitored throughout the study. Total duration of follow-up: 12 weeks.

Key secondary outcome(s))

- 1. Liverpool Seizure Severity Scale at baseline, 8 and 12 weeks
- 2. Hospital Anxiety and Depression Scale at baseline, 8 and 12 weeks
- 3. Quality of Life in Epilepsy Inventory-31 (QOLIE-31) at baseline, 8 and 12 weeks
- 4. Health service usage for previous 8 weeks, assessed using the Client Service Receipt Inventory (CSRI) at baseline, beginning of treatment, 8-week follow-up and for previous 4 weeks at 12 week follow up
- 5. Health status, assessed by Euroqol EQ-5D at baseline, 8 and 12 weeks

Completion date

31/03/2010

Eligibility

Key inclusion criteria

- 1. Both males and females, age 18-65 years
- 2. Clinical diagnosis (with supporting electroencephalography [EEG]) of epilepsy
- 3. A maximum of 40 seizures/month involving alteration/loss of consciousness (i.e. complex partial or generalised seizures) with no limit to simple partial seizure frequency
- 4. Stable medication for the month prior to recruitment
- 5. Able to attend weekly/fortnightly sessions
- 6. Willing to complete questionnaires regularly to monitor progress

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

65 years

Sex

All

Total final enrolment

Key exclusion criteria

- 1. History of, or current actual/suspected non-epileptic seizures
- 2. <2 seizures per month in each of the preceding four months
- 3. Active major psychiatric disorder
- 4. Drug/alcohol dependence
- 5. Receiving active vagus nerve stimulation (VNS) for seizures that has been adjusted within the previous 12 months
- 6. Insufficiently fluent in English to be able to undertake treatment and complete questionnaires without the assistance of an interpreter
- 7. Established learning disability (or where this has been measured formally, IQ <70)

Date of first enrolment

01/12/2008

Date of final enrolment

31/03/2010

Locations

Countries of recruitment

United Kingdom

England

Study participating centre Department of Psychology

London United Kingdom SE5 8AF

Sponsor information

Organisation

Institute of Psychiatry, King's College London (UK)

ROR

https://ror.org/0220mzb33

Funder(s)

Funder type

Charity

Funder Name

Epilepsy Research UK (UK) (ref: P0805)

Alternative Name(s)

Epilepsy Research UK, The Epilepsy Research Institute, ERUK

Funding Body Type

Private sector organisation

Funding Body Subtype

Other non-profit organizations

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Basic results		07/07/2017	07/08/2020	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes